

**Integrated Management Plan  
of  
East Kolkata Wetlands**  
(Management Action Plan 2021-2026)



**Department of Environment  
Government of West Bengal**





## **East Kolkata Wetlands Management Authority**

The East Kolkata Wetlands (Conservation and Management) Act, 2006 represents an important landmark as it paved way for establishment of the East Kolkata Wetlands Management Authority (EKWMA) for conservation and management of the East Kolkata Wetlands (EKW) and for matters connected therewith and incidental thereto. The EKWMA was constituted under Section 3(1) of the EKW(C&M) Act, 2006.

As per the East Kolkata Wetlands (Conservation and Management) (Amendment) Act, 2017 notified by the State Government vide no. 304-L dated 17.03.2017, the EKWMA has been reconstituted. According to the said Amendment Act, 2017, the EKWMA is now a thirteen (13) member body headed by the Hon'ble Minister-in Charge, Department of Environment, Government of West Bengal along with the Chief Secretary and Secretaries of different Departments of State Government as well as four experts each in the areas of wetland ecology, hydrology, fisheries and socio-economics.

The EKWMA has been entrusted with wide range of responsibilities including demarcation of wetland boundary, prevention of unauthorised land use of EKW, pollution abatement and conservation of flora and fauna, awareness generation about the utility of the wetland, to promote basic conservation principles of sewage fed pisciculture and research on various aspects of EKW.



## **Wetlands International South Asia**

Wetlands International South Asia is a non-government organisation working for sustaining and restoring wetlands, their resources and biodiversity in the South Asia region. Its office in New Delhi (India) was established in 1996 as a part of Wetlands International network. Wetlands International is a global, independent, non-profit organisation dedicated to conservation and restoration of wetlands, and presently works in over 100 countries through a network of 18 regional and national offices and expert networks headquartered in The Netherlands. Wetlands International is also one of the five International Organization Partners of the Ramsar Convention. In 2005, Wetlands International South Asia was registered under the Societies Registration Act of Government of India (retaining remit of South Asia region), consequently gaining an Indian legal entity while subscribing to the goals and targets of the Wetlands International network. The organisation endeavours to use a mix of approaches including technical knowledge, policy dialogue and field demonstrations for addressing various issues related to wetland management. To leverage change, the organisation works with national and state governments, knowledge centres, civil society as well as the private sector, often acting as catalysts to enable joined up actions. Given that securing a positive change in the status of wetlands and linked livelihoods takes considerable time, the organisation works for long-term engagement, forging strategic and innovative partnerships.

# Integrated Management Plan of East Kolkata Wetlands (Management Action Plan 2021 – 2026)



2021

This plan was drafted under the supervision of the Technical Committee, State Wetlands Authority, West Bengal comprising:

1. Dr. Kalyan Rudra, Chairman, West Bengal Pollution Control Board
2. Sri. Niraj Singhal, Chief Environment Officer, Environment Department (from 24.08.2018 to 26.02.2021)
3. Sri. Kaliyamurthi Balamurugan, Chief Environment Officer, Environment Department (from 26.02.2021 to till date)
4. Smt. Esha Sengupta, Director of Fisheries, Fisheries Department
5. Sri. Sampad Ranjan Patra, Director of Agriculture, Agriculture Department
6. Prof. Arunabha Majumder, Expert Member Ecology, State Wetlands Authority
7. Dr. Pratap Kumar Mukhopadhyay, Expert Member Fisheries, State Wetlands Authority
8. Prof. Basab Chaudhuri, Expert Member Socio-economics, State Wetlands Authority

In the preparation of the Integrated Management Plan of East Kolkata Wetlands (Management Action Plan 2021-2026), the valuable contribution of the following experts is acknowledged:

Dr Ritesh Kumar, Director, Wetlands International South Asia  
Mr Harsh Ganapathi, Technical Officer, Wetlands International South Asia  
Mr Dushyant Mohil, Project Manager, Partners for Resilience, Wetlands International South Asia  
Mr Kamal Dalakoti, Junior Technical Officer, Wetlands International South Asia

**Report ID:** EKW-2021-01

**Suggested Citation:** EKWMA and WISA (2021). Integrated Management Plan of East Kolkata Wetlands (Management Action Plan 2021-2026). East Kolkata Wetlands Management Authority and Wetlands International South Asia.

**Disclaimer:** Usual disclaimers apply.

**Photograph credits:** EKWMA and WISA Photo Library unless cited

**Cover:** Wetland features of East Kolkata Wetlands

**Back Cover:** Water Hyacinth used as biological screens at East Kolkata Wetlands

**First Edition:** February, 2021

**Second Revised Edition:** August, 2021

**Prof. (Dr.) Saumen Kumar Mahapatra**



**MINISTER-IN-CHARGE**  
**Department of Environment**  
Government of West Bengal  
Pranisampad Bhawan, 5th Floor, LB-2 Block,  
Salt Lake, Kolkata-700 106  
Tel. ☎ (033) 2335-0238, Fax : (033) 2335-2329  
&  
**Department of Public Health Engineering**  
Government of West Bengal  
New Secretariat Building, 7th Floor,  
1, K. S. Roy Road, Kolkata-700 001,  
Tel. ☎ (033) 2248-2130, Fax : (033) 2248-9843

### **MESSAGE**

The East Kolkata Wetlands are situated on the eastern fringes of Kolkata city. The wetlands naturally treat sewage generated from the city, through the shallow pisciculture ponds, where the sewage is fed on regulated basis. Besides the production of fish and vegetables, the wetlands act as a Carbon sink, help in flood buffering and climate regulation of the region, harbour rich biodiversity and provide many more environmental benefits. The wetlands save substantial capital and recurring costs involved in setting up of conventional Sewage Treatment Plants (STPs).

Water is life, and wetlands are the life support systems that ensure optimal functioning of the water cycle. Despite their tremendous value, wetlands are getting degraded globally due to natural and anthropogenic impacts. Considering the numerous challenges to the East Kolkata Wetlands, it was felt necessary to formulate an integrated Management Plan of the East Kolkata Wetlands for their conservation and sustainable development, ensuring livelihood opportunities for the wetlands communities.

The East Kolkata Wetlands Management Action Plan 2021-2026 comprises four basic components viz. Institutions and Governance, Water Management and Pollution Abatement, Conservation of Species and Habitats, and Sustainable Resource Development. The Plan is a major step forward towards the maintenance of East Kolkata Wetlands in a healthy condition to enable delivery of their full range of ecosystem services and sustaining biological diversity.

I would like to thank all officials and staff of the Environment Department and the East Kolkata Wetlands Management Authority for their contribution in preparing the East Kolkata Wetlands Management Action Plan 2021-2026.

Prof. (Dr.) Saumen Kumar Mahapatra  
Minister-in-Charge, Environment Department  
and  
Chairman, East Kolkata Wetlands Management Authority



## Summary

---

Wetland Name	East Kolkata Wetlands
Wetland Area (in ha)	12500 ha
Location	South 24 Parganas and North 24 Parganas
Area of the direct catchment (in ha)	162500 ha
Name of the nodal agency for management plan implementation	East Kolkata Wetlands Management Authority, Department of Environment, Government of West Bengal
Management plan period	2021-2026
Date on which approval of East Kolkata Wetlands Management Authority was obtained	25.01.2021
Total budget	Rs. 110.7198 Crores
Funds requested from State Government	Rs. 44.2898 Crores
Funds requested from MoEFCC	Rs. 66.43 Crores





## Contents

<b>Message: Chairman, East Kolkata Wetlands Management Authority</b>	<b>1</b>
<b>Summary</b>	<b>3</b>
<b>List of Figures</b>	<b>8</b>
<b>List of Maps</b>	<b>9</b>
<b>List of Tables</b>	<b>10</b>
<b>List of Pictures</b>	<b>11</b>
<b>Acronyms</b>	<b>12</b>
<b>Executive Summary</b>	<b>15</b>
<b>1 Introduction</b>	<b>21</b>
<b>1.1 East Kolkata Wetlands</b>	<b>21</b>
<b>1.2 Management Planning Purpose</b>	<b>21</b>
<b>1.3 Management to-date: An overview</b>	<b>22</b>
<b>1.4 Approach and Methodology</b>	<b>26</b>
<b>1.5 Management Plan Structure</b>	<b>28</b>
<b>2 Description of Wetland Features</b>	<b>29</b>
<b>2.1 Physical regime</b>	<b>29</b>
<b>2.2 Wetland catchments</b>	<b>37</b>
<b>2.3 Hydrological regimes</b>	<b>46</b>
<b>2.4 Species and Habitat</b>	<b>62</b>
<b>2.5 Livelihoods and Resource Linkages</b>	<b>67</b>
<b>3 Evaluation of Wetland Features</b>	<b>84</b>
<b>3.1 Ecological Character Description</b>	<b>85</b>
<b>3.2 Status and trends</b>	<b>87</b>
<b>3.3 Risk of adverse change in ecological character</b>	<b>93</b>
<b>3.4 Key knowledge gaps</b>	<b>95</b>
<b>4. Institutional Arrangements</b>	<b>96</b>
<b>4.1 Existing Institutional and Governance Arrangements</b>	<b>96</b>
<b>4.2 Gaps</b>	<b>101</b>
<b>4.3 Proposed Arrangements</b>	<b>104</b>

<b>5 Management Framework</b>	<b>105</b>
<b>5.1 Management Goal and Purpose</b>	<b>105</b>
<b>5.2 Management Strategy</b>	<b>106</b>
<b>5.3 Management Objectives and Performance indicators</b>	<b>110</b>
<b>5.4 Risks and Risk Mitigation Options</b>	<b>114</b>
<b>6 Monitoring Plan</b>	<b>116</b>
<b>6.1 Monitoring Objective</b>	<b>116</b>
<b>6.2 Monitoring Strategy</b>	<b>117</b>
<b>6.3 Assessing management effectiveness</b>	<b>126</b>
<b>6.4 Infrastructure and human resources requirements</b>	<b>127</b>
<b>6.5 Reporting</b>	<b>127</b>
<b>6.6 Quality control</b>	<b>128</b>
<b>6.7 Review and adaptation</b>	<b>128</b>
<b>7 Action Plan</b>	<b>129</b>
<b>Component 1. Institutions and Governance</b>	<b>131</b>
<b>Component 2. Water Management and Pollution Abatement</b>	<b>138</b>
<b>Component 3. Conservation of Species and Habitats</b>	<b>141</b>
<b>Component 4. Sustainable Resource Development</b>	<b>144</b>
<b>8 Budget and Financing</b>	<b>146</b>
<b>8.1 Budget</b>	<b>146</b>
<b>8.2 Financing</b>	<b>152</b>
<b>8.3 Phasing of Activities</b>	<b>158</b>
<b>8.4 Financing Arrangements</b>	<b>162</b>
<b>References</b>	<b>163</b>
<b>Annexes</b>	
<b>A1 Brief Document (as per Wetlands (Conservation and Management) Rules, 2017)</b>	<b>167</b>
<b>A2 Wetland Health Card</b>	<b>182</b>
<b>A3 List of Mouzas</b>	<b>185</b>

<b>A4 List of species</b>	<b>186</b>
1. List of Protozoa: Free-living Ciliates	186
2. List of Protozoa: Free-living Testate Amoebae	187
3. List of Rotifers	189
4. List of Soil Nematodes	190
5. List of Soil Mites	192
6. List of Spiders	194
7. List of Crustacea: Cladocerans	195
8. List of Crustacea: Ostracods	196
9. List of Crustacea: Copepods	196
10. List of Crustacea: Crabs and Shrimps	197
11. List of Apterygotan fauna	200
12. List of Odonata: Dragonflies and Damselflies	201
13. List of Orthoptera: Grasshoppers and Crickets	202
14. List of Blattodea: Isoptera- Termites	206
15. List of Terrestrial Bugs (Hemiptera)	207
16. List of Aquatic and Semi-Aquatic (Heteroptera)	209
17. List of Ants	211
18. List of Hymenoptera: Vespidae	213
19. List of Chalcidids	213
20. List of Encyrtidae	213
21. List of Coleopterans: Beetles	214
22. List of Butterflies	215
23. List of Moths	217
24. List of Dipterans: True Flies	224
25. List of Gastropods and bivalves	231
26. List of Fish	233
27. List of Amphibians and Reptiles	238
28. List of Birds	240
29. List of Mammals	244
30. List of Fungi	245
31. List of Freshwater Algae	248
32. List of Bryophytes	252
33. List of Agro-Flora (Cultivation crops)	252
34. List of Floristic Diversity	254
35. List of Invasive Alien Species (Vascular Plants)	261
<b>A5 Wetland Mitra</b>	<b>264</b>
<b>A6 Schedule of Rates</b>	<b>266</b>
1. Boundary Pillar	266
2. Dredging and desilting	270

## List of Figures

---

Figure 1. Financial flows of East Kolkata Wetlands Management Authority (2006-2019)	25
Figure 2. Step-wise wetland management planning process	27
Figure 3. Land use and Land Cover transformation in East Kolkata Wetlands (2019)	29
Figure 4. Schematic diagram of East Kolkata Wetlands resource recovery system	31
Figure 5. Changes in area under different land use and land cover categories in East Kolkata Wetlands (2000-2019)	33
Figure 6. Land use transformation in East Kolkata Wetlands (2000-2019)	36
Figure 7. Precipitation trend in East Kolkata Wetlands direct catchment (1989 - 2020) (Stackhouse, 2020)	45
Figure 8. Temperature and evapotranspiration trend in East Kolkata Wetlands Basin (1989 – 2020) estimated using Thornthwaite’s equation (Stackhouse, 2020)	46
Figure 9. Inflows in East Kolkata Wetlands (Modelled)	49
Figure 10. BOD at various stations in East Kolkata Wetlands (2018-20)	57
Figure 11. COD in various stations of East Kolkata Wetlands (2018-2020)	58
Figure 12. Dissolved Oxygen levels in various stations of East Kolkata Wetlands (2018-2019)	58
Figure 13. Total Nitrogen levels in various stations of East Kolkata Wetlands (2018-2019)	59
Figure 14. Nitrate levels in various stations of East Kolkata Wetlands (2018-2019)	59
Figure 15. Phosphate levels at various stations of East Kolkata Wetlands (2018-2019)	60
Figure 16. Ammonium levels at various stations in East Kolkata Wetlands (2018-2019)	60
Figure 17. Distribution of noteworthy plant species found in East Kolkata Wetlands	64
Figure 18. Bird count and species reported from Nalban Bheri	65
Figure 19. Population Density and Growth in East Kolkata Wetlands	68
Figure 20. Education level of communities living in East Kolkata Wetlands	71
Figure 21. Possession of assets by communities living in East Kolkata Wetlands	71
Figure 22. Community’s perception score (6 to -6) of wetlands condition in the past decade	81
Figure 23. Community’s perception score (6 to -6) of threat to wetlands	82
Figure 24. Stakeholder responses on agencies responsible for implementing various interventions in East Kolkata Wetlands	83
Figure 25. Perception ranking of responsibility for implementing wetland management measures	83
Figure 26. Major risks assessed in East Kolkata Wetlands	93
Figure 27. Embedding the concept of BwN in EKW management plan	106
Figure 28. The Adaptive Management Cycle (after Allen et al, 2011)	109
Figure 29. Wise use of wetlands for the maintenance of ecological character at East Kolkata Wetlands	110
Figure 30. Management interventions proposed at East Kolkata Wetlands	130

## List of Maps

---

Map 1. East Kolkata Wetlands	24
Map 2. Seasonal transformation in land use and land cover of East Kolkata Wetlands (2019)	32
Map 3. Land use and land cover of East Kolkata Wetlands (November 2000 and November 2019)	34
Map 4. Wetlands of Southern Fraction of Ganga-Brahmaputra-Meghna Delta	39
Map 5. Catchment of East Kolkata Wetlands	40
Map 6. Direct catchment of East Kolkata Wetlands	41
Map 7. Aquifer system in East Kolkata Wetlands Catchment (India-WRIS 2020)	43
Map 8. Elevation profile of East Kolkata Wetlands catchment	44
Map 9. Sewage treatment systems in KMC Area	47
Map 10. Hydraulic structures in East Kolkata Wetlands Basin	48
Map 11. Canal system of East Kolkata Wetlands	50
Map 12. Groundwater levels in East Kolkata Wetland	53
Map 13. Groundwater levels and flow direction in East Kolkata Wetlands	55
Map 14. Water quality sampling stations in East Kolkata Wetland	61
Map 15. Mouza within East Kolkata Wetlands	69
Map 16. Proposed monitoring stations for water flow, water quality, air quality and biodiversity	119
Map 17. Proposed management zones of East Kolkata Wetlands	133
Map 18. Canals proposed to be desilted in East Kolkata Wetlands	139

## List of Tables

---

Table 1. Perception ranking of responsibility for implementing wetland management measures	70
Table 2. Concentration of heavy metals (mg/kg) for select fish samples	76
Table 3. Heavy metal concentrations in select vegetable samples of East Kolkata Wetlands	79
Table 4. Ecological Character Description	87
Table 5. Threats and associated impacts on East Kolkata Wetlands	94
Table 6. Staff strength in East Kolkata Wetlands Management Authority	97
Table 7. SWOT Analysis of current institutional arrangements for EKW management	102
Table 8. Benefits offered by natural infrastructure within the BwN concept	107
Table 9. Management objectives, performance indicators and targets	110
Table 10. Mapping of management objectives against wetland values and threats	113
Table 11. Risks and risk mitigation measures	114
Table 12. Inventory, assessment and monitoring needs for managing East Kolkata Wetlands	120
Table 13. Monitoring and assessment parameters and indicators	122
Table 14. Management plan components	129
Table 15. Canals to be desilted in East Kolkata Wetlands	138
Table 16. Summary of budget	146
Table 17. Analysis of convergence funding	147
Table 18. Detailed activity wise budget	148
Table 19. Yearwise breakup of requirement of funds required from NPCA	152
Table 20. Yearwise breakup of requirement of funds required from Government of West Bengal	154
Table 21. Phasing of activities	158

## List of Pictures

---

Picture 1. East Kolkata Wetlands provide immense ecological subsidies to the City of Kolkata	22
Picture 2. Part of Dhapa landfill in East Kolkata Wetlands which has been treated	30
Picture 3. Fishing activities in East Kolkata Wetlands	35
Picture 4. The Storm Water Flow Channel carries the monsoon runoff directly to Kulti River	38
Picture 5. The Dry Weather Flow Channel	51
Picture 6. The Bantala regulator from where sewage is diverted to the wetland	52
Picture 7. Diverse habitats in East Kolkata Wetlands	63
Picture 8. Black-winged Stilt in East Kolkata Wetlands (Credit: Kaushik Mukhopadhyay)	65
Picture 9. A Marsh mongoose in East Kolkata Wetlands	66
Picture 10. Life in East Kolkata Wetlands is closely linked to wetland health	68
Picture 11. Sewage fed fish farming has evolved traditionally in East Kolkata Wetlands	72
Picture 12. A bamboo screen filters solid waste from sewage before being brought into the fish farm	73
Picture 13. Nutrient in sewage is directed to vegetation growth and fish	74
Picture 14. Agriculture forms the terminal use of sewage in East Kolkata Wetlands	77
Picture 15. Vegetable cultivation in Dhapa region	78
Picture 16. Community participation during survey of EKW	80
Picture 17. Demolition of illegal structures in East Kolkata Wetlands is a major task of the Authority	98

## Acronyms

---

AMSL	Above Mean Sea Level
BMC	Bidhannagar Municipal Corporation
BOD	Biochemical Oxygen Demand
BSI	Botanical Survey of India
BwN	Building with Nature
CAG	Community Action Groups
CBO	Community Based Organisations
CGS	Central Government Scheme
CIFRI	Central Inland Fisheries Research Institute
CO <sub>2</sub>	Carbon Dioxide
COD	Chemical Oxygen Demand
COVID	Corona Virus Disease
CPCB	Central Pollution Control Board
CSO	Civil Society Organizations
DO	Dissolved Oxygen
DWF	Dry Weather Flow
EC	Electrical Conductivity
ECD	Ecological Character Description
EKW	East Kolkata Wetlands
EKWMA	East Kolkata Wetlands Management Authority
EKW Act	East Kolkata Wetlands (Conservation and Management) Act, 2006
FC	Faecal Coliform
FFC	Fishery Feed Canal
FIR	First Information Report
GHG	Greenhouse Gas
GP	Gram Panchayat
GPS	Global Positioning System
IESWM	Institute of Environmental Studies and Wetland Management
IMC	Indian Major Carps
IMP	Integrated Management Plan
IWMED	Institute of Wetland Management and Ecological Design
KLC	Kolkata Leather Complex



KMC	Kolkata Municipal Corporation
MCM	Million Cubic Metres
MLD	Million Litres per Day
MoEFCC	Ministry of Environment, Forest and Climate Change
MPN	Most Probable Number
MT	Metric Tonnes
NGO	Non-Government Organizations
NOC	No Objection Certificate
NPCA	National Plan for Conservation of Aquatic Ecosystems
NWC	National Wetlands Committee
pH	Potential of Hydrogen
PWD	Public Works Department
R-METT	Ramsar Site Management Effectiveness Tracking Tool
RSIS	Ramsar Sites Information Service
SHG	Self Help Group
SPM	Suspended Particulate Matter
SWF	Storm Weather Flow
SWOT	Strength, Weakness, Opportunity, and Threat
TC	Total Coliform
TON	Total Organic Nitrogen
TS	Total Solids
TSS	Total Suspended Solids
WBBD	West Bengal Biodiversity Board
WBPCB	West Bengal Pollution Control Board
WIAMS	Wetland Inventory, Assessment and Monitoring System
WISA	Wetlands International South Asia
WRR	Waste Recycling Region
ZSI	Zoological Survey of India



## Executive Summary

---

Located to the eastern fringes of Kolkata City and spanning 12,500 ha, East Kolkata Wetlands (EKW) is a mosaic of landforms including predominantly water dominated areas (used as fish farms) to land centric usages for agriculture, horticulture and settlements. The existing wetland regime is a remnant of series of brackish wetlands connected to the freshwater as well as marine environments of the Gangetic Delta and the Bay of Bengal, in an ecological continuum with the Sundarbans. Over 260 shallow fish ponds in the EKW receive over 900 MLD pre-settled sewage from the Kolkata Metropolitan region through a network of locally excavated secondary and tertiary canals, which is used to produce annually 20,000 MT of fish, 50,000 MT of vegetables and irrigate 2850 ha of paddy lands. As the nutrient-rich effluent moves through the system, it is progressively cleaned, and nutrients are redirected to the growth of algae or agricultural products grown along the pond edges and agricultural lands. Algae and other aquatic plants are used to feed up to 17 species of fish cultured in these ponds, which in turn create nitrogen and phosphorus-rich water to irrigate the adjacent rice fields. The traditionally evolved natural water purification waste recovery practice saves the Kolkata City nearly Rs. 4,680 million annually in terms of the treatment cost of up to 65% of the City's sewage. These wetlands also lock in over 60% of carbon from wastewater, thus reducing harmful Green House Gas emissions from the region.

The wetland is inhabited by diverse species. Atleast 380 species under major flora including 93 plant families, 10 amphibians, 29 reptiles, 123 birds, 79 fish, 24 crustaceans, and 13 mammal species have been recorded from these wetlands. Marsh mongoose *Herpestes auro punctatus* is endemic to the region and also included in the schedule II of Indian Wildlife Protection Act, 1972. The traditional waste recovery practice provides subsistence opportunities for a large, economically underprivileged population of 0.15 million living in over 37 mouzas within its boundaries. EKW is also one of the few natural habitats providing recreational avenues for the urban and peri-urban population.

East Kolkata Wetlands Management Authority is the nodal government agency mandated to ensure wise use of the wetland within the regulatory framework defined by the Wetlands (Conservation and Management) Rules, 2017 (notified under the Environment (Protection) Act, 1986) and the East Kolkata Wetlands (Conservation and Management) Act, 2006 and rules.

The ability of EKW to provide ecosystem services and sustain biological diversity is enabled by:

- a) Ingenious use of pre-settled sewage received from Kolkata City for resource recovery through aquaculture, horticulture and irrigation.
- b) Hydrological gradient which enables distribution of sewage to different parts of the wetland.
- c) Water-hyacinth mediated phytoremediation enabling wastewater treatment.
- d) Indigenous knowledge of the fish-farmers on management of fish ponds for cultivation of diverse species of fish.
- e) Diverse habitats which enables a range of plant and animal species to survive.
- f) Land use control as set under the provisions of East Kolkata Wetlands (Conservation and Management) Act, 2006 and Wetlands (Conservation and Management) Rules, 2017.

India, as a signatory of the Ramsar Convention, is committed to achieving wise use of all wetlands in her territory. Wise use of wetlands is defined in the text of Ramsar Convention as „the maintenance of their ecological character, achieved through the implementation of ecosystem approaches, within the context of sustainable development“. The evaluation of various wetland features has indicated following trends:

**Adverse land use change:** During 2000 – 2019, the area under fish farms, settlements and landfill has been observed to increase, while the area under horticulture and agriculture declined. The increase in area under fish farm area is largely on account of transformation from agriculture (more prominent after 2015). There is a conspicuous pressure on the conversion of fish farms in areas adjoining EM bypass.

**Waste treatment function of the wetland being compromised:** An increase in the presence of heavy metals along with the organic nutrients in the supply sewerage has been observed predominantly due to mixing of Kolkata city's industrial effluents. Wetland function of treating raw sewage can get affected by settling of heavy metals in the fish farm beds and also lead to bioaccumulation of heavy metals in fish and plant species. Presence of heavy metal contaminants in wetlands pose health risk to the producers and consumers dependent on resource harvest from wetlands.

**Disturbed habitats:** There is a reduction reported in the number of waterbird species being sighted in recent times. Increased urbanization in the periphery of wetland and shifting climate patterns has led to the reduction of key wetland species sighted in the area.

**Invasive fish species:** The rapid spread and population increase of suckermouth armoured catfishes in EKW in recent times is of concern, because of the notable possibility that these non-native catfishes may be adversely affecting fish germplasm and commercial fishery of this unique ecosystem.

**Increasing population pressure:** A rapid rise in population has been recorded in 2001-2011 which has grown as much as 4 times in some mouzas with population density, ranging from 100 to 4500 persons per square kilometre. This increased numbers have led to a greater demand for the wetland resources, particularly fish and food crops. The number of people dependent on fisheries has grown from 12000 to 20000 in the last two decades.

**Increasing climate risks:** The intensity of rainfall has increased while the duration has decreased meaning variations in freshwater flows. Wetland communities are highly vulnerable to impacts of climate change, including the risk of high floods and increase in temperature due to varying climate. The area around Salt Lake and Bantala is reported to have high rates of subsidence which can alter the topography and natural hydraulics of the wetland.

**Solid waste challenge:** The garbage dumping site at Mollar bheri within EKW has been stopped completely and fencing has been done along the water body to prevent spillage of legacy waste material into the water. Bio-mining of legacy waste has also been initiated at the site. Sustained efforts will be needed in this regard to protect EKW.

EKW needs to be managed for conservation of its natural remediation and resource recovery ability, biological diversity as well as securing sustained provision of its full range of ecosystem services which support livelihoods of dependent communities. The effectiveness of management will be reflected in the ability to sustain multiple use of the wetland, based on the traditional knowledge of resource recovery developed over time, without undermining the key ecological and social processes that underpin the functioning of EKW social-ecological system.

Management also needs to be dynamic and adaptive so as to accommodate uncertainties and challenges that emerge from multiple drivers and pressures, and allow for suitable modification based on continuous site monitoring and amalgamation of new information.

### Management Framework

The goal of management planning is to „maintain East Kolkata Wetlands in a healthy condition to enable delivery of its full range of ecosystem services and sustain biological diversity values’.

The purpose of management is to ‘enable natural infrastructure services to the Kolkata city in the form of wastewater treatment, flood buffering and climate regulation; providing livelihood opportunities for wetlands communities in food production; and sustain diversity of biota in the landscape’.

Management of EKW in 2021-2026 is structured around following eight objectives:

Objectives	Desired outcome
<p><b>Objective 1.</b> Land use and land cover of the wetland to be maintained in line with regulatory requirements under Wetlands (Conservation and Management) Rules, 2017 and East Kolkata Wetlands (Conservation and Management) Act, 2006.</p>	<p>No illegal transformation of land use</p>
<p><b>Objective 2.</b> Sewage quantity and quality received within the wetland to be efficiently treated applying traditional waste recovery practices.</p>	<p>Tertiary channels to deliver sewage without use of pumps</p> <p>Equitable distribution of sewage</p> <p>Heavy metals concentration in sewage to be brought to safe levels</p>
<p><b>Objective 3.</b> Maintaining the biodiversity within East Kolkata Wetlands.</p>	<p>No species extirpation</p> <ul style="list-style-type: none"> <li>• Improving the richness of species</li> <li>• Recover of species, if any</li> </ul> <p>Counts to be maintained in the range of 20% deviation from average of last five years</p> <p>Sighting to be maintained in the range of 20% deviation from average of last five years</p>
<p><b>Objective 4.</b> Species invasion threats to fisheries to be reduced.</p>	<p>To be reduced to a manageable level</p>
<p><b>Objective 5.</b> Sustainable livelihood with risk reduction.</p>	<p>Non-declining harvest of fish and vegetables</p> <p>Wetland communities having income in the lower quintiles to gain additional sources of income</p>

<p><b>Objective 6.</b> Individual and collective capacity and opportunities for stakeholders and wetland communities to participate in wetland management and contribute to wetland wise use to be enhanced.</p>	<p>Integration of community, rights and capacities in management plan implementation and monitoring</p> <p>An Integrated Wetlands Inventory, Assessment and Monitoring System to be put in place and used.</p> <p>Local action for addressing solid waste or preventing encroachment</p> <p>Community norms for preventing land use change or overharvesting of resources</p>
<p><b>Objective 7.</b> Systematic wetlands inventory, assessment and monitoring system is used to inform management decisions and assess effectiveness.</p>	<p>Data to be available on all priority wetland features and threats to them</p> <p>Data to be systematically analyzed and presented in EKWMA meeting</p>
<p><b>Objective 8.</b> Integration of multiple values of wetlands in sectoral developmental planning is enhanced.</p>	<p>EKWMA to meet regularly and consider implications of sectoral plans</p> <p>EKWMA to enable integration of role of EKW in programmes and actions plans on climate change, urban development, and disaster risk reduction</p>

Following actions are proposed under four management components, namely: a) Institutions and Governance, b) Water Management and Pollution Abatement, c) Species and habitats, and d) Sustainable Resource Development and Livelihoods.

#### Component I. Institutions and Governance

- Reorganizing and strengthening EKWMA with an efficient and results-oriented institutional structure, and gradual shift from an enforcement to a strategic role in wetland management. Reorganized set-up to have the following units: a) planning and design, b) enforcement, c) monitoring, and d) community engagement and communications.
- Demarcation of wetlands boundary by placing geo-tagged pillars, etc.
- Zoning of EKW.

- Setting up an integrated wetland inventory, assessment and monitoring system to address the overall information needs of wetland management and to provide robust decision support system.
- Annual Ecosystem Health Report Card publication and communication of wetland monitoring information to decision-makers and stakeholders.
- Research on carbon and GHG flux, nutrient budgets, bioaccumulation and multiple values, etc.
- Communication, Education, Participation and Public Awareness through signage at major points, webpage, establishment of community advisory group, resource material and workshop and public events, etc.
- Mid-term and end-term review of management plan implementation to assess the extent to which stipulated objectives have been achieved with a high degree of resource efficiency and in participation with stakeholders.

#### Component 2. Water Management and Pollution Abatement

- Dredging 14 highly silted canals measuring 43.8 km, to enable flow of sewage by natural gradient.
- Construction of a solid waste segregator near Bantala lockgate to regulate flow of solid waste.
- Development of constructed wetland system at Kulti outfall to augment waste water treatment beyond EKW region.
- Consider the feasibility of establishment of a plastic waste recycling unit in Dhapa region.
- Declaration of EKW region as a no-plastic zone.

#### Component 3. Species and habitats

- Regular monitoring of waterbird population following standard protocols of the Asian Waterbird Census at all the major congregation sites within and around EKW
- Culture of indigenous fish species in Goltala bheri.
- Mapping invasion pathways of Suckermouth Catfish, and undertake screening at various points along the DWF to prevent further spread.
- Revegetating peripheral areas of Nalban, Goltala Captain Bheri and some identified locations with Phragmites, Typha, Shola and other native species to improvise habitat of marsh mongoose, amphibian and reptilian species.
- Construction of a wetland interpretation centre at Bantala sedimentation tank area with facilities such as: exhibits, viewing gallery, watchtowers, waste recycling models, children play area, auditorium and souvenir shop.

#### Component 4. Sustainable Resource Development and Livelihoods

- Desilting bheri under cooperative ownership to increase productivity.
- Construction of 4 hatcheries at Dhapa Manpur, Tardha Kapasati, Kharki and Kantipota for production of seeds of Indian major carps and air breathing catfishes.
- On a demonstration basis, promoting crop diversification in agriculture and horticulture farms.
- Strengthening community health infrastructure and comprehensive access to safe drinking water and sanitation facilities.

## Budget and Financing

Over 2021-2026, implementation of management plan entails a budget of Rs. 110.7198 Crore. Of this, the component on institutions and governance is allocated 31.72%. This is followed by allocation of 26.80% for implementing actions under component for sustainable resource development. The components on water management and pollution abatement and conservation of species and habitats have been allocated 21.98% and 19.50% of the budget respectively. Being aligned with the objectives of National Mission of Clean Ganga (of the Jal Shakti Mantralaya) and the National Plan for Conservation of Aquatic Ecosystems (of the Ministry of Environment, Forest and Climate Change), the East Kolkata Wetlands Management Authority may consider seeking funding from these sources, along with allocations from the state budget.



## I Introduction

---

### I.1 East Kolkata Wetlands

The East Kolkata Wetlands<sup>1</sup> (EKW), located on the eastern fringes of Kolkata city, sustain one of the world's largest integrated resource recovery systems based on a combination of aquaculture, agriculture and horticulture (Map 1). The Government of India designated EKW as Wetland of International Importance under Criteria I of the Ramsar Convention in 2002 (as an example of wetland wise-use).

EKW is a critical natural infrastructure for Kolkata City. The wetland treats over 900 million litres of sewage generated by Kolkata Municipal Corporation every day (approximately 65% of total sewage generated in the metropolitan area) saving the city nearly Rs. 4,680 million annually in terms of treatment costs saved, as well as providing a much-needed flood buffer on the peri-urban interface. As the nutrient-rich effluent moves through a maze of fish farms, horticulture and agriculture system, it is progressively cleaned and nutrients are redirected to produce annually 20,000 MT of fish, 50,000 MT of vegetables and irrigate nearly 2850 ha of paddy lands. Over 60% of carbon from wastewater is also locked in various forms in the production process, thus reducing harmful Green House Gas emissions from the region. At least 380 species under major flora including 93 plant families, 10 amphibians, 29 reptiles, 123 birds, 79 fish, 24 crustaceans, and 13 mammal species have been recorded from these wetlands. The traditional waste recovery practice provides subsistence opportunities for a large, economically underprivileged population of 0.15 million living in the 37 revenue villages (locally called mouza) within its boundaries. EKW is also one of the few natural habitats providing recreational space for the urban and peri-urban population. The wetland provides strong arguments for the adoption of traditional knowledge of local communities for wise-use.

### I.2 Management Planning Purpose

The Management Plan outlines the commitment of the Government of West Bengal for conservation of the East Kolkata Wetlands through wise- use of this natural endowment and a series of developmental interventions. Located on the peri-urban interface of Kolkata City, the wetland has been under constant pressure of conversion for settlements and agriculture. Changes in land use, siltation of channels, pollution and stakeholder conflicts can impair wetland functioning. Mapping of the conservation area boundary for the wetland including the waste recycling region was done in

1985 by the State Planning Board, Government of West Bengal. Subsequently, the High Court of Kolkata passed an order in 1992 prohibiting changes in land use of the area and directed the state government to take recourse to statutory cover to protect EKW.

The notification of the East Kolkata Wetlands (Conservation and Management) Act in 2006 led to creation of the East Kolkata Wetland Management Authority (EKWMA) as a nodal agency for systematic implementation of wise use principles for the management of this Ramsar Site.

---

<sup>1</sup>The Ramsar Site was designated in 2002 with the name "East Calcutta Wetlands" and remains so in the Ramsar database. In 2001, the Government of West Bengal decided to change the name of its capital city to Kolkata. The Rules governing the wetland is titled 'East Kolkata (Conservation and Management) Rules, 2006' and the Authority as East Kolkata Wetlands Management Authority.



**Picture 1. East Kolkata Wetlands provide immense ecological subsidies to the City of Kolkata**

The Act took explicit cognizance of EKW as a Wetland of International Importance and its various ecosystem services, including the ability for the regulation of water regimes, wastewater treatment, source of groundwater recharge and other socio-cultural values. The Act defined the land use within the wetland, identifying each land parcel (as substantially water dominated, under agriculture, horticulture or settlements), and banned any further diminution of wetland area, change in ecological character, and overall land use. EKWMA was entrusted with the responsibility of implementing the Act, which included regulating land-use change, preventing unauthorized development and promoting integrated management of the wetland system. The current management plan is a response to the needs identified by the EKWMA to put in place an integrated strategy and action plan for ensuring wise use of the EKW.


### **1.3 Management to-date: An overview**

Upon its establishment in 2006, EKWMA collaborated with Wetlands International South Asia for formulating a management plan for EKW in 2008 for a period of five years. The goal of the plan was 'conservation and sustainable utilization of ecosystem services and biodiversity of EKW for ecological security and economic improvement of stakeholders'. The purpose was to 'establish effective management practices for EKW through coordinated actions at river basin level integrating coastal processes'. Seven management objectives related to institutional development, management zoning, water management, biodiversity conservation, sustainable fisheries development, sustainable agriculture development and livelihood improvement were laid down under the plan, to be implemented with a budget of Rs. 304 crores over five years (including funds from convergence sources).

The management plan was submitted to the MoEFCC for financial support under the National Wetlands Programme in 2009, and funding of Rs. 2.29 crores received during 2007-2013 for a limited number of activities, such as desilting of sewage channels, plantation, monitoring and implementation of zoning plan.


In 2010, the Wetlands (Conservation and Management) Rules were notified by the Ministry of Environment, Forest and Climate Change (MoEFCC), under the provisions of which discharge of sewage, dumping of solid waste and construction of permanent nature (except boat jetties) were prohibited in all notified wetlands, including all Ramsar Sites. Since some of these conditions were at variance with the State's East Kolkata Wetlands (Conservation and Management) Act, 2006 and hence remained unfulfilled for EKW, further funding from MoEFCC was stopped. Various interventions as envisaged in the management plan could not be implemented, resulting in the 2008 management plan implementation being kept in abeyance.

In 2017, the MoEFCC notified the Wetlands (Conservation and Management) Rules in supersession of the 2010 Rules. However, there was no change in the list of prohibited activities. The EKWMA requested the National Wetlands Committee (NWC) of the MoEFCC for granting special permission for EKW, as utilization of sewage from Kolkata was the core driver of wetland functioning. The Committee considered the request of EKWMA in its two meetings (held on July 26, 2018 and November 5, 2019) and asked the CPCB to define the optimum quantity of sewage that may be permitted to flow into EKW to maintain ecological functions of these wetlands. The EKWMA was advised to frame a comprehensive management action plan to address development as well as conservation aspects related to EKW.



**Map I. East Kolkata Wetlands**

The entire financing of activities of EKWMA since 2012 has been with the support of the Government of West Bengal, which has allocated Rs. 5.35 crores on the basis of annual plans considered in the Authority's meetings at the beginning of each financial year. The activities undertaken include: a) desilting of sewage canals, b) demolition of illegal constructions (demolition is done by Kolkata Municipal Corporation on the basis of the recommendation and financial support of the Authority), c) monitoring and research, d) wetland delineation and e) communication and outreach. An overview of receipt and expenses of the Authority for 2006-2019 is summarized in Fig I.



**Figure I. Financial flows of East Kolkata Wetlands Management Authority (2006-2019)**

Apart from these activities, the Authority has considered several developmental projects to date, notable being the proposal for construction of an elevated corridor over the wetlands to ease traffic congestion in the Kolkata City. The Authority forwarded the proposal for consideration of the National Wetlands Committee, which in turn in their meeting of July 26, 2018, and November 05, 2019 recommended a detailed Environmental Impact Assessment of the proposed intervention.

In 2019, a case against the EKWMA and Government of West Bengal on encroachment of the EKW area and violation of the East Kolkata Wetlands (Conservation and Management) Act, 2006 was filed in The Hon'ble National Green Tribunal. A task force was constituted to look into the solid waste dumping at Dhapa and Mollar Bhery area, encroachments and illegal plastic and industrial waste processing units along Basanti Highway. Following the report of the task force, the Government of West Bengal has submitted an affidavit to the Court confirming that a range of measures to control solid waste dumping were being taken up, including fencing of Mollar Bheri area and bio-mining of waste, provision of waste segregation units, organic waste processor units, and integrated waste processing unit. The Court also directed the Environment Department to submit an integrated management plan for wetland wise use. This plan has also been formulated taking into consideration the NGT's direction. The eastern zone bench of the NGT in their order dated 24 June, 2021 also directed the Additional Chief Secretary of the Department of Environment, Government of West Bengal to ensure immediate submission of the management plan to the MoEFCC for approval and necessary funding.

## 1.4 Approach and Methodology

Article 3.1 of the Ramsar Convention commits the Contracting Parties to put in place management arrangements to ensure wise use of all wetlands within their jurisdiction. The text of the Ramsar Convention defines wise use as "the maintenance of their ecological character, achieved through the implementation of ecosystem approaches, within the context of sustainable development".

Wise use is the longest established example amongst intergovernmental processes, implementation of which have become known as ecosystem approaches for conservation and sustainable development of natural resources, including wetlands (Finlayson et al. 2011). The approach recognizes human interdependency with wetland functioning and accommodates sustainable utilization of these ecosystems for the benefit of humankind in a way compatible with the maintenance of natural properties of the ecosystem. Wise use encourages stakeholder engagement and transparency in negotiating trade-offs and determining equitable outcomes for wetland conservation while promoting maintenance of environmental, economic and social sustainability (Finlayson 2012). The purpose of management planning is to outline the pathway through which wetland wise use can be achieved (Ramsar 2010).

The term wise use is often interpreted to indicate that human use of all wetlands is promoted by the Ramsar Convention. However, this needs careful consideration. The most recent update of the wise use definition was in 2005, wherein along with the definition cited in the previous paragraph, two footnotes were also placed. The first clarifies that 'ecosystem approaches' include the elements elaborated by the Convention on Biological Diversity – integrated management, stakeholders' participation in decision-making, transparency about tradeoffs, and equitability of the outcomes. Mechanisms such as integrated river basin management, integrated coastal zone management respond to this aspect.

The second footnote expands the phrase 'in the context of sustainable development' to recognize that development, though inevitable in most cases, is not an objective for every wetland. Wherever development is to take place, it has to be facilitated in sustainable ways by approaches elaborated in the Convention. Thus, when the concept of wise use is examined from the lens of sustainability, the elements of wetlands' conservation' and 'use' are reconciled to ensure that ecosystem retains capability for use now and into future, rather than 'using' or developing the wetlands at present. The onus of elaborating a pathway for achieving wise use outcome is on the management planning process.

The Ramsar Convention, in its Resolution XI.11 of 2012 has laid down principles for planning and management of urban and peri-urban wetlands. The resolution recommends that: a) thematic planning should be used as an essential tool to safeguard wetlands and their ecosystem services both within and beyond urban settlements; b) the consideration of wetlands within urban planning needs to be integrated fully with wider elements of spatial planning (such as Integrated River Basin Management, water resource management, the development of transport infrastructure, agriculture production, fuel supply, etc.), and c) alternative locations need to be identified for planned urban developments (both formal and informal built development) which do not lead to wetlands, or other natural ecosystems, being degraded or lost. It is suggested that environmental appraisal committees at government level should invariably involve wetland experts to gather valid opinion about the area before the site is granted clearance for developmental projects.

Conservation and sustainable development of EKW requires integrated planning and resource management at the river basin level recognizing the interconnectedness of the wetland system with its catchments. River basin-level planning requires an understanding of the carrying capacity of the basin to produce desired outputs (goods and services) from limited resource base and achieving equitable quality of life while maintaining desired environmental quality in the region. The planning for management calls for trade-off between desired production and consumption levels. It also emphasizes on the development of supportive mechanisms within the generative capacity while maintaining the environmental quality. The challenge, therefore, is to conserve the ecological character and full range of ecosystem services of EKW while providing sustained benefits to the communities dependent upon the wetland for their sustenance.

The methodology for management planning is based on the New Guidelines for Management Planning for Ramsar Sites and Other Wetlands as adopted by the Contracting Parties to the Ramsar Convention on Wetlands in 2002. These guidelines also form the basis of the wetlands management planning guidelines of the MoEFCC's National Plan for Conservation of Aquatic Ecosystems (NPCA).



**Figure 2. Step-wise wetland management planning process**

The NPCA guidelines recommend following a diagnostic approach – wherein the selection of management interventions is guided by knowledge of wetlands features and factors governing these features, and their relationship with broader societal conservation and development goal that wetland wise use is contributing to (Fig 2). Wetlands features are its ecological, social and institutional attributes, which collectively characterize a wetland. Wetlands are dynamic systems, and thus their features undergo cyclical and temporal changes. Factors (natural as well as anthropogenic) cause the wetland to move along a specific trajectory.

As the management plan was framed during COVID-19 epidemic, extensive fieldwork and stakeholder consultations were not feasible. The first draft of the management plan was reviewed by the Technical Committee of the Authority in a virtual meeting held on July 27, 2020. Then followed a series of informal consultation with the wetland experts and internal reviews. It is also envisaged that this plan will be widely circulated and feedback of stakeholders taken on board prior to initiating implementation.

### 1.5 Management Plan Structure

The management plan includes eight chapters, in line with the structure recommended by the MoEFCC. The current chapter provides the management planning context. The chapters 2 and 3 contain description and evaluation of wetland features and assessment of risk of adverse change. The effectiveness of current institutional arrangements in preventing the adverse change in wetland ecological character is analysed in Chapter 4. The management planning framework is discussed in Chapter 5, and the monitoring plan in Chapter 6. The action plan is in Chapter 7 and budget and financing discussed in Chapter 8 of the plan.




## 2 Description of Wetland Features

### 2.1 Physical regime

Located between 22°25'N to 22°35'N and 88°20'E to 88°35'E, with the city of Kolkata on the west and River Kulti on the east, the EKW is a mosaic of landforms including predominantly water dominated areas (used as fish farms, locally known as bheri) to land centric usages for agriculture and horticulture (Figure 4). Settlements are interspersed between various land uses. The use of sewage as the basis of aquaculture, agriculture and horticulture production systems provides the rationale for considering the entire area as a single management unit. The wetland boundary has been defined in the East Kolkata Wetlands (Conservation and Management) Act, 2006.

As per images of May 2019, the wetland has 48% area under fish farms, 27% under agriculture and 20% under settlements. Canals and horticulture areas account for 2% of the area respectively, whereas the landfills account for the remaining 1%. Being highly regulated, the wetland does not undergo much transformation in land use and landcover during pre and post monsoon season (Map 2 and Figure 3).



**Figure 3. Land use and Land Cover transformation in East Kolkata Wetlands (2019)**

The current wetland area is remnant of series of brackish water wetlands, which till the beginning of 20<sup>th</sup> century maintained active connection with the freshwater as well as marine environments of the Gangetic Delta Rivers and the Bay of Bengal, in an ecological continuum with the Sundarbans (also referred as Bidyadhari wetlands). The Kolkata City grew on the levees of River Hooghly in the sixteenth century, virtually without any drainage system. Initially, all the solid waste and sewerage was dumped into the river. Frequent outbreaks of malaria, plague and other diseases forced the city planners to abandon this practice and construct a system of sewers to discharge waste into EKW – then considered a malarious jungle with limited societal value. The wetland system acted as a vast spill area for the flood and tidal flows for the discharge of Hoogly and Bidyadhari Rivers, in the process attenuating floods of the Kolkata City. However, the wetlands gradually lost their connectivity to freshwater and tidal environments due to deltaic processes. Human interference primarily in the form of channelization further quickened the process of silt deposition within the channels, and finally

the Bidyadhari became defunct by the end of 18th century. With this, the sewage from the city became the dominant flow, which was used as an opportunity by private entrepreneurs to establish wastewater aquaculture, horticulture and agriculture in the late 18th and early 19th centuries. In 1864, a portion of the salt lakes was acquired for dumping solid wastes. Thus, over a period of time, the wetlands transformed from brackish water lakes to sewage fed fish farms, with large areas being converted for settlements and agriculture development. The post-independence surge of refugees to Kolkata City made the town planners further look into expansion of the urban area. This prompted reclaiming of more than 1,000 ha of the northern portion of the wetland for the establishment of the Salt Lake City. In 1969, redistribution of land under land reforms led to further filling up of approximately 2,500 ha of water bodies for conversion into paddy fields. It was only during the eighties that the role of over 260 fish farms in treating wastewater caught planners' attention, and the events leading to mapping of the WRR took place and the site designated as Ramsar in 2002.

The EKWMA, after its establishment in 2006 was entrusted with the task of boundary delineation based on the East Kolkata (Conservation and Management) Rules, 2006. Till then, the map prepared by the State Planning Board in 1985 and later on surveyed by Creative Research Group in 1996 and translated into geospatial framework was the basis for planning. During 2005, the Department of Environment delineated the wetland boundary on cadastral maps at 1: 4,000 scale (104 such sheets were prepared).



**Picture 2. Part of Dhapa landfill in East Kolkata Wetlands which has been treated**






Figure 4. Schematic diagram of East Kolkata Wetlands resource recovery system




**Map 2. Seasonal transformation in land use and land cover of East Kolkata Wetlands (2019)**

To capture land use and land cover changes in the two decades since Ramsar designation, a change assessment was conducted using satellite data of November 2000 (Landsat 5, 30 m resolution, November 9, 2000) and November 2019 (Sentinel 2A, 10 m resolution, November 21, 2019) (Map 3 and Figure 5). During this period, the area under fish farms, settlements and landfill has been observed to increase, while the area under horticulture and agriculture declined. Along the southern margins of the wetland (around Tardaha and Tardaha-Kapashati), extensive transformation of agricultural lands to fish farms can be seen, primarily attributed to the increased availability of sewage as an outcome of canal desilting operations. The area under waste dump have also increased, from just one location within the wetland (Dhapa) in 2000 to creation of a new site at Mollar Bheri. However, due to environmental concerns, waste dumping at Mollar Bheri has been stopped completely and steps have been taken to convert the legacy waste into assets.

The increase in area under fish farms during 2000 – 2019 is striking and significant. A longer term change assessment based on remote sensing images of 1973, 1989 and 2010 had indicated a continuous decline in fish farm area (1890 ha during 1973-1989 and an additional 1959 ha during 1989 – 2010) (Parihar et al. 2013). Similarly, a change assessment for 1990 – 2011 period had indicted a loss of 1,476 ha of area under wetlands (fish farms) to cropland and expansion of settlements (Mondal et al. 2017). A land use and land-cover change analysis matrix presented in Figure 6 reveals that increase in fish farm area is largely on account of transformation from agriculture, which has become more prominent after 2015 and in all likeliness due to the enhanced productivity of water bodies vis-à-vis farm land due to increased flow of waste water in the area.



**Figure 5. Changes in area under different land use and land cover categories in East Kolkata Wetlands (2000-2019)**



**Map 3. Land use and land cover of East Kolkata Wetlands (November 2000 to November 2019)**



**Picture 3. Fishing activities in East Kolkata Wetlands**

The boundary demarcation based on physical verification of the site and examination of land records is an ongoing process. The EKWMA, in consultation with Land and Land Reforms Department (currently known as Land & Land Reforms and Refugee Relief and Rehabilitation Department) has identified 2540 waypoints with geographical coordinates using geo-rectified satellite imageries and ground truthing. The work is nearing completion. The Authority plans to place physical pillars to demarcate the wetland boundary on ground.

The Wetlands (Conservation and Management) Rules, 2010 subsequently replaced with the Wetlands (Conservation and Management) Rules, 2017 prohibit permanent construction of any nature within the wetland due to which the EKWMA has stopped admitting any applications for land use conversion. However, some conversions have taken place illegally. The Authority has filed as many as 357 FIRs in the eight police stations to report and take action against illegal land conversion and encroachment. However, the scale of encroachment is much larger considering the growing pressure on land. Some gaps between the records of the Land and Land Reforms Department and the Schedule appended to the East Kolkata Wetlands (Conservation and Management) Act, 2006 have also been noted.




Figure 6. Land use transformation in East Kolkata Wetlands (2000-2019)



## 2.2 Wetland catchments

The EKW forms a part of the wetland regime of the Gangetic Delta. The definition of Gangetic delta extent has evolved over a period of time. A group of experts belonging to an older school delineated the triangular tract encompassed by the Bhagirathi-Hugli in the west, the Ganga- Padma in the north, combined flow of the Jamuna and the Meghna in the east and the Bay of Bengal in the south as the Ganga delta (Bagchi, 1944). Considering the role of the Brahmaputra in delta formation, the scholars of modern school treat the entire alluvial plain of Bengal as the Ganga- Brahmaputra (GB) delta (Allison et al.2003; Bandyopadhyaya, 2007). It may be more appropriate to refer to the delta as Ganga-Brahmaputra-Meghna (GBM) delta, considering also the role of the latter in the depositional process. GBM delta is essentially a coalesce of deltas of the three rivers (Rudra, 2018), with an estimated deposit of half a million kilometer<sup>3</sup> of sediments.

EKW is an area of incomplete morphogenesis where fluvial and marine land-building processes were juxtaposed. These wetlands were connected till about a century ago with the Bay of Bengal through Matla-Piyali-Adiganga distributaries, and was influenced by tidal action. The Bidyadhari, Sunti and Noai brought in freshwater into the wetland from the upstream reaches.

Over time, Jamuna, a cross channel connecting Hugli with Ichhamati River became moribund. Changes also took place in the drainage of River Bidyadhari. The river originates from the Mathura bill near Kalyani and bifurcates at Tehatta. The western branch of the river, which used to flow through the present site of Salt Lake City and discharged through Matla estuary can hardly be traced, whereas the eastern branch continues to flow into the river Kulti Gang. During late 18<sup>th</sup> century Tolly's Nala was excavated between Garia to Samukpota, connecting Adiganga with the western branch of Bidyadhari. However, owing to the shifting of the main course of the Ganges from Bhagirathi to Padma and westward migration of the former near Kalyani, the freshwater flows to the Bidyadhari rapidly reduced leading to gradual domination of the tidal process over fluvial one.


The north south hydrological exchange between the freshwater and brackish water regimes was further altered when a system of locks and pumps were put in place in Kolkata City to drain its stormwater and sewage into the EKW. The flow regimes of the wetland have thus come to be dominated by sewage from the city, and their utilization for fisheries and agriculture. Therefore, since the mid-19<sup>th</sup> Century both fresh and brackish water regimes became feeble. The drainages of Bidyadhari in the upstream gradually merged into the Kulti estuary, which is connected to the Bay of Bengal.

Considering the hydrological regimes, the southern part of GBM Delta forms an indirect catchment of the EKW. Yet, Kolkata, Piyali-Adiganga, Bidyadhari-Sunti-Noai (often referred as Kulti Upper), and Kulti lower basins are most significant (Map 5). Since the excavation of east-flowing wastewater canal, the hydrology of the EKW drastically changed and it was metamorphosed into a wastewater wetland. Hence, EKW receives the flows from the Kolkata Basin, i.e. those arising from the core areas of city system, suburban system and Manicktala, Topsia-Tangra, Tollygunge-Panchannagram systems, North and South Salt Lake basin and a very little part of lower Bagjola Basin (Map 6). Thus, the direct catchment of EKW comprises the following six sub-basins of Kolkata Basin:


- a) Kolkata Sub Basin which contributes sewage flow as well as storm flow into the wetlands through an intricate network of channels, ultimately terminating into Dry Weather Flow and Storm Weather Flow channels.
- b) North Salt Lake Basin comprises of the Kestopur canal. Bhangor Kata Khal, Circular canal.
- c) South Salt Lake Basin consists of a low-lying area subject to inundation during the wet season. Nodor khal, Deyara Khal, Lalkuthi khal are flowing through this basin.
- d) Kheyadaha-Sumidgiri Basin carries the sewage mainly by Sumidgiri canal, Bidyadhari canal, Boye Nala. Tilly's Nala is the extreme south boundary of both the South Salt Lake Basin and the Kheadaha -Sumidgiri basin.
- e) Tollygunge-Panchannagram (T.P.) Basin carries the sewages of Jadavpur, Santoshpur, Kasba, Tiljala area of KMC. This area was basically a very low marshy land. Tolly's Nala and SWF are the southern and northern boundary of this basin respectively. A large number of intercepting channels and tributaries crisscrossed with T.P. Channel are passing through this basin.
- f) Lower Bagjola Basin. Very little of this basin falls within the EKW.




**Picture 4. The Storm Water Flow Channel carries the monsoon runoff directly to Kulti River**



**Map 4. Wetlands of Southern Fraction of Ganga-Brahmaputra-Meghna Delta**



**Map 5. Catchment of East Kolkata Wetlands**



**Map 6. Direct catchment of East Kolkata Wetlands**

### 2.2.1 Land use and land cover

Agriculture is the predominant land use within the indirect catchment of EKW, accounting for 59% of the total area, followed by settlements. Wetlands constitute 17% of the indirect catchment area. There is hardly any perennial vegetative cover within this region. The EKW located within the Kolkata Basin displays a complex mosaic of land use categories including water spread areas, agriculture, horticulture and settlements.


The land use within the East Kolkata Wetlands has been changing with the growing developmental pressures from the metropolitan area as well as population within. As compared to 1959, there is a discernible conversion of wetlands for non-wetland purposes. The reclamation of Salt Lake city and construction of Eastern Metropolitan Bypass has also led to shrinkage of wetlands on the western boundary. Since 2000, a distinct conversion of wetlands to agricultural land and settlements can also be recorded (Mondal et al. 2017; Parihar et al. 2013).

### 2.2.2 Geology and Geomorphology


Located within the lower deltaic alluvial plain of the Ganga-Brahmaputra-Meghna Delta, the EKW is underlain with sediments deposited in successive stages by the river and consists of layers of clay, clay and calcareous concretions, peaty intercalations, sandy and silty clay and fine to coarse sand. The water table throughout the basin is high.

Based on lithological, floral, faunal and radiocarbon dating (Chatterji et al. 1959; Banerjee et al. 1984; Sen and Banerjee 1990; Barui and Chanda 1992; Hait et al. 1996) clay silt with a peaty intercalation is present below the wetlands at two depth ranges from the surface 2-5 m and 12-12.6 m in the Holocene series of the Quaternary system, Sand, fine to coarse with clay lenses, gravel and calcareous concretions is present at a depth of 40 -246 m in the Pleistocene series of the Quaternary system and Clay layers are present from 246 – 614 m in the Pliocene series of the Tertiary system. Piezometric wells with screens placed at 40-220 m indicate that groundwater flows in a confined condition within a sandy aquifer sandwiched between two clayey beds (Sikdar 2000; Sikdar et al. 2002).

The catchment slopes from northwest to south east, the relief in the northern side is over 11 m amsl and in the southern side it is under 3 m amsl and pockets of areas within the catchments are below sea level. EKW is a part of the lower deltaic plain of the Ganga– Bhagirathi–Hugli River system wherein the land surface with its elevation of 2–6.5 m amsl slopes gradually towards the south and southeast (Map 8). This elevation differs locally because of paleo-levees, paleo-courses and stream channels.



Map 7. Aquifer system in East Kolkata Wetlands Catchment (India-WRIS 2020)




**Map 8. Elevation profile of East Kolkata Wetlands catchment**




### 2.2.3 Climate

EKW catchment has a Tropical wet-and-dry climate (Köppen climate classification Aw). The annual mean temperature is 25 °C; monthly mean temperatures ranges between 17 °C and 32 °C (Stackhouse, 2020). Summers are hot and humid with temperatures in early 30's and during dry spells the maximum temperatures often exceed 40 °C during May and June. Winter tends to last for only about two and a half months, with seasonal lows dipping to 10 °C between December and January. The highest recorded temperature in the past 30 years is 46.6 °C and the lowest is 3.2 °C (Stackhouse, 2020). Often during early summer, dusty squalls followed by spells of thunderstorm and heavy rains fall in the catchment, bringing relief from the humid heat. These thunderstorms are convective in nature, and is locally known as *Kal baisakhi* (Nor'westers). Rains brought by the Bay of Bengal branch of South-West monsoon fall between June and September and supplies the catchment with most of its annual rainfall of ~1300 mm, the highest rainfall occurs during the monsoon in August ~306 mm (Stackhouse, 2020). The annual Average rainfall is ~1600 mm with an average humidity of 80% and wind speed ranging from 2.9 to 7.0 km per hour. Solar radiation varies between 150 and 250 Langley/day (Ghosh 2018). The average evapotranspiration ranges from 150- 200 mm per year (Stackhouse, 2020). The climate of the area is predominantly influenced by northeast and southwest monsoons.



**Figure 7. Precipitation trend in East Kolkata Wetlands direct catchment (1989 - 2020) (Stackhouse, 2020)**

The City of Kolkata is perennially exposed to urban flooding which is the most critical climate-related hazard in Kolkata. Hydrological and hydraulic impacts resulting from increased precipitation in a changing climate by 2050 indicate that the low lying areas of the city, essentially wetlands, are expected to be under prolonged inundation (Dasgupta et al. 2013).




**Figure 8. Temperature and evapotranspiration trend in East Kolkata Wetlands Basin (1989 – 2020) estimated using Thornthwaite’s equation (Stackhouse, 2020)**

## 2.3 Hydrological regimes


### 2.3.1 Hydrological set-up

Hydrological regimes of the EKW are highly modified and occur through drainage and sewerage channels which connect the urban and peri urban Kolkata located on the bank of Hooghly to the outfall system in Kulti. The flows of Kolkata sub basin, of which the wetland forms a part, are ultimately discharged to the two rivers, i.e. Hooghly and Kulti Rivers through its 1,412 km length of drainage system and interconnected network of drains and channels. Flows from the urban centers are transferred into the lead channels through a system of pumps. As the two rivers are subject to tidal actions, regulators have been constructed at the outfalls of channels to enable controlled discharge of the sewage.

Located within a deltaic area, and within a system subject to tidal action, the inflow and outflow are conditioned through various hydraulic structures, including sewage pumps and regulators (Maps 9 and 10). The operation of these structures is governed by the need to ensure minimum waterlogging by transporting stormwater quickly from the Kolkata City and secondarily to ensure sewage availability within the fish farms and agriculture areas of the wetland. During the monsoons, all structures are synchronized to ensure water drainage to Kulti bypassing the wetland, with the locks ensuring that no tidal influx takes place towards the system. During the lead seasons, the structures are operated to pump sewage towards the wetland and thereafter to Kulti River. The Kolkata Municipal Corporation generates approximately 4500 MT of solid waste daily which is nearly entirely dumped at the Dhapa landfill located within the EKW.




**Map 9. Sewage treatment systems in KMC Area**



Map 10. Hydraulic structures in East Kolkata Wetlands Basin


### 2.3.2 Water and sediment regime

Presently, the overall inflows from the upstream watersheds to the wetland primarily comprise two elements, i.e. dry weather flows and storm weather flows. Wastewater generated in Kolkata City accounts ~1400 MLD (Ghosh 2018) of which the total sewage treated in the EKW fisheries has been estimated as ~910 MLD to 930 MLD (KMC Drainage Report, 2020), which is roughly 65% of daily sewage generation. However, given the fact that the Tollygunge – Panchannagram Basin and Topsia – Tangria Basin discharge directly into the SWF channel, which is then bypassed into the Kulti River, only 78% of the DWF emanating from the Kolkata city and its sub urban areas is drained into the wetlands, and is available for nutrient recycling. The storm weather flow emanating from these watersheds is linked to the rainfall patterns. An analysis of the rainfall within the Kolkata Basin indicates a concentration of 75% of the average rainfall of 1,600 mm within June – September. The total SWF generation from the upstream watersheds has been estimated to be 136 MCM. Including the dry weather flows, the overall flow generation from the upstream watersheds is 498 MCM (Figure 9).



**Figure 9. Inflows in East Kolkata Wetlands (Modelled)**

An assessment of inflows from the upstream watersheds indicates that about 40% of upstream flows generated gets into the wetland. During the lean seasons, the wetland receives the DWF flows from the municipal region. During monsoons, the hydraulic structures divert the entire flows to the SWF, reducing the overall upstream inflow by nearly 85%. As the level of SWF is lower than the DWF, the flows naturally get into the former, and drains to Sundarbans through Kulti.



**Map 11. Canal system of East Kolkata Wetlands**



**Picture 5. The Dry Weather Flow Channel**

One of the key consequences of flow regulation through extensive channelization has been rapid enhancement in the natural siltation and sedimentation process within the Kolkata City sewerage system. In a natural course, the sediments would be expected to be accumulated within the wetlands, but due to extensive regulation of flows, there has been a higher rate of sedimentation within the upper watersheds of East Kolkata Basin. Another impact of extensive channelization and subsequent flow reduction is loss of self-cleansing velocity which conveys wastewater or storm water without long term deposition of solid material. A common design criterion is to maintain a value of 0.80 m/ sec with peak flow and 0.60 m/sec at average flows. However, with extensive siltation the hydraulic capacity of the various canals has got reduced by 15% to 50% and this self-cleansing velocity cannot be maintained, which further augments siltation. In downstream reaches, within the channels of the wetland, the entire monsoon flows from upstream do not flow through the secondary channels, and therefore no flushing takes place. The limited monsoon rains available over the fish ponds are completely insufficient to ensure channel flushing. As a result, the ponds have to use pumps for lifting sewage, which would otherwise be available through normal gradient flow.

The EKW being a natural depression between the Kolkata city and Kulti estuary provides an enormous water holding capacity. In its natural regimes, this water holding capacity enabled the wetland to regulate the flow regimes and attenuate floods by storing peak monsoon flows as well as tidal flows. However, this natural function of the wetland has not been integrated into spatial planning, and instead reliance is on engineering structures wherein the monsoon flows are transported out from the basin into the downstream Kulti River. Given the fact that the delta building process within the banks has been prematurely interrupted and the city has only marginal elevations as compared to other reaches, the flood attenuation function of the wetlands would have provided a natural buffer against high flows.

Water use within the wetland is predominantly governed by its use for fisheries and agriculture within the EKW. Fisheries within the wetland are exclusively dependant on the sewage inflows, whereas agriculture uses monsoon runoff as well as ground water, apart from effluent discharge from the fish farms. The current allocation pattern within different users has not been assessed comprehensively. Sewage for agricultural purposes is only accessible to mouzas fringing the fish farms. Agricultural areas lying at the tail end of the DWF channel and towards the north have no access to sewage and therefore depend on monsoon or groundwater through shallow pumps. In the long run, this has also been responsible for gradual reduction of agriculture in the wetland.




**Picture 6. The Bantala regulator from where sewage is diverted to the wetland**

### 2.3.3 Groundwater

The groundwater in EKW region is available from an average depth from the surface at 4-8 m in pre monsoon, 1-9 m in the monsoon and 1-10 m in the post monsoon periods. Due to the shallow levels of groundwater, an alluvial media and a hydraulic conductivity, fluctuations between the groundwater levels at shallow ranges are highly variable.






**Map 12. Groundwater levels in East Kolkata Wetland**

Groundwater levels in post monsoon (Map 12) indicate that a groundwater trough exists within the wetland near Dhapa–Manpur region and the groundwater moves towards this depression from the surrounding region. Several groundwater mounds are found in the adjoining areas of the wetlands from which the water flows in all directions. In and around Bantala the piezometric levels are within a meter below the surface, groundwater abstraction must be restricted in this region as it would result in volumetric compression the underlying aquiclude. If this water contains toxic material then the freshwater aquifer will be polluted. But in general, the groundwater flows from east to west.

An increasing trend of groundwater extraction in Kolkata has been observed since 1950s till present. In 1956, the groundwater pumping rate was ~ 55 MLD. With development of borewell technologies, the screens of the drinking water wells were placed within a depth interval of 100 to 140 for deep bore wells and 30-60m for shallow wells, for domestic use in 2010. Kolkata now pumps about 310 MLD of groundwater. It is anticipated that by 2025, 385 MLD of groundwater may be required for drinking, domestic and industrial purposes in Kolkata. This rate was calculated by projecting the past trends of groundwater abstraction at Kolkata for the period 1956–2006 (Sikdar 2000, Central Ground Water Board 2004) by the projected population in 2025.

In Salt Lake City, pumping began in 1969 with the construction of a single tube well with a discharge of 2.4 MLD. At present, 19 MLD of groundwater is extracted for drinking and domestic purposes. At the KLC Project area 5 MLD of groundwater is extracted today to meet the demand of the various industrial units. Industrial activity in the KLC Project area is likely to increase, requiring an additional 30 MLD of water by 2025. In the absence of any other source of water, this additional need will be met from 16 new tube wells planned to be constructed in the near future (Sahu et al, 2013). At present to meet the growing demand for agriculture and aquaculture, groundwater is being extracted at fast rate.

Unsustainable withdrawal of groundwater in the EKW due to rapid urbanization, aquaculture, agricultural and industrial development has posed a risk of land subsidence. The subsurface geology of the area consists of quaternary sediments of various grades and as groundwater occurs mostly under confined conditions except in those places where the top aquitard has been obscured due to the paleo channels. Currently, the hydraulic head exhibits a falling trend and it accelerates due to further overuse of groundwater, which leads to aquitard compression leading to land subsidence. Estimated land subsidence rate of the area ranges between 1.1 and 43.8 mm/year with an average of 13.5 mm/year (Sahu and Sikdar 2011).



**Map 13. Groundwater levels and flow direction in East Kolkata Wetlands**

### 2.3.4 Water quality

Wastewater of the city is discharged into the fish farms locally known as bheri. This wetland system exhibits immense potential in remediating the water quality by reducing the high amounts of biochemical oxygen demand (BOD) and chemical oxygen demand (COD) it receives from the inflow on a daily basis. Each hectare of the shallow waterbody has the ability to remove 237 kg of BOD per day (Ghosh 2018). The bheri being shallow allows full vertical circulation of water to the surface where algal blooms occur. A solar radiation that is about 250 Langley's a day, allows sufficient photosynthesis which augments reoxygenation to allow for efficient BOD and pathogen/faecal coliform reduction. Factors that play role in water purification are the shallow ponds acting as stabilization ponds, abundant water hyacinth that absorbs heavy metals, sun light penetrating to the bed of the water body and other microbial components that help in bioremediation (Raychaudhuri et al, 2008).

The characteristics of wastewater is not uniform round the year, it actually varies not only in the different sections of the wetland but also from hour to hour at the same place depending on the weather conditions, availability of water and other agro-economic conditions.

Water quality is monitored by West Bengal Pollution Control Board on a monthly basis. The seasonal trend analysis of 2018-2019 period is illustrated in Figures 10 -16. Significant BOD and COD reduction is noticed as the inflows travel from Bantala lock gate to Ghaskhali lock gate passing through the fish farms. During monsoon with the dilution from rainwater falling directly over the surface of the wetland the overall concentration of BOD and COD is relatively lower as compared with other seasons. Thus, with the reduction of the BOD and COD the DO levels also rise during monsoon. The DO is observed to be at its best at the Bamanghata Jalpath outlet station.

On the other hand, when the water flows from Bantala lock gate to Ghusiaghata lock gate travelling approximately 28 km, initial reduction of BOD, COD and increase in DO is observed, but, the dynamics is changed when the water travels around the Kolkata Leather Complex zone where a spike of BOD, COD takes place due to the occasional discharge of tannery effluents, which gradually subsides as it reaches Ghusiaghata lock gate. The Kolkata Leather Complex has 4 CETPs of 20 MLD capacity functional and 4 more modules of 20 MLD capacity are expected to be functional by March 2022. These are positive developments for EKW.

Total organic nitrogen (TON) is lowered by almost 30% from levels recorded at Har Har Bheri inlet to Lock gate at Ghaskhali. Nitrate and ammonium levels are somewhat reduced due to the wetland function in the Uttar Hatgachha. Their levels are nearly uniform for post monsoon and summer seasons. The levels drop in the monsoon season and rise by almost a factor of two by the winter season. Phosphate levels drop 50% from summer to monsoon and rise by a factor of three in post monsoon and winter season. The observed phosphate levels at Uttar Hatgachha is significantly lower when compared with the levels at other monitoring stations.

The presence of heavy metals in inflowing sewage and fish farms has been debated on several occasions. Monitoring records of WBPCB do not indicate presence of heavy metals in any of the sampling sites. However, several independent studies contradict these findings (Nandi et al., 2013; Dutta et al., 2016). On the other hand, studies confirm bioaccumulation of heavy metals in fish and vegetables produced in EKW (Joystu 2017; CPCB 2019).

A study conducted by (Sahu and Sikdar, 2008) for 40 water samples from different hydrochemical facies around EKW indicated that 7.5% groundwater samples show "excellent" quality, a majority of the samples, about 52.5%, fell in the category of "good water"; 27.5% of the water samples were of the "poor" quality and "very poor" quality is shown by 7.5% of the

collected groundwater samples. The study also pointed out that majority of the poor and water unsuitable for drinking was found within Bantala area and its buffers approximately up to 10 kilometres. More recent and recurrent assessments of groundwater quality is required to ensure that the water used by the communities living in the wetland is safe for use.




Figure 10. BOD at various stations in East Kolkata Wetlands (2018-20)




Figure 11. COD in various stations of East Kolkata Wetlands (2018-2020)




Figure 12. Dissolved Oxygen levels in various stations of East Kolkata Wetlands (2018-2019)




Figure 13. Total Nitrogen levels in various stations of East Kolkata Wetlands (2018-2019)




Figure 14. Nitrate levels in various stations of East Kolkata Wetlands (2018-2019)




Figure 15. Phosphate levels at various stations of East Kolkata Wetlands (2018-2019)





Figure 16. Ammonium levels at various stations in East Kolkata Wetlands (2018-2019)





Map 14. Water quality sampling stations in East Kolkata Wetland

### 2.3.5 Climate regulation

The low lying EKW act as storage reservoirs by absorbing majority of the excess flow from rainwater runoff through its storm-water flow channel (SWF) and protecting the city from frequent monsoon flooding. The water holding capacity of the aquaculture ponds increases by an average depth of 10 to 15 cm during the monsoon season. The area has not experienced flooding for the last 3 decades, regardless of the intensity of rain or quantum of flow of water into the wetlands. In addition, a large number of outlet channels syphoned to divert excess water have been designed by the local residents of the area, which contributes greatly towards averting floods (Banerjee 2017).

EKW is also referred to as the lungs of the Kolkata city. The air flowing over the waterbody humidifies the vapours into aerosol form, which captures dust and pollution particles and removes them permanently from the air stream. A study (Disha 1996) indicated that the air samples collected over Dhapa dumping site had reduced SPM concentration over 67% in a village site downwind in the EKW system (Banerjee 2017).

The wetland system contributes to significant (over 60%) carbon reduction of the Kolkata city's emission. Unique phenomenon of algae–bacteria symbiosis occurs where algal photosynthesis, in the presence of sunlight, releases oxygen resulting in reduction of BOD. The solar energy trapped by the phytoplankton (blue-green algae) is consumed by the fishes. The plankton plays an important role in degrading the organic matter in the sewage discharged into the wetlands. Over-growth of plankton might cause a problem of algal bloom, but the fishes grazing over the planktons keep it in check. The wastewater treatment in fish ponds is a direct contributor in reducing the GHGs to a great extent. A single STP treating the same amount sewage with similar efficiency to EKW would emit ~3500 tonnes of CO<sub>2</sub> per year. EKW on the other hand due to its natural system of using solar energy is saving Kolkata city from ~3500 tonnes of CO<sub>2</sub> per year (Mukherjee and Bardhan, 2019). Therefore, urban wetlands like EKW contribute towards urban resilience which is crucial for climate change mitigation and urban sustainability.

## 2.4 Species and Habitat

The biota of EKW co-exists within a mosaic of densely populated and intensely fragmented landscape. Over the last hundred years or more, the EKW has undergone changes in salinity regimes with corresponding drastic changes in floral and faunal diversity. The review of literature indicates that wetland harboured a rich floral and faunal diversity when connected to the freshwater and tidal regimes. But presently, with the wetland being predominantly managed as a sewage fed aquaculture system, only a limited range of species is supported. Disappearance of marshes for development of aquaculture has seriously affected diversity.

### 2.4.1 Vegetation

The BSI has prepared a list of noteworthy plant species of EKW which is also incorporated in Chandra, Raghunathan and Mao, 2020. The list includes 381 species of 93 families (Fig 17). The first comprehensive study on flora of Salt Lake was carried out in the early nineteen twenties, wherein 70 species of plants from various zones distributed within embankments, main wetland and terrestrial area were recorded (Biswas 1927). The vegetation of the embankments and bunds was mostly colonized by the plant species dominated by *Fimbristylus ferruginea*, *Suaeda maritime*, *Acanthus illicifolius*, *Excoecaria agalocha*, *Avicennia officinalis* etc. The main




**Picture 7. Diverse habitats in East Kolkata Wetlands**

wetland area was dominated by numerous algal species, in the deeper pockets of the wetland area and tall vegetation dominated by *Phragmites karka*, *Aegiceras majus*, *Typha elephantine* etc. The swamps and dry lands were mostly dominated by oligihaline and mesohaline shrubby plant species and several halophytic trees like *Sonnerata apetala*, *Avicennia officianalis*. A number of filamentous algae *Enteromorpha intestinalis*, *E. prolifera* etc) have been also reported. (De et al., 1989) recorded 97 plant species belonging to 41 families, among which 34 species were confined to saline water habitat. He also mentioned the presence of mangrove flora in the region. However, the recent study carried out by (Ghosh and Ghosh, 2003) a total of 106 aquatic plants belonging to 70 genera and 36 families have been reported. Some other studies also reported 96 species under 79 genera and 38 families (IWMED 2004). Among which there is hardly any submerged vegetation in the core fishing area except the plankton communities.

From the earlier studies it is inferred that in the past, Salt Lake supported rich vegetation including mangroves and other brackish-water species. Till date there are some remnants of earlier salt water vegetation like *Excoecaria agallocha*, *Achrostichum aureum* and *Acanthus ilicifolius*. However, distribution of these species is very sparse.

EKW harbours a variety of economically important plant species. Some of these plants have tremendous medicinal value and used traditionally by the communities for human and veterinary consumption. Besides several plant species such as *Bacopa monnieri*, *Enhydra fluctuans*, *Ipomea aquatica*, *Marsilea minuta* are used as vegetables by the local communities. *Cyperous rotundous*, *Phragmites karka* and *Typha angustifolia* are used by the local communities for thatching as well as for pulp, fibre and other uses. Several aquatic plants are used as green compost and manure apart from their usage as food for fish and water purification.



**Figure 17. Distribution of noteworthy plant species found in East Kolkata Wetlands**

### 2.4.2 Amphibia and Reptiles


ZSI has listed the presence of 10 amphibians and 29 reptiles from EKW (EKWMA, 2020). The reptiles include 9 lizard, 1 turtle, and 19 snake species. The invertebrates as well as lower vertebrates including fish and amphibian provide food to the reptilian species. A survey carried out between 1980 –1995 indicated occurrence of 19 reptilian species overall including 2 species of snake, 2 of monitor lizard, 3 of common lizard and 1 species of fresh water tortoise.

### 2.4.3 Birds

In 2016, the EKWMA published a compilation of 267 bird species recorded in EKW since 1964 (EKWMA 2016). The list includes 93 waterbirds of which 36 are migratory. Zoological Survey of India carried out one of the first surveys of birds in EKW during 1964 - 1969, wherein 248 species were recorded. Of these, only 162 species have been variably noticed during last 30 years. Moreover, at least 16 species have not been recorded after 1978-83 survey. Prakriti Samsad recorded 123 species of birds from Salt Lakes during 1978-83. Some of the land birds, bush lark, red-winged bush lark, franklin's wren warbler have been recorded during this survey, but never before or after. It is likely that the reclamation of marshes and changes inflicted on aquatic and other species resulted in absence of larger species of birds like the Openbill Stork, Spoonbill and many other species of ducks and teals including Red crested Pochard, Tufted Pochard, Baer's Pochard, Brahminy Duck, Comb Duck, Bar-headed Goose etc. Additionally, the birds of prey like Brahminy Kite, Pallas's Fishing Eagle, Osprey and Laggar Falcon (and now Vultures) which used to be common in recent past are now no longer seen. The observation from annual bird population estimation under Asian Waterbird Census in Nalban bheri shows that still a handful of Gadwals, Garganeys, Snipes, Gulls, Terns, Egrets and Cormorants also occur throughout the year though showing declining trends in population.



Figure 18. Bird count and species reported from Nalban Bheri



Picture 8. Black-winged Stilt in East Kolkata Wetlands (Credit: Kaushik Mukhopadhyay)

#### 2.4.4 Mammals

ZSI has enlisted 13 mammal species from EKW. Marsh mongoose *Herpestes auro-punctatus* is endemic to the region and also included in the schedule II of Indian Wildlife Protection Act, 1972 (Mallick 2009). It borrows along the slopes of ponds and feeds on fishes and aquatic snails within the wetland.



**Picture 9. A Marsh mongoose in East Kolkata Wetlands**

#### 2.4.5 Fish

The commercially important fin and shellfish in the EKW include 79 species of fish (Chandra, Raghunathan and Mao 2020), 11 species of prawns, 3 species of crabs and 20 species of molluscs (Mahapatra 2015). Among the fish species, 17 are culture species and 41 are wild species.

Traditionally, aquaculture in the EKW was predominantly of indigenous Indian Major Carps (IMC), Rohu (*Labeo rohita*), Catla (*Catla catla*) and Mrigal (*Cirrhinus mrigala*), and sometimes minor carp Bata (*Labeo bata*). In the 60s, exotic fish species were introduced, which included Common carp (*Cyprinus carpio*), Silver carp (*Hypophthalmichthys molitrix*), Grass carp (*Ctenopharyngodon idella*) and Mozambique tilapia (*Oreochromis mossambicus*). The Nile tilapia (*Oreochromis niloticus*) has replaced Mozambique tilapia in recent times. In limited quantities, Walking catfish (*Clarias batrachus*) and striped catfish (*Pangasianodon hypophthalmus*) are also produced, and attempts made to introduce Seabass (*Lates calcarifer*) enhance farm incomes (Bunting, Pretty, and Edwards 2010). 27 ornamental fish species of 21 families have also been recorded from EKW (Mahapatra and Lakra 2014).

Previously the fish fauna in the EKW system comprised both brackish water and fresh water forms. (Biswas 1969) reported occurrence of 80 species of fishes from the Salt Lake. The low-lying region with saltwater lakes acting as spill reservoirs for the Bidyadhari were utilized for farming of brackish water fish such as Bhetki (*Lates calcarifer*), Parse (*Mugil parsia*), Bhangar (*Mugil tada*) and Prawns (*Macrobrachium rosenbergii*), etc. (Ghosh and Sen, 1987). (De et al., 1989), has reported occurrence of 40 fish species from EKW. CIFRI has observed entry of 4 exotic species (Crocodile fish) in EKW in recent years. Exotics *Clarius guriepinus* and *Pangasius sutchi* have also been recorded from the aqua culture farm.

Indigenous species *Nandous nandus* and *Xenentodon cancila* which were abundant during 1980's have not been recorded at present. The rapid spread and population increase of suckermouth armoured catfishes belonging to the genus *Pterygoplichthys* (*Loricariidae*) in EKW in recent times is of increasing concern, because of the notable possibility that these non-native catfishes are adversely affecting fish germplasm and commercial fishery of this unique ecosystem (Hussan et al. 2019).

## 2.5 Livelihoods and Resource Linkages

### 2.5.1 Demographic features

The EKW has 37 mouzas or revenue villages (Map 15), covering the districts of North 24 Parganas and South 24 Parganas Districts. The EKW area falls under two Municipal Corporations and seven Gram Panchayats. In South 24 Parganas, there are seven Gram Panchayats – Beonta I, Beonta II, Bamanghata, Tardaha, Kheyadaha I, Kheyadaha II, and Pratapnagar – that cover a majority of these mouzas. The majority of the wetlands area in North 24 Parganas comes under the jurisdiction of the Ward 35 and 36 of the Bidhannagar Municipal Corporation. There are six mouzas within the Kolkata Municipal Corporation jurisdiction (Chakraborty and Gupta 2019).


The human settlements fringe the fish farms and are located mostly towards south of the DWF channel or towards the eastern parts of the wetland. Settlements on the western periphery adjoining the Eastern Metropolitan Bypass Road are comparatively larger as compared to those on the east. The total population of the 37 villages as per 2011 census was 0.15 million.

The overall population density of the wetland is 866 people per sq.km. as per 2011 census. Scheduled caste and tribes form 83% of the population. Livelihoods of the wetland communities are distinctly linked to wetland resources, with 74% of the working population drawing sustenance through engagement in fish farming, agriculture and horticulture. The rest of the population seeks livelihoods through seeking employment within the metropolis and its associated areas.

Assessment of trends in population growth rates indicates a continuous increase since 1951 with a marginal decline during the decade 1981-91. This trend broadly is commensurate with the overall trend of stabilization of population within Kolkata City, and rapid expansion of sub urban and peri-urban interfaces of the city, including the population within the wetland.



**Picture 10. Life in East Kolkata Wetlands is closely linked to wetland health**



**Figure 19. Population Density and Growth in East Kolkata Wetlands**





**Map 15. Mouza within East Kolkata Wetlands**


For the purpose of management planning, a household survey of 30 respondents was undertaken covering a range of stakeholders to assess demographic trends, map their perceptions of wetland condition and their recommendations for wetland management. It is true that the initiative cannot be even termed as a survey due to sheer limitation of the number of respondents. This was more like a dip stick study which helped the authority gauge the perception of different stakeholders associated with the wetlands which could be built into the Management Action Plan. The limited survey was done due to restrictions imposed by CoVID-19 pandemic. Despite its limitations, the study threw significant light on the perception of the communities on the wetland and its use.

The assessment indicates fish farming to be the primary source of income for 50% of the respondents, followed by agriculture and rag picking (Table 1).

**Table 1. Perception ranking of responsibility for implementing wetland management measures**


Occupation	Stakeholders					
	Agriculture farmers	Big Fish Farmers (area > 10 ha)	Cooperative fishers	Small Fish Farmers (area < 10 ha)	Vegetable farmers	Waste-collectors
Fish farming	7%	100%	100%	100%		
Agriculture	100%		20%	40%		
Vegetable farming					100%	
Small Shopkeeping						
Labour	100%					
Government Service						
Private Service						
Others					20%	100%

Access to social amenities within the wetland communities is limited leading to lower quality of life. The survey data shows very low levels of literacy amongst the respondents with 23% of the respondents illiterate and 33% have attended class till V.



**Figure 20. Education level of communities living in East Kolkata Wetlands**

The survey also highlights inadequacy amongst stakeholders in terms of assets owned and access to social amenities. As per the survey, 59% of the families have kachha houses and 30% of the families do not have separate bathrooms and cooking areas. Most striking disparities exist between rag pickers and the small bheri workers in terms of assets. Of all the respondents 94% of land ownership within EKW is of big bheri owners.



**Figure 21. Possession of assets by communities living in East Kolkata Wetlands**

### 2.5.2 Resource linkages

Ecosystem services, the benefits people receive from wetlands, provide the foundation of assessing wetlands-livelihoods inter linkages. Direct dependence on EKW is in the form of fish farming, agriculture and horticulture.



**Picture 11. Sewage fed fish farming has evolved traditionally in East Kolkata Wetlands**

### 2.5.2.1 Sewage fed fish farming

#### The traditional technology

Sewage fed fisheries form the dominant resource use within EKW. Over 250 fishponds (locally called bheri) receive sewage for culturing fish, and in the process remove nutrients and other heavy metals rendering wastewater treatment to the City of Kolkata. The practise has been continuing since the 1940s after an enterprising fish farmer discovered the virtue of using sewage water for fish farming.

Dr Dhrubajyoti Ghosh, who is credited with bringing national and international attention to this waste recycling system has described the ecological processes underpinning this fish farming practice in his book „Ecology and Traditional Wetland Practice“ (Ghosh 2005). The process is paraphrased in the following paragraphs.

The network of pumps brings in the sewage from the Kolkata City to the open channels called the Dry Weather Flow channels of the Kolkata Drainage Outfall System. A regulator at Bantala raises the level of sewage so that further movement to fish ponds can be ensured by a natural hydraulic gradient. The drainage network consisting an intricate web of primary, secondary, and tertiary canals was engineered in such a way that it could distribute wastewater using gravity to the majority of fisheries.

The fish farms receive sewage in wastewater stabilisation pond. These shallow ponds create more favourable conditions for photosynthesis and full vertical circulation of water to the surface where algal blooms occur. The algae–bacteria symbiosis allows for efficient BOD and pathogen/faecal coliform reduction. The sunlight supports photosynthesis and the rapid growth of plankton, which helps degrade organic matter. Water hyacinth plays a special role in the functioning of this complex ecosystem by leaching out heavy metal ions from the surrounding water, with its dense root system acting as biofilters for the passive remediation of wastewater. The overgrowth of plankton is controlled by fish which grazes the plankton.

Once every three to four years in January, the ponds are allowed to dry out in rotation through natural evaporation; the embankments are repaired and vegetable is grown on the dry bed. After harvesting the vegetables, the fishers dig silt traps, wherein the deposited silt is repairing banks. Just prior to the release of the sewage into the bheri, the dry ground is tilled, and lime is scattered over it to decrease the acidity of the pond waters, making them more suitable for the fish.

The fish species used for aquaculture occupy different ecological niches of the pond system. When the nutrient-rich effluent moves through the system, it is progressively cleaned, and nutrients are redirected to the growth of algae or agricultural products grown along the pond edges. The natural water purification saves the Kolkata nearly Rs. 4,680 million annually (Dey and Banerjee 2018).



**Picture 12. A bamboo screen filters solid waste from sewage before being brought into the fish farm**

#### *Fish production and productivity*

The last detailed assessment of the number of bheri in EKW was done close to 2000, wherein approximately 260 numbers were counted. The size of bheri ranged roughly between 5 and 50 ha. Smaller fisheries under 2 ha exist mainly in Sonarpur and Tiljala, while large fisheries over 40 ha predominate in Bidhannagar; Bhangar is characterised by a mixture of small to medium-sized fisheries. Over 80 per cent of the bheri are under 20 ha in size, occupying approximately 30–40% of the total fishery area. The two state-owned fisheries, Nalban and Goltala, are the largest, comprising about 165 ha and 115 ha, respectively.

Fish yield in the EKW is two–four times higher than that of ordinary fish ponds. The yield is positively correlated to the area of the fish pond, increasing up to 6.5 MT where the effective area of the water body was above 70 ha (Ghosh 1999). The average annual fish production from the EKW during 2015-2020 (as per records of the Fisheries Department) has been estimated to be 22,000 MT. This compares well to the production of around 18,000 MT reported in 2002 (Bunting, S W, Kundu, N, and Mukherjee, M 2002; Bunting, Pretty, and Edwards 2010).

The location of a bheri, i.e., its distance from the DWF canal is also a key variable determining the productivity and profit returns. Usually, a bheri located further from the DWF canal and smaller in size have a lower profitability as compared with those which are larger and located closer to the main canal.



**Picture 13. Nutrient in sewage is directed to vegetation growth and fish**

#### *Fisheries management*

The bheries are under three forms of ownership: a) government; b) cooperatives, and c) private. Bheri ownership by cooperatives has gradually reduced (from 27% to 6% during 2002-15) (Mukherjee 2020). Private fish pond owners rarely participate directly in fish growing but have leased out to leaseholders, who basically concentrate on how to minimise the cost of production. Many worker-run non-registered cooperatives have also leased out to leaseholders due to inability to generate capital (Chakraborty and Gupta 2019).

Fish is traded through seven auction markets: Chingrighata, Kestopur, Bamanghata, Bantala, Chowbaga, Garia, and Gangajoara. Chingrighata is the biggest market with an average daily sale of about 13 MT.

A multi-stakeholder analysis for EKW fisheries conducted by Forum for Policy Dialogue on Water Conflicts in India (Chakraborty and Gupta 2019) listed out the following issues related to the management of EKW fisheries:

- Cooperatives are apprehensive of the imminent threats of corporatization of cooperatives since the Fisheries Department has opted for the tender route of awarding license in recent years.
- The decline in sewage quantity and quality affects the fisheries as well as of the fishers.
- De-siltation of the fish ponds along with re-excavation of the canal distribution network

#### *Food safety and health concerns*

The safety of fish produced for human consumption from EKW has been questioned in recent times, as the sewage quality has changed. The 2019 CPCB report to the National Wetlands Committee (CPCB 2019) highlights the transformed character of sewage in its report as thus:

“We are no longer speaking of the raw sewage of the kind that entered the East Kolkata Wetlands when the bheries were established in 1920-30. Today industrial effluent makes up about 20% of Kolkata's daily total liquid waste and most of the untreated liquid mixes with Kolkata's stormwater and sewage, industrial runoff and household liquids are all mixed and end up in multiple canals that drain the city. The sewage that enters the Wetlands is a composite biodegradable and toxic, non-biodegradable matter, including a range of heavy metals.”

Monitoring records of WBPCB do not indicate presence of heavy metals in any of the sampling sites, However, heavy metal contamination in sewage has been indicated by several recent studies (Vicziány, Chattopadhyay, and Bhattacharyya 2017; Nandi et al., 2013; Sarkar et al., 2011; Joystu 2017). Though, there have been arguments that heavy metal accumulation was within safe levels within the fish muscles (the part that is generally eaten), and cooking at high temperatures reduces the concentrations, this fact has been contested (Nath and Bhounik 2013) and close monitoring of bioaccumulation recommended (Kumar et al., 2010). Table 2 reproduces data on heavy metal concentrations in select fish species of EKW from CPCB 2019.

Epidemiological evidences suggest that the communities living within the wetland are subjected to various kinds of diseases. Workers employed on the fish farms experience a range of health problems predominantly diarrhoea and those related to nutritional deficiencies revealed by the presence of anaemia, oedema, night blindness, and bleeding gums.

*Vibrio parahaemolyticus*, one of the most important diarrhoea-causing agent (after *V. cholerae*) in the Kolkata area, has been found in the intestines of fish from EKW (Das and Mandal 2018) suggesting why there is a high rate of diarrhoea amongst the workers.

**Table 2. Concentration of heavy metals (mg/kg) for select fish samples**

Fish species	Muscle	Liver	Gill
<i>Tilapia Nilotica</i>	Pb: 0.52 – 7.91 Cd: 0.07 – 0.75 Cr: 41.02 – 65.29	Pb: 0 – 7.93 Cd: 0.14 – 5.18 Cr: 0 – 33.09	Pb: 2.64 – 14.77 Cd: 0.06 – 1.08 Cr: 3.69 – 57.35
<i>Tilapia Mosambica</i>	Pb: 0 – 4.52 Cd: 0.04 – 0.09 Cr: 0 – 4.55	Pb: 0 – 4.66 Cd: 0.24 – 4.36 Cr: 16.82 – 43.54	Pb: 3.76 – 18.19 Cd: 0.08 – 1.19 Cr: 0 – 28.56
<i>Labeo Rohita</i>	Pb: 0 – 7.38 Cd: 0.05 – 0.67 Cr: 0 – 63.29	Pb: 0 – 6.63 Cd: 0.16 – 0.2 Cr: 0 – 29.9	Pb: 6.94 – 26.50 Cd: 0.2 – 0.21 Cr: 0 – 42.93

(WHO and FAO permissible levels in fish tissue: Pb = 0.5 mg/kg; Cd = 0.5 mg/kg and Cr = 0.15 mg/kg)

Industrial effluents within the vicinity of EKW include the Bantala Leather Complex, which is located outside the boundaries of the wetland, and several old tanneries located in the eastern part of Kolkata Municipal Corporation. The Bantala Leather Complex was a response to Supreme Court order of 1992 to relocate the over 500 tanneries that had come about in Topsia and Tangra stretches of the municipality. However, leather shavings dumped along the EKW are a cause for worry and need to be monitored regularly.

#### 2.5.2.2 Agriculture

Rice is cultivated using effluent water from the bheri, covering an area of 2637 ha. Rice cultivation is one of the major livelihoods within the wetland followed by cultivation of vegetables. In a study by Mukherjee et al 2013 efforts were made compare rice cultivation from untreated sewage water from the bheri and rice cultivation using ground water. The study concludes that rice harvest is less profitable when ground water is used.

Labour employed within the paddy fields are employed on a casual basis and therefore lack security of regular income. Labourers working for rice farmers usually have their own farms, yet work on the fields of larger farmers to earn extra incomes.





**Picture 14. Agriculture forms the terminal use of sewage in East Kolkata Wetlands**

### 2.5.2.3 Horticulture

Approximately 244 ha of agricultural farms are located within the EKW, and particularly in and around the Dhapa landfill area. There are around 3000 farm plots in Dhapa, ranging between 0.03 and 0.2 ha, employing over 4000 people (Mukherjee and Ghosh 2015). Tradition of organic waste reuse forms the base of horticulture concentrated within Dhapa landfill areas. Organic waste makes the soil conditions extremely fertile and it is a common practice for farmers to crop 3-5 different varieties of vegetables on the same land. 55% area is used for double-cropping, while the rest is used for three crops in a year (Kundu & Chakraborty 2017, Mukherjee 2020). Almost 15 different types of vegetable are grown in EKW; production is estimated at 150 Metric Tonnes annually (Mukherjee 2020).



**Picture 15. Vegetable cultivation in Dhapa region**

Vegetable production from EKW meets nearly 20% of the vegetable requirement of the Kolkata city. Proximity to the city, availability of water makes horticulture a household activity in EKW, farmers even rent small plots for raising vegetables (Kundu and Chakraborty 2017). Labour employment within the horticulture sector is on a casual basis depending upon the availability from the manager's own household and workload; during busy periods of planting and harvesting more labour is likely to be employed. There is also a tendency of seeking employment into neighbouring farms for extra incomes.

As in fisheries, concerns have been raised on the levels of heavy metal accumulation in the vegetables grown in EKW. Table 3 presents data from a 2019 report of CPCB indicating that such risks do exist.

**Table 3. Heavy metal concentrations in select vegetable samples of East Kolkata Wetlands**

Vegetable	Heavy metal concentration (mg/kg)
Brinjal ( <i>Solanum melongena</i> )	Pb: 6.32 – 10.2 Cd: 0.175-0.176 Cr: 23.89-26.62 Hg: BDL
Lal Shak (Root) ( <i>Amaranthus cruentus</i> )	Pb: 17.16-89.82 Cd: 0.35-0.36 Cr: 35.22-67.73 Hg: 0-0.021
Lal Shak (Leaves) ( <i>Amaranthus cruentus</i> )	Pb: 11.47-60.23 Cd: 0.175-0.176 Cr: 14.9-16.74 Hg: 0-0.011
Cauliflower ( <i>Brassica oleraceae</i> )	Pb: 5-10.96 Cd: 0.175 Cr: 7.52-13.33 Hg: BDL
Maize ( <i>Zea mays</i> )	Pb: 0-5.63 Cd: 0.174-0.176 Cr: 0-9.26 Hg: BDL

(WHO and FAO permissible levels in vegetables: Pb = 5 mg/kg; Cd = 0.1 mg/kg; Cr = 20 mg/kg and Hg = 0.03 mg/kg)

#### 2.5.2.4 Waste collectors

The Dhapa in EKW was initially conceived as garbage-dumping ground. Kolkata generates 1.6 million tonnes of municipal solid waste annually and 4500 tonnes a day, which is entirely dumped into Dhapa dumping area within EKW.

Previously Dhapa site employed over 25000 people, waste collectors looking for objects that could be re-recycled, collecting approximately 10% of the total solid waste.

### 2.5.3 Community perception on wetlands conditions

There is a general perception that the health of EKW needs to be protected (Figure 22). Nearly half of the respondents (47%) indicated that EKW condition is on the decline. Reduced inflow of sewage in EKW and siltation of bheri over the last 10 years were indicated as significant trends. Instances of diseases amongst family members have increased with 50% respondents agreeing. Respondents perceptions whether pressure of builders and realtors have increased in EKW in last 10 years seem to vary with 23% strongly agreeing and 27% agreeing and 23% disagreeing.



**Picture 16. Community participation during survey on EKW**




Figure 22. Community's perception score (6 to -6) of wetlands condition in the past decade

Perceptions of different occupation groups seem to vary when it comes to the condition of EKW over the last ten years (Figure 23). Agriculturists and waste collectors seem to disagree that the area of bheri has decreased over the last ten years whereas respondents belonging to the groups of big bheri and fish cooperative seem to agree that bheri areas have decreased over the last 10 years.

Agriculture, big and small bheri groups seem to agree that diseases in fish may have increased over the last ten years. Apart from horticulture groups all the others state an increase in silt in the bheri. All the occupation groups do not strongly agree over the increase in agriculture and horticulture production. A majority of agriculture and big bheri groups agree that

agriculture production has increased over the last ten years, whereas rest of the groups disagree over the production increase of agriculture. All groups other than agriculturists and horticulturists perceive the threat of realtors has increased over the last 10 years.





Figure 23. Community's perception score (6 to -6) of threat to wetlands

#### 2.5.4 Community recommendations for wetland management

Improving income generation activities within EKW and provision of toilets and drinking water were the topmost choices for restoration measures followed by desiltation of bheri and channels (Figure 24). Improving awareness and stopping industrial discharge into the sewage were the other most sought out options for wetland management followed by improving tourism facilities within and strict enforcement of the EKW Act.


Community recommendations for wetland management vary across the stakeholder groups with a big bheri, fishing cooperatives and small bheri groups all choosing the options of stopping the Dhapa landfill. Small bheri groups seem to prefer reduction in fishing followed by stopping the Dhapa landfill and preventing encroachment from builders. The perceived threat

from builders to small bheri operators and fish cooperatives seem larger compared with horticulture and agriculture groups. Agriculturists and waste collectors' top most choice for wetland management is the enforcement of the EKW Act. Stopping of industrial discharge in sewage were preferred by all groups in varying degree except the waste collectors.



**Figure 24. Stakeholder responses on agencies responsible for implementing various interventions in East Kolkata Wetlands**

The respondents rank responsibility of Government of West Bengal as the highest for implementing the recommended measures followed by EKWMA and the central government (Figure 25). Implementation of measures such as desiltation of bheri and channels, income generation activities, provision of toilets and drinking water were all ranked to be implemented by Govt of West Bengal followed by EKWMA.



**Figure 25. Perception ranking of responsibility for implementing wetland management measures**

### 3 Evaluation of Wetland Features

---

India, as a signatory of the Ramsar Convention, is committed to achieving wise use of all wetlands in her territory. Wise use of wetlands is defined in the text of Ramsar Convention as 'the maintenance of their ecological character, achieved through the implementation of ecosystem approaches, within the context of sustainable development'. Ecological character is 'the combination of ecosystem components<sup>2</sup>, processes<sup>3</sup> and services<sup>4</sup> that characterize the wetland at any given point in time.' Ecosystem Management of wetlands thus seeks to achieve the goal of 'maintenance of ecological character' or 'wetland wise use'.

Though sounding counter-intuitive, wise use as a wetland management approach is much wider than use of a wetland. The phrase 'in the context of sustainable development' recognizes that development, which may be inevitable in some cases, is not an objective for every wetland. Wherever development is to take place, it has to be facilitated in sustainable ways using approaches elaborated in the Convention (managing at basin scale, mainstreaming in urban developmental planning, participatory management amongst others). "Ecosystem approaches" include the elements elaborated by the Convention on Biological Diversity – integrated management, stakeholders' participation in decision-making, transparency about trade-offs, and equitability of the outcomes. In totality, wise use is about 'maintaining the capability of the wetland' to support human well-being at present and in future, rather than 'use' or 'development' at present.

---

<sup>2</sup>The living (biotic) and non-living (abiotic) constituents of wetland ecosystem. These include: Geomorphic setting (landscape, catchment, river basin); Climate (precipitation, wind, temperature, evaporation, humidity); Physical setting (area, boundaries, topography, shape, bathymetry, habitat type and connectivity); Water regime (inflow, outflow, balance, surface – groundwater interactions, inundation regime, tidal regime, quality); Wetland Soil (texture, chemical and biological properties); and Biota (Plant and animal communities)

<sup>3</sup> Processes that occur between organisms and within and between populations and communities, including interactions with non-living environment, that result in existing ecosystem state and bring about changes in ecosystems over time. These include: Physical processes (water stratification, mixing, sedimentation, erosion); Energy – nutrient dynamics (primary production, nutrient cycling, carbon cycling, decomposition, oxidation – reduction); Processes that maintain animal and plant population (recruitment, migration); and Species interaction (Competition, predation, succession, herbivory)

<sup>4</sup> Benefits obtained by humans from ecosystems, categorized as: Provisioning (fisheries, use of aquatic vegetation for economic propose, wetland agriculture, biochemical products); Regulating (maintenance of hydrological regimes) and Cultural (recreation and tourism, spiritual, scientific and educational value). Supporting services have been included in definition of ecosystem processes.



Changes to ecological character of wetlands outside natural variation may signal that uses of the site are unsustainable, and may lead to the breakdown of its ecological, biological and hydrological functioning (Ramsar Convention 1996, Resolution VI.1). Assessing and responding to risks of human induced adverse change in ecological character is therefore fundamental to achieving wise use of a wetland.

For wetland managers to be able to implement management that ensures maintenance of ecological character, it is important to identify and retain the site's essential ecological functions which underpin the wetland's ecosystem services and biodiversity. Implicit within this recommendation is the need to identify key elements of ecological character, maintaining which would constitute site's wise use. The extent to which ecological character is maintained and adverse human-induced changes prevented is reflected in these key features. Furthermore, a social-ecological systems perspective for defining and assessing ecological character enables consideration of the interactions social actors and institutions have with biophysical components of wetlands.

The Ramsar Convention's Guidelines for ecological character description are contained in Ramsar Resolution X.15. These elements have also been formally and systematically included in the 2015 revision of the Ramsar Information Sheet (RSIS) Format, which needs to be updated every six years.

This chapter of management plan presents an evaluation of ecological character of EKW, based on assessment of wetland features presented in the previous chapter. A description of ecological character is provided herein, highlighting key elements, maintenance of which may be the focus of management. Status and trends in the key elements are discussed next and followed by an analysis of threats and risks of adverse change in ecological character. The chapter concludes with a listing of the knowledge gaps.

### 3.1 Ecological Character Description

Located on the eastern fringes of Kolkata City and spanning 12,500 ha, East Kolkata Wetlands are a mosaic of landforms including predominantly water dominated areas (used as fish farms) to land centric usages for agriculture, horticulture and settlements. The existing wetland regime is a remnant of series of brackish wetlands connected to the freshwater as well as marine environments of the Gangetic Delta and the Bay of Bengal, in an ecological continuum with the Sundarbans.

Over 260 shallow fish ponds in the EKW receive over 900 MLD pre-settled sewage from the Kolkata Metropolitan region through a network of locally excavated secondary and tertiary canals, which is used to produce 20,000 MT of fish, 50,000 MT of vegetables and irrigate 4700 ha of paddy lands. As the nutrient-rich effluent moves through the system, it is progressively cleaned and nutrients are redirected to the growth of algae or agricultural products grown along the pond edges and agricultural lands. Algae and other aquatic plants are used to feed upto 17 species of fish cultured in these ponds, which in turn create nitrogen and phosphorus-rich water to irrigate the adjacent rice fields. The traditionally evolved natural water purification waste recovery practice saves the Kolkata City nearly Rs. 4,680 million annually in terms of treatment cost of upto 65% of the City's sewage. These wetlands also lock in over 60% of carbon from wastewater, thus reducing harmful Green House Gas emissions from the region.

The wetland is inhabited by diverse species. At least 120 plant, ten amphibians, 29 reptiles, 260 bird, 58 fish, 11 prawn, 3 mollusc, and 11 mammal species have been recorded from these wetlands. Marsh mongoose *Herpestes auro-punctatus* endemic to the region and also included in the schedule II of Indian Wildlife Protection Act, 1972.

The traditional waste recovery practice provides subsistence opportunities for a large, economically underprivileged population of 0.15 million living in over 37 mouzas within its boundaries. EKW is also one of the few natural habitats providing recreational avenues for the urban and peri-urban population.

East Kolkata Wetlands Management Authority is the nodal government agency mandated to ensure wise use of the wetland within the regulatory framework defined by the Wetlands (Conservation and Management) Rules, 2017 (notified under the Environment (Protection) Act, 1986) and the East Kolkata Wetlands (Conservation and Management) Act, 2006 and rules.

The ability of EKW to provide ecosystem services and sustain biological diversity is enabled by:

- a) Ingenious use of pre-settled sewage received from Kolkata City for resource recovery through aquaculture, horticulture and agriculture.
- b) Hydrological gradient which enables distribution of sewage to different parts of the wetland.
- c) Water-hyacinth mediated phytoremediation enabling wastewater treatment.
- d) Indigenous knowledge of the fish-farmers on management of fish ponds for cultivation of diverse species of fish.
- e) Diverse habitats which enable a range of plant and animal species to survive.
- f) Land use control as set under the provisions of East Kolkata (Conservation and Management) Act, 2006 and Wetlands (Conservation and Management) Rules, 2017.

### 3.2 Status and trends

Status and trends in key wetland features are presented in the Table below, with reference to their condition at the time of Ramsar Site designation in 2002.

**Table 4. Ecological Character Description**

<b>ECD descriptor</b>	<b>Current Condition (2018-2020)</b>	<b>Data source</b>	<b>Condition at time of Ramsar Site designation (2000 – 2002)</b>	<b>Data Source</b>	<b>Change description and evaluation (Adverse, Positive, Not Evaluated)</b>
<b>Wetland extent and regime</b>					
Area	12500 ha	Sentinel 2B May 2019, Sentinel 2A November 2019	12500 ha	Landsat 5 November 2000	No change
Shape	Amoeba like	Sentinel 2B May 2019, Sentinel 2A November 2019	Amoeba like		No change
Area under fish farms	6100 ha	Sentinel 2B May 2019, Sentinel 2A November 2019	4431.21 ha	Landsat 5 November 2000	Increased. Positive change

Area under agriculture	3,350 ha	Sentinel 2B May 2019, Sentinel 2A November 2019	4,300 ha	Landsat 5 November 2000	A decrease in area under agriculture may reduce the capability of the landscape to treat wastewater.  Adverse Change
Area under solid-waste dump	91 ha	Sentinel 2B May 2019, Sentinel 2A November 2019	49 ha	Landsat 5 <sup>TM</sup> November 2000	Almost doubled in 2 decades  Adverse Change
Settlement area inside wetlands	2400 ha	Sentinel 2B May 2019, Sentinel 2A November 2019	1560 ha	Landsat 5 <sup>TM</sup> November 2000	Increased by over 50 %.  An increase in settlements increases intensity of resource use as well as increases vulnerability of wetland communities.  Adverse Change
<b>Hydrological regimes</b>					
Rainfall	1601 mm	Indian Meteorological Department	1383 mm	Indian Meteorological Department	The rainfall has increased.  Not evaluated

Sewage inflow into wetland	Flow rate: ~ 900 MLD	CPCB, 2019	Flow rate: ~ 750 MLD	Ghosh, 2002	Positive change
Sewage distribution within different bhery and parts of wetlands	Fish farms in the southern use pumps to draw sewage				Not evaluated
Sedimentation	TSS: Inflows: 6-342 ppm Interim: 4-302 ppm Outflows: 8-462 ppm				Not evaluated
Sewage quality	<i>Inflows</i> BOD: 22 ppm (Average of 12 months)  COD: 66.5 ppm (Average of 12 months)  <i>Outflows</i> BOD: 11.8 ppm (Average of 12 months)  COD: 47.1 ppm (Average of 12 months)	WBPCB, 2019-20  Inflows: Bantala to Bananghata Jalpath  Interim: Kestopur canal after passing through the bhery.  Outflows: Ghushighata after CLC	<i>Inflows</i> BOD:23 ppm  COD:105 ppm  <i>Outflows:</i> BOD:13 ppm  COD: 80 ppm	Ghosh, 2002  Inflows: Bantala lock gate  Outflows: Ghusiaghata lock gate	No Change.

<b>Wetland Catchment</b>					
Population density within catchment	866 people per km <sup>2</sup>		859 people km <sup>2</sup>		No change.
Development of waste treatment infrastructure within catchment	179 MLD	KMC Drainage Report 2020	120 MLD	Dey and Banerjee 2017	Impact not evaluated
Land use and land cover in the region adjoining the wetland	Surrounded by permanent settlements, built up areas dominant, especially to the north and west.		59% of the area is agriculture followed by settlements.	MAP 2008	Built-up area around the wetland has increased, thus increasing runoff.  Impact not evaluated
Species and Habitats	120 species 37 families of plants 10 amphibians 29 reptiles	ZSI			No baseline
Number of water-birds	267 bird species recorded since 1964, 90 aquatic 11 semi aquatic	EKWMA			Adverse change as nearly 30 species not sighted in recent times
<b>Livelihood systems</b>					
Number of people dependent on fisheries	20, 000	Mandal and Jana 2018	12, 000	MAP, 2008	Impact not evaluated


Number of people dependent on horticulture	4000	Mukherjee and Ghosh 2015	2640	Chakraborty and Gupta 2019	Impact not evaluated
Number of people engaged in waste collection and sorting	25,000	Mukherjee and Ghosh 2015	5000 people (2010)	Chakraborty and Gupta 2019	Impact not evaluated
Fish production	20,000 MT	Fisheries Department	18,000 MT	Bunting et al, 2010	No change
Vegetable production	50,000 Metric Tonnes annually, 150 MT per day	Kundu and Chakraborty 2017			Not assessed
Fish quality	Presence of heavy metals indicated	CPCB, 2019			Bioaccumulation may have increased health risks. Final report yet to come. Not evaluated

Health of wetland communities	<p>Antibiotic residue and microbial counts are reported to be less than the permissible limit. Risks due to heavy metal contamination</p> <p>Skin problems of fish producers or catchers or other labourers who work in sewage ponds</p> <p>Fever, cholera, diarrhoea</p> <p>Edema of eyelids, tumour, congestion of nasal mucous membranes and pharynx, stuffiness of the head, and gastrointestinal, muscular, reproductive, neurological, and genetic malfunctions</p>	<p>Vicziany et al., 2017</p> <p>Jana et al., 2018</p> <p>Das and Mandal 2018</p> <p>WBPCB 2014</p>			Not evaluated
<b>Institutions and Governance regimes</b>					
Implementation of regulatory regimes	Over 350 violations of East Kolkata Wetlands (Conservation and Management) Act, 2006 recorded.				Adverse Change



### 3.3 Risk of adverse change in ecological character

Based on the analysis of status and trends, following factors of adverse change in ecological character have been identified. The management needs to address these risks through specific interventions.



**Figure 26. Major risks assessed in East Kolkata Wetlands**

**Table 5. Threats and associated impacts on East Kolkata Wetlands**

Threats	Impacts (known / likely)
Encroachments	<ul style="list-style-type: none"> <li>• Reduced waste treatment capability</li> <li>• Increased pressure on species habitats</li> <li>• Increased vulnerability of assets</li> </ul>
Heavy metal contamination	<ul style="list-style-type: none"> <li>• Health risks due to bioaccumulation</li> </ul>
<p><b>Solid waste dumping</b></p> <p><b>A few plastic reprocessing units have cropped up along the Bantala, Basanti High Way Road side areas, in violation of the Plastic Waste Management Rules, 2016 and the Plastic Waste Management (Amendment) Rules, 2018. These illegal units emit toxic plumes, and create health risks for the communities.</b></p>	<p>Reduced waste assimilation capacity of the wetland</p> <p>Adverse impacts on health of wetland communities</p>
<p><b>Land subsidence</b></p> <p><b>The area around Salt Lake and Bantala is reported to have high rates of subsidence which can alter the topography and natural hydraulics of the wetland.</b></p>	<p>Altered topography may adversely impact natural hydrology</p>
<p><b>Sedimentation of channels</b></p> <p><b>Fish farms located away from Dry Weather Flow channel are not flushed naturally and have relatively higher rates of sedimentation. Several fish farms in the Southern part of the wetland need to use electrical pumps to draw sewage.</b></p>	<p>Reduced access to sewage increases energy usage for pumping, reducing overall efficiency of waste water treatment</p>
<p><b>Invasive species</b></p> <p><b>The rapid spread and population increase of suckermouth armoured catfishes belonging to the genus <i>Pterygoplichthys</i> (<i>Loricariidae</i>) in EKW in recent times is of increasing concern, because of the notable possibility that these non-native catfishes may adversely affect fish germplasm and commercial fishery of this unique ecosystem.</b></p>	<p>Adverse impacts on wetlands biota.</p>

<p>Climate Change impacts</p> <p>Wetland communities are highly vulnerable to impacts of climate change, including risk of high floods and increase in temperature.</p>	<p>Increased vulnerability of wetland communities</p>
---	---

### 3.4 Key knowledge gaps

The current monitoring of wetland is limited to assessing land-use change through remote sensing, select water quality parameters, and fish production. Some of the major knowledge gaps include:

- a) Climate vulnerability of wetland and risk mitigation options
- b) Impacts of heavy metals on food production systems and consequent health risks
- c) Impacts of solid waste dumping and plastic waste on wetland environment

## 4. Institutional Arrangements

---

The wise use principle of wetland management recognises that addressing wetland degradation requires incorporation of linkages human societies have with wetlands within all spheres of management. Institutions and governance systems structure the relationship human societies have with wetlands, and thus, have an important bearing on wise use outcomes.

Institutions encompass all formal interactions among stakeholders and social structures that determine how decisions are taken and implemented, how power is exercised, and how responsibilities are distributed. Various collections of institutions come together to form governance systems, which include interactions between different centres of power in the society at different scales from local to global. Institutions and governance systems determine, to various degrees, the access to, and the control, allocation and distribution of components of nature and anthropogenic assets and benefits to people. Formal institutions are often embedded in informal institutions, which collectively define the overall institutional environment. Most importantly, institutions and governance influence the direct and indirect drivers of change in a wetland ecosystem. The degree of fit of institutions and governance systems with functioning of EKW at basin scale is one of the key determinants of wise use.

Wetlands are open systems and subject to the influence of a range of developmental planning and resources management decisions being implemented within their river basins and beyond. An important institutional pre-requisite is to ensure arrangements for cross-sectoral coordination, so that various programmes pursued by the different government as well as private agencies do not work for cross-purposes, generating adverse outcomes for wetlands. Institutions also need to be adaptable, to be able to accommodate new information and perspectives on wetlands, needs of diverse stakeholder groups, and an uncertain political environment. Since communities living in and around a wetland have a direct bearing on the ecological state, and often have nuanced management systems based on traditional knowledge, institutional arrangements need to have adequate representation of such communities and knowledge systems. The institutions also need an adequate regulatory backing to be able to enforce measures required to prevent detrimental activities around wetlands.

Institutional requirements for conservation and sustainable management of the EKW is defined by the ability to ensure integration of site management within broad scale environmental and developmental programming (at river basin scale), and enabling stakeholder participation. This section of the management plan contains an analysis of institutional arrangements in the context of integrated management of EKW.

### 4.1 Existing Institutional and Governance Arrangements

#### 4.1.1 Government organizations

*East Kolkata Wetlands Management Authority* is the nodal agency constituted as per the provisions under the East Kolkata Wetlands (Conservation and Management) Act, 2006, most recently amended in March 2017 through The East Kolkata Wetlands (Conservation and Management) (Amendment) Act, 2017. The authority is constituted with the Secretaries of different state government departments (Environment, Land and Land Reforms and Refugee Relief and Rehabilitation, Fisheries, Forests, Panchayat and Rural Development, Irrigation and Waterways

and Tourism) under the chairmanship of Minister-in-Charge, Environment Department of Government of West Bengal. Chief Secretary to Government of West Bengal is the Vice-Chairperson of the EKWMA. Secretary, Environment Department is the Member Secretary of the Authority. The Authority also includes four experts drawn from the fields of wetland ecology, hydrology, fisheries and socioeconomics. The authority has been entrusted with a range of responsibilities including:

- Demarcation of wetland boundaries
- Prevention of unauthorized development projects or unauthorized use of wetland area
- Enforcement of land use control and prevention, prohibition or regulation of developmental activities
- Management of wetland based on conservation principles and guided by management plan prepared in conformity with Ramsar guidelines
- Pollution abatement and conservation of flora and fauna
- Research on various aspects of wetlands
- Communication and outreach on values and functions of wetlands
- Networking with other Ramsar sites

The office of the Authority is housed within the Department of Environment and is headed by the Member Secretary. The Authority currently has 17 staff, functioning under the Chief Technical Officer (Table 6). Of these, three (including the Chief Technical Officer) are employees of Government of West Bengal and the rest are employed on ad-hoc basis.

**Table 6. Staff strength in East Kolkata Wetlands Management Authority**

Division	Number of officers	Function
Legal	1 Senior Law Officer *(@) 1 Technical Officer (Law) 1 Technical Assistant	Address all legal matters resulting from enforcement of EKWCM Act
Monitoring and Research	1 Technical Officer* (@) 1 Scientific Officer 2 Field Supervisors 5 Technical Assistants	GIS and Remote Sensing Environmental monitoring Scientific and technical documentation
Accounts	2 Technical Assistants	Maintenance of accounts and preparation of financial statement of the Authority
Office Support	3 peons	Office assistance

(\* = head of the division, @ = permanent employee of Government of West Bengal)



Picture 17. Demolition of illegal structures in East Kolkata Wetlands is a major task of the Authority

Police authorities keep vigil on any unauthorised activities in the EKW, submit charge sheet within stipulated time and provide security in case of violation of law and order situation. The Authority also receives support from the District Administration, Municipal Corporations, PWD and other Government Departments and agencies in the implementation of various activities.

Department of Environment is overall responsible for protection of the EKW as EKWMA is set up under this Department. West Bengal Pollution Control Board (WBPCB), Institute of Environmental Studies and Wetland Management (erstwhile IWMED) and West Bengal Biodiversity Board (WBBB) (three parastatal organisations within the Department) are responsible for research and monitoring activities. The Department has also constituted the West Bengal State Wetlands Authority as the nodal institutions for policy, regulation and management of wetlands at the state level.

Department of Land & Land Reforms has the authority for land transfer and conversion and has a pivotal role in management of land use in the EKW area. As per the operative legislation, all applications for conversion of land within the EKW are scrutinized by the EKWMA and based on their recommendations, such proposals are sanctioned by the Land & Land Reforms Department in exercise of powers under section 4C of The West Bengal Land Reforms Act (1955, most recently amended in 2017).

Department of Irrigation & Waterways manages the main drainage channels in EKW area namely dry weather flow (DWF), Storm Weather Flow (SWF) and Fishery Feeding canal (FFC). Fisheries get sewage from the intrinsic network of canals originating through these channels. This department also has the responsibility for flow of sewage, regulation of lock gates at Bantala and construction of lock gates and pumping stations.

Department of Fisheries is responsible for the production and improvement of the sewage fed fisheries in this area. The Department also operates two fisheries at Nalban and Goltala, and oversees functioning of the fishery co-operatives.

Department of Forest is responsible for conservation and maintenance of various kinds of trees /plantation grown in non-forest areas included in EKW.

Department of Panchayats & Rural Development is focused on the non-urban areas which include parts of the EKW, and are authorized to sanction the building plans under The West Bengal Panchayat Act, 1973 as amended from time to time. According to the order of the Panchayat Department dated 04.09.2014 "no Panchayat body shall accord approval to any building plan within the East Kolkata Wetlands area except in strict compliance with the provision of the East Kolkata Wetlands (Conservation and Management) Act, 2006 and the relevant rules under the said Act". The district planning committees are also empowered to co-ordinate development activities in parts of wetlands under their control. Gram Panchayat (GP), Panchayat Samity and Zilla Parishad (ZP) are elected tiers in rural areas. GPs are amalgamations of 10-15 villages, Panchayat Samitis are coterminous with the area of the blocks and ZPs consist of a number of Panchayat Samity. GPs and other tiers of panchyati-raj system have a critical role at the lowest levels of govt. in the WRR and are direct representatives of the residents of the WRR.

The Urban Development and Municipal Affairs Department sanctions building plans (including those for public amenities) within the urban areas of the EKW under Kolkata Municipal Corporation (KMC) and Bidhannagar Municipal Corporation (BMC). KMC is authorised to sanction building plan for its areas in the EKW. However NOC from the EKWMA is necessary according to the KMC order no. 18 of 2013-14 dated 31.01.2014. KMC also manages the discharge of city's sewage water from its pumping stations, solid waste disposal and land ownership of garbage farms at Dhapa.

Agriculture Department works for development of agriculture and allied sector in a holistic manner by ensuring farmers' access to skills, technologies, markets and financial inclusion.

#### [Different parastatals of the Department of Environment](#)

*Institute of Environmental Studies and Wetland Management (IESWM)* was established in 1986 by the Department of Fisheries. It is one of the most prominent research organizations in the Waste Recycling Region (WRR). Currently, IESWM is under the Department of Environment, Government of West Bengal.

*West Bengal Pollution Control Board (WBPCB)* established in 1974, is the statutory authority for abatement and control of pollution. *West Bengal Biodiversity Board (WBBD)* is focused on inventorization and documentation of biodiversity of the State.

Scientific and research support is also provided by *Zoological Survey of India, Botanical Survey of India and Central Inland Fisheries Research Institute (CIFRI)*. A number of Universities in West Bengal are carrying out research projects and other activities related to sewage fisheries.

#### **4.1.2 Fishery Cooperatives**

Cooperatives (government as well as private) form a small proportion of EKW fishery, and their proportion has declined from 28% to 6% during 2002-2015 (Mukherjee 2020). Trends in increasing privatization of bhery has also been recorded in other studies (Chakraborty and Gupta 2019). Much of their interest is in production activities, and their engagement in EKW management issues is limited to that of sewage distribution and desiltation of inflowing channels.

### 4.1.3 Rules and Regulations

The East Kolkata Wetlands (Conservation and Management) Act, 2006 defines the boundary of the wetland and also sets the regulatory regime for EKW. The Act stipulates maintenance of preservation of wetland area, maintenance of its ecological character and land use as settled on the date of promulgation of rules. A process for change in land use has also been laid down – although the broader mandate remains in favour of preserving current/existing land use. The EKWMA has been regularly filing cases upon contravention of these rules, and as on date, over 350 FIRs have been filed by the Authority in eight police stations having jurisdiction over the wetland area.

The West Bengal Inland Fisheries Act, 1984 (amended in 1993) accords the right to the State Government for regulating fisheries for the purpose of conservation and propagation of fish by restricting a range of activities including engines, hydrological regulation, net size, and fishing methods. Section 17A specifies that no person shall put any water area including embankment, which is capable of being used as fisheries or any naturally or artificially depressed land holding, which retains water for more than six months in a year to such use, other than fisheries.... It further states that no person shall fill up any water area including embankment or naturally or artificially depressed land holding as aforesaid, with a view to converting into solid land for the purpose of any building thereon or for any other purpose, or divide any water area so as to make any such part for any purpose other than pisciculture. Section 8 of the Act provides for state government taking over management and control of any pisciculture tank, if the norms as laid under the Act are not being complied with, or taking over the fishery is in public interest.

The West Bengal Land Reform Act (1955) and its amendments regulate land use and assign the land rights holder the responsibility to ensure preservation of the character of land in line with the purpose for which the rights were given in the first place (Section 4B). A process for change in land use has been specified in the said Act. According to the notification vide no. 4297-LR/1A-05/GE(M) dated 17.09.2009, the District Land and Land Reforms Officer shall not make any order directing change of character of any land having water body unless he has made a prior consultation in writing with the Fisheries Department and the Environment Department. A 2016 notification by Environment Department (425/EN/O-42/2014) necessitates creation of compensatory waterbody in case where waterbodies are reclaimed for developmental purposes, and authorizes West Bengal Pollution Control Board, through its regional offices in different locations to receive, examine and authorize proposal received from respective district land and land reforms officers for conversion or alteration of water bodies, except EKW.

In 2017, the Ministry of Environment, Forest and Climate Change (MoEFCC) notified the Wetlands (Conservation and Management) Rules, 2017 under the Environment (Protection) Act, 1986. The Act is applicable to all Ramsar sites and wetlands notified under these rules. Solid waste dumping and discharge of untreated waste and effluents from industries, cities, towns and villages have been listed under prohibited activities in the notified wetlands. Similarly, construction of any permanent nature has been prohibited. Meeting these conditions in EKW is difficult as the sewage from Kolkata is the major inflow source, and the wetland is inhabited by nearly 0.15 million people.

Besides the Wetlands Rules, The Indian Forest Act, 1927 and the Indian Wildlife (Protection) Act, 1972, The Indian Fisheries Act, 1897, The Water (Prevention and Control of Pollution) Act, 1974, The Environment (Protection) Act, 1986 and The Biological Diversity Act, 2002



provide substantive legal and regulatory framework for conservation of Indian wetlands including EKW.

## 4.2 Gaps

The constitution of EKWMA has been hailed as a landmark in wetlands conservation in India, as it was the first wetlands authority to be backed by legislation. With increasing developmental pressures, the need for integrated management has become pertinent. The multiplicity of organizations and laws effecting different parts of the wetlands and different ecosystem components (water, land, fish, biota) makes compliance a challenge.

Strength, weakness, opportunities and threats (SWOT) analysis of EKWMA are summarized in Table 7.

**Table 7. SWOT Analysis of current institutional arrangements for EKW management**

Strength	Weakness
<ul style="list-style-type: none"> <li>• <b>EKW presents a unique wetland wise use model – balancing human use with goal of maintenance of ecological character.</b></li> <li>• <b>Ramsar designation commits the state government and the central government to conserving wetland character and putting in place effective management arrangements for the same.</b></li> <li>• <b>EKWMA provides an institutional architecture for resolving inter-departmental conflicts.</b></li> <li>• <b>Wetlands (Conservation and Management) Rules, 2017 and East Kolkata Wetlands (Conservation and Management) Act, 2006 provide a robust regulatory framework for addressing developmental threats on the wetland.</b></li> </ul>	<ul style="list-style-type: none"> <li>• EKWMA has modest human and financial resources to deliver wetland management related functions.</li> <li>• Much of available management resources are used for enforcement of extant rules and regulation.</li> <li>• Current structure of EKWMA does not allow for robust stakeholder engagement.</li> <li>• EKWMA does not have a comprehensive wetlands inventory, assessment and monitoring system to assess changes in wetland character and determine management response options.</li> <li>• Penal measures/procedures are not explicitly defined in the Wetlands (Conservation and Management) Rules, 2017 which is currently used for regulating different developmental activities in the wetland. However, it is clearly mentioned in the East Kolkata Wetlands (Conservation &amp; Management) Act, 2006.</li> <li>• EKWMA operates through district administrative machineries, which are burdened with a large number of other responsibilities.</li> <li>• Multiplicity of agencies and government departments managing different aspects of wetland requires much coordination.</li> <li>• Integration of wetlands conservation with plans for climate change and disaster risk reduction is limited.</li> <li>• Mechanisms for aligning developmental planning within the inflowing catchments and downstream areas needs strengthening.</li> </ul>
Opportunities	Threats
<ul style="list-style-type: none"> <li>• <b>The information on wetland functioning and biodiversity values has improved thus creating a robust basis for adaptive management.</b></li> <li>• <b>Management zoning may help resolve conflicts over land use and also help balance multiple interests.</b></li> <li>• <b>A vibrant civil society network exists which can play a bridging</b></li> </ul>	<ul style="list-style-type: none"> <li>• Commercial interests due to high market value of the land in Kolkata Metro pose a threat to wetlands conservation.</li> <li>• Legal framework for EKW is litigated. The State's East Kolkata Wetlands (Conservation and Management) Act, 2006 and the Wetlands (Conservation and Management) Rules, 2017 are the subject matter of writ petitions in the Hon'ble High Court.</li> </ul>

<p><b>role between government and EKW communities to support integrated management of the wetland.</b></p>	
--	--

### 4.3 Proposed Arrangements

Reorganization of EKWMA: EKWMA requires a more efficient and results-oriented institutional structure, and gradual shift from an enforcement role to a strategic role in wetland management. A reorganised set up may include following units:

- a) Wetland monitoring unit: Responsible for systematic wetland inventory, assessment and monitoring.
- b) Engineering unit: Responsible for carrying out minor engineering and construction works.
- c) Community engagement unit: Responsible for liaison with community and civil society organisations, and ensuring integration of community concerns within management actions.
- d) Enforcement unit: Responsible for assessing adherence to the various regulatory provisions, and liaison with concerned state government department for ensuring action in case of violations.
- e) Communication and outreach unit: Responsible for external communications, by following a communications plan.

Sustainable financing mechanism: A gradual shift to financing through appropriate allocations within the state budgets would need to be made so that core wetland monitoring and management functions are not interrupted if the financing of projects is not available. EKWMA should also be capable of rendering consultancies and support to other states on wetland management issues, which could be an additional source of revenue.

Policy coherence: Development in and around EKW should be based on a single and coherent policy, which should guide various agencies in the design of various programmes. EKWMA should be consulted before implementation of developmental programmes, to ensure that no adverse impacts on the wetlands result from such interventions.

East Kolkata Wetlands Management Authority works in close coordination with the other organisations as mentioned in 4.1.1 and 4.1.2. After completion of the period of this IMP, various Departments such as Department of Fisheries, Agriculture Department, Department of Forest, Department of Irrigation & Waterways, Department of Water Resources Investigation and Development (WRIDD), Department of Panchayats & Rural Development, Urban Development and Municipal Affairs Department, Department of Land & Land Reforms would play a major role in maintaining the eco-system of the East Kolkata Wetlands in the years to come. For management of the EKW further planning and developing of IMP would be required in the subsequent years.

## 5 Management Framework

---

Management of EKW needs to be based on recognition of the full range of ecosystem services and biodiversity values of the wetland and their mainstreaming into developmental planning at all levels. The effectiveness of management will be reflected in the ability to sustain multiple use of the wetland, based on the traditional knowledge of resource recovery developed over time, without undermining the key ecological and social processes that underpin the functioning of EKW social-ecological system. Wise use of EKW will be realized when the capability of the wetland to provide its ecosystem services and sustain rich biota is maintained now as well as in future, on pathways that are aligned with ecosystem principles and guided by sustainable development. Sustainable development framework, as adopted in 2015, call for addressing five elements – People, Partnership, Planet, Peace and Prosperity.

The evaluation of wetland features, as summarized in Chapter 2 and 3 of the management plan and the institutional arrangements in Chapter 4 indicate that the current management arrangements are focused on regulation and select provisioning ecosystem services of the wetland, at the cost of regulating and supporting services. Most of the wetland features have been oriented to support production systems within the wetland. The sustainability of such a management approach is limited, as it is centres on a few wetland processes (primarily nutrient cycling) which are known to operate within ecological thresholds and if not managed within limits can cause the wetland system to change its state and functions.

As the impacts of climate change unfold over the Kolkata City and the Gangetic Delta, extreme precipitations, increase in temperatures and tropical cyclogenesis are likely to become more frequent. The ability of EKW to buffer extreme events, regulate local hydrology, as well as prevent release of harmful GHGs indicate the relevance of these wetlands in mitigation as well as adapting to climate change. At the same time, climate change and the ongoing developmental pressures may lead to intensification of existing risks as well as creating new risks of the ecosystem, thereby calling for systematic monitoring, and periodic adaptation in management approaches on the basis of new information that is generated in the process.

The current chapter sets out the management planning framework, including setting the management goal and purpose, management strategy, objectives, targets and indicators, and likely risks and risk mitigation options pertaining to implementation of the management plan.

### 5.1 Management Goal and Purpose

The goal of management planning is to *'maintain East Kolkata Wetlands in a healthy condition to enable delivery of its full range of ecosystem services and sustain biological diversity values'*.

The purpose of management is to *'enable natural infrastructure services to the Kolkata city in the form of wastewater treatment, flood buffering and climate regulation including reduction in the urban heat island effect; providing livelihood opportunities for wetlands communities in food production; and sustain diversity of biota in the landscape'*.


## 5.2 Management Strategy

### 5.2.1 Building with Nature

Building with Nature (BwN) is a comprehensive engineering approach that seeks to enhance the use of natural ecological processes to achieve efficient and sustainable infrastructural designs to provide a set of desired benefits. The BwN concept for EKW is motivated and supported by nature that may also offer environmental, economic, and social benefits, while increasing resiliency.

EKW offers wide range of natural infrastructure that have multiple functionalities like that of a water treatment plant, flood buffer, carbon sink, sediment trap, fish farms, agricultural farms, horticulture farms, temperature regulators, rich biodiversity etc.

BwN will be enabled in EKW management through the following process



**Figure 27. Embedding the concept of BwN in EKW management plan**

Implementing BwN yields a wide variety of benefits for various ecosystem services, some of which include:

**Table 8. Benefits offered by natural infrastructure within the BwN concept**

Benefit	Natural infrastructure that provide benefit
Carbon sequestration	With no treatment plant present for Kolkata city EKW helps reduce the carbon footprint ~3500 tonnes of CO <sub>2</sub> per year which is nearly 60% of Kolkata city's emission.
Improved water quality	Devised by local fishermen and farmers, these wetlands serve as the natural sewage treatment plant for the city treating over 900 MLD of raw sewage. The algae and fish consume the organic nutrients and lower its concentration by significant amounts.
Enhanced water availability	Spanning over 12500 ha EKW acts like a sponge soaking up excess water thereby recharging groundwater and storing water in the surface for agro-economic activities.
Retained soil moisture	The plant belt along the bheri help maintain soil moisture by natural adsorption process.
Reduced sedimentation	The depressions in the wetlands trap sediments and enhance the soil quality.
Controlled soil erosion	The plant belt along the bheri controls soil erosion by holding on to the soil by their roots.
Regulated air temperature	The relative humidity over EKW is higher than the surrounding area. The air transports these water droplets to the city of Kolkata thereby cooling the air. This also reduces the urban heat island effect to a considerable extent.
Controlled floods	Placed in the flood plains of Kulti and Hooghly rivers, the vast spanning wetlands trap excess water during floods which are then used for various purposes.
Bioremediation	Organic and Inorganic nutrients are significantly reduced by the water hyacinth wall around the boundaries of the bheri.
Habitat for rich Biodiversity	The EKW provides suitable habitat areas for a variety of resident and migratory species.

### 5.2.2 Management zoning for multiple use

The ability of EKW to provide the diverse ecosystem services while also harbouring diverse biota is linked with multiple use the wetland is put to. The balance of land use between aquatic usages (fish farms) and non-aquatic usages (horticulture, agriculture and settlements) is key to ecosystem functioning. The land use control principle of the East Kolkata Wetlands (Conservation and Management) Act, 2006 and Wetlands (Conservation and Management) Rules, 2017 provide the regulatory architecture under which management zoning of EKW can be framed for meeting conservation, development and regulation related outcomes. Since productive activities in the wetlands are mostly inter-twined with the banks of the waterbodies supporting horticulture and paddy cultivation using the nutrient-rich water released by the bheris, zoning activities would primarily focus on settlement areas and non-settlement areas.

### 5.2.3 Adaptive management


EKW, as several other wetland ecosystems, have an inherent uncertainty and unpredictability in their behaviour owing to complex and multi-scalar ecological, social and institutional interactions that shape their features and governing factors. There are a number of reasons, including:

- Environmental variation that is uncontrollable (such as increasing intensity of precipitation)
- Partial observability (as not all wetland features and factors can be monitored)
- Partial controllability of actions (as management interventions are implemented through a number of agencies)
- Structural uncertainty arising out of lack of complete understanding of how the ecosystem functions

Given that the knowledge on ecosystem is always likely to be incomplete, an adaptive management based on iterative learning, and using that learning to improve management using a goal-oriented and structured process (Figure 28) shall be applied. Adaptive management will be enabled in management of EKW by a combination of processes, such as:

- Structured decision making to clarify management goals, objectives and actions, involving stakeholders
- Investing into monitoring and learning for management. Each management intervention in reality is an experiment based on a working hypothesis of ecosystem functioning. Monitoring enables assessment whether the hypothesis works in reality (for example, whether changing sewage composition is leading to epidemiological concerns for the wetlands communities)
- Investing into cross-scale communication. Understanding change at multiple scales (such as City Environments, Gangetic Delta) may help getting a better understanding of ecosystem functioning and variability.
- Adaptive governance, based on collaborative and participatory management which has the flexibility of sharing management responsibilities. Successful adaptive governance requires leadership with a vision, systematic monitoring, complementary legislation framework which allows for adaptive management, information flow amongst stakeholders, and clear opportunities for stakeholders to collaborate.





**Figure 28. The Adaptive Management Cycle (after Allen et al, 2011)**

#### 5.2.4 Multiple values of nature and nature's contribution to people


Management plan implementation will take into account the diverse ways in which nature and nature's contribution to people support well-being. These will include intrinsic values (the value of EKW as an ecosystem with its complex ecological functions), instrumental values (the value of EKW towards meeting food security, water quality regulation and climate moderation), and relational values (the values linked with a sense of place and cultural identity which communities attribute to EKW). The management plan will be built on the full range of values to bring in multiple perspectives in decision-making and implementation of programmes.

#### 5.2.5 Focus on behaviour change

The management plan entails a shift from a high emphasis on wetland regulation to inducing positive behaviour within wetlands communities and stakeholders which are aligned with wise use. The EKWMA will use strategic communication to inform stakeholders on the role of EKW in their overall well-being, and the ways these groups can engage in ensuring that the wetland continues to deliver their wide-ranging services. Information, education and communication tools, tailor-made to the needs of various stakeholder groups, would be proactively used to trigger behaviour change, along with building capacities and opportunities for participation in wetlands management.

## 5.3 Management Objectives and Performance indicators

The management strategies have been translated into eight objectives which reflect the desired state of key features of wetlands. For each feature, the performance indicators are the attributes which can indicate change. The desired outcome reflects the target which the management plan envisages to achieve within five years, thus providing a monitoring framework to assess effectiveness. These indicators and targets have also been integrated into the monitoring plan discussed in Chapter 6 of the management plan.



**Figure 29. Wise use of wetlands for the maintenance of ecological character at East Kolkata Wetlands**

**Table 9. Management objectives, performance indicators and targets**

Objectives	Performance Indicators	Desired outcome
Objective I. Land use and land cover of the wetland is maintained in line with the regulatory requirements under Wetlands (Conservation and Management) Rules, 2017 and East Kolkata Wetlands (Conservation and Management) Act, 2006.	Land use and land cover change in comparison with the 2000 baseline  Number of violations of the Wetlands (Conservation and Management) Rules, 2017	No illegal transformation of land use  Violations are reduced by atleast 60%

	and East Kolkata Wetlands (Conservation and Management) Act, 2006	
<b>Objective 2. Sewage quantity and quality received within the wetland is efficiently treated applying traditional waste recovery practices.</b>	<p>Sewage availability through natural hydrological gradient</p> <p>Quality of sewage</p>	<p>Tertiary channels deliver sewage without use of pumps</p> <p>Equitable distribution of sewage.</p> <p>Heavy metals concentration in sewage is brought to safe levels</p>
<b>Objective 3. Maintaining the biodiversity within East Kolkata Wetlands.</b>	<p>Species richness</p> <p>Population of migratory waterbirds</p> <p>Population and habitat of marsh mongoose</p>	<p>No species extirpation</p> <ul style="list-style-type: none"> <li>Improving the richness of species</li> <li>Recover of species, if any</li> </ul> <p>Counts are maintained in the range of 20% deviation from average of last five years</p> <p>Sighting is maintained in the range of 20% deviation from average of last five years</p>
<b>Objective 4. Species invasion threats to fisheries is reduced.</b>	Occurrence of invasive species in fish catch	Reduced by at least 80% of occurrence in 2020
<b>Objective 5. Sustainable livelihood with risk reduction.</b>	<p>Resource productivity (fish catch, vegetable harvest)</p> <p>Instances of waterborne diseases</p> <p>Diversification of income sources</p>	<p>Non-declining harvest of fish and vegetables</p> <p>Reduced by at least 80% of occurrence in 2020</p> <p>Wetland communities having income in the lower quintiles gain additional sources of income</p>
<b>Objective 6. Individual and</b>	Participation of wetland	Integration of community,

<p><b>collective capacity and opportunities for stakeholders and wetland communities to participate in wetland management and contribute to wetland wise use is enhanced.</b></p>	<p>communities in management plan implementation</p> <p>Effective wetlands inventory</p> <p>Evidences of affirmative behaviour change within communities living in and around EKW supporting wise use of wetlands</p> <p>Evidences of community-led collaborative action for wise use of wetlands</p>	<p>rights and capacities in management plan implementation and monitoring</p> <p>An Integrated Wetlands Inventory, Assessment and Monitoring System is put in place and used.</p> <p>Local action for addressing solid waste or preventing encroachment</p> <p>Community norms for preventing land use change or overharvesting of resources</p>
<p><b>Objective 7. Systematic wetlands inventory, assessment and monitoring system is used to inform management decisions and assess effectiveness.</b></p>	<p>Availability of time-series data on wetland ecological character</p> <p>Availability of data on threats leading to adverse change in ecological character.</p> <p>Evidences of use of data generated from wetlands inventory, assessment and monitoring system in decision-making.</p>	<p>Data is available on all priority wetland features</p> <p>Data is available on threats</p> <p>Data is systematically analysed and presented in EKWMA meeting.</p>
<p><b>Objective 8. Integration of multiple values of wetlands in</b></p>	<p>Number of sectoral policies (wherein</p>	<p>EKMWA meets regularly and considers implications</p>

<b>sectoral developmental planning is enhanced.</b>	integration of wetlands is relevant) which take into account wetlands values.	of sectoral plans.  EKWMA enables integration of role of EKW in programmes and actions plans on climate change, urban development, and disaster risk reduction.
---	---	---

In the table below, a mapping of each of the objectives against the values (derived from Chapter 2), threats (derived from Chapter 3) and indirect drivers (derived from Chapter 4) is presented.

**Table 10. Mapping of management objectives against wetland values and threats**

	Management Plan Objectives							
	Obj. 1	Obj. 2	Obj. 3	Obj. 4	Obj. 5	Obj. 6	Obj. 7	Obj. 8
Ecosystem Services								
<b>Wastewater treatment</b>	++	+++			+	+		
<b>Carbon sequestration</b>	++	+++				+		
<b>Flood buffering</b>	++				+	+		
<b>Wetland products</b>	++	+++	+	+	++	+		
<b>Recreation</b>			++	++	+++	++		
<b>Relational values</b>					+++	+++		
Biodiversity								
<b>Diverse biota</b>			+++	+++				
Threats (Direct)								
<b>Encroachments</b>	+++					+	++	
<b>Heavy metal contamination in sewage</b>		+++						
<b>Increased solid waste dumping</b>	+++							
<b>Land subsidence</b>							++	
<b>Sedimentation of channels</b>		+++						
<b>Invasive species</b>			+++	+++				

<b>Climate change local impacts</b>		++					++	
Threats (Indirect)								
<b>Lack of clarity on regulation</b>						+	++	
<b>Limited stakeholder engagement in governance</b>					+++	+++	+++	
<b>Limited capacity within EKWMA for integrated management</b>						+++	++	+++
<b>Limited consideration of full range of wetland values in sectoral planning and decision making</b>			+	+		++	+++	++

## 5.4 Risks and Risk Mitigation Options

The management plan design is based on certain assumption. The table below identifies the risks of these assumptions impacting management plan implementation adversely, and possible risk management measures.

**Table 11. Risks and risk mitigation measures**

Risks	Risk management measures
At goal level	
<b>Local communities participate in the conservation and management of the EKW</b>	A Community Advisory Group will be constituted to advice regarding management plan implementation and disseminate information within communities
At objective level	
<b>Trained human resources are available for management of EKW</b>	EKWMA staff will be imparted training on various aspects of wetlands management, and hand-holding support provided.
<b>Communities are sufficiently organized to engage in and take ownership of the project activities</b>	The Community Advisory Group will be used as a mechanism to support and strengthen community networks.
<b>A long-term financial framework is established for maintaining the expanded EKWMA programme</b>	Funding from various convergence sources will be leveraged, including from private sector organizations.

At Output level	
<b>Qualified trainers and experts are available</b>	A roster of experts and trainers will be prepared for supporting capacity development.
<b>Strategies offer flexibility for adaptation</b>	EKWMA will present progress of management plan implementation in the meetings. Data from monitoring will also be analysed to assess effectiveness of interventions.
<b>Robust monitoring mechanisms and application of scientific approach for resolving conflicting issues</b>	A Wetlands Inventory, Assessment and Monitoring system will be put in place in the first year of the management plan implementation
<b>Capable NGOs are available to assist community-based activities</b>	EKWMA will put in place a process of selecting and inducting NGOs in management plan implementation

## 6 Monitoring Plan

---

Management of EKW is aimed at maintaining its ecological character, and in doing so, retaining those essential ecological and hydrological functions which ultimately enable the wetland to provide its provisioning, regulating and cultural services. Having a system to describe, monitor and detect changes in ecological character is therefore critical to support decision making for wise use of this Ramsar Site. Equally important is to be able to assess effectiveness of management in terms of ability to develop and implement an integrated planning, management and evaluation system to secure wise use of the wetland.

The current section of the management plan describes a monitoring framework for EKW to support integrated management for wetland wise use. The section details monitoring purpose and strategy and associated resource requirements. The monitoring plan is proposed to be applied both at the scale of wetland ecosystem, as well as institutional arrangements supporting management. Thus, a section outlining strategy and framework for assessing management effectiveness is also included. The cost implications of the monitoring plan have been factored in the Chapter 7 (management planning framework) and Chapter 8 (budget and financing).

### 6.1 Monitoring Objective

Developing a monitoring plan for EKW requires addressing the following inter-related requirements of wetland inventory and wetland assessment. It is imperative therefore to put in place an integrated Wetland Inventory, Assessment and Monitoring System (WIAMS) to address the overall information needs for wetland management, and to provide a robust decision support system for the same. The ambit of monitoring is also envisaged to include assessment of management effectiveness. Following are the specific objectives for establishing WIAMS:

- Developing up-to-date and scientifically valid information on status and trends of wetland features and influencing factors
- Establishing a baseline for measuring change in ecosystem components, processes and services
- Informing decision makers and stakeholders on the status and trends in biodiversity, ecological functioning and ecosystem services of the wetland
- Supporting compliance to national and state legal requirements and regulatory regimes
- Determining impacts of developmental projects on ecosystem components, processes and services
- Identifying risks to ecological character and support development of response strategies
- Assessing effectiveness of wetland management



## 6.2 Monitoring Strategy

Monitoring strategy responds to the following information needs EKW:

- inventory - to establish the ecological character baseline
- assessment – to establish status, trends and threats to wetland using inventory information
- monitoring – to assess changes in status and trends, including reduction in existing threats or appearance of new threats, or even changes in management effectiveness

As this information pertain to various spatial scales, the overall information requirements can be classified into three hierarchical levels:

- the aquatic environment of EKW
- East Kolkata Wetlands
- EKW Basin

A hierarchical classification of inventory, assessment and monitoring needs for EKW is presented in Table I2. The information needs for inventory are derived from the core datasets needed to establish a baseline on ecological character<sup>5</sup> for EKW, and contain all the essential ecosystem components, processes and services, as well as management related parameters that characterize the site. Within the aquatic environment, information needs pertain to sewage inflow, quality, its uptake for resource recovery, and ecosystem services such as wastewater treatment, flood buffering and carbon sequestration. At the level of wetland, information needs pertain to land-use and land cover change, and threats such as solid waste dumping. At the basin scale, the information requirement is related to geo-morphological and climatological setup, as well as basin wide management arrangements, particularly those related to land and water resources and urban planning. At all levels, information on institutional arrangements and management practices is included so as to enable creation of a baseline on sectoral programmes, and the linked stakeholders, which are likely / have an impact on the wetland state. While not explicitly mentioned, strategic environmental assessments can be commissioned for any developmental project that has / likely to have negative impact on the wetlands.

Information needs for monitoring EKW have been derived from assessment of ecological character carried out for development of the management plan. Four cluster of needs have been identified: a) land use and land cover change, to assess the dynamics of land use within the catchment; b) hydrological regimes, to assess the flux of water, sediments and nutrients; c) ecological components and processes, to assess the biodiversity, habitat quality and resource productivity; and d) socioeconomics and livelihoods to assess the trends in ecosystem services – livelihoods interlinkages.

This monitoring information adequately addresses the needs of East Kolkata Wetlands (Conservation and Management) Act, 2006 and Wetland (Conservation and Management)

---


<sup>5</sup>Derived from the core inventory fields required for ecological character description as per Ramsar Convention Resolution X.15: Describing the ecological character of wetlands, and data needs and formats for core inventory: harmonized scientific and technical guidance. These fields have been further integrated into guidance related to information requirement for describing Ramsar site at the time of designation and subsequent updates (Ramsar Convention Resolution XI.8 and XI.8 annex 1)

Rules, 2017 of the Ministry of Environment, Forests and Climate Change. A list of wetland features, indicators and corresponding methodology and data collection frequency is provided as Table 13. Proposed stations for hydrological, air quality and biodiversity monitoring are indicated in Map 16.

The monitoring and assessment needs are envisaged to be addressed by a dedicated monitoring programme and specific research and assessment projects. Inventory, being based on collated information on identified wetland features and management practices, will be developed based on the monitoring and assessment information, as well as secondary sources.

Inventory, assessment and monitoring form an integral part of wetland management, and thereby core activity of EKWMA. The management plan proposes to establish a dedicated wetland monitoring unit with adequate infrastructure support to effectively deliver this function.

Linkages also need to be developed so that data from the existing monitoring networks of different agencies (for example, inundation and flooding information from Central Water Commission and Department of Water Resources; groundwater quality and quantity from Central Ground Water Board; select surface water quality parameters from West Bengal Pollution Control Board; and fish production from Fisheries Department) can be accessed and shared. Similarly, provision for participation of NGOs and civil society in monitoring programme has also been built, especially for socioeconomics and livelihoods aspects and biodiversity monitoring (for example, waterbird census being implemented by NGOs under the aegis of Asian Waterbird Census). Thematic management needs-based research can be taken up by specialized agencies such as ZSI, BSI, CIFRI, Indian Statistical Institute, Kolkata to complement the monitoring programme.



**Map 16. Proposed monitoring stations for water flow, water quality, air quality and biodiversity**

**Table 12. Inventory, assessment and monitoring needs for managing East Kolkata Wetlands**

	<b>Information Purpose</b>		
<b>Information Scale</b>	<b>Inventory</b>	<b>Assessment</b>	<b>Monitoring</b>
<b>East Kolkata Wetlands (aquatic region)</b>	<p><b>Physical extent</b> (area, depth)</p> <p><b>Sewage quantity and quality</b></p> <p><b>Status of canals</b> (maintenance of natural gradient)</p> <p><b>Harvest</b> (fish and aquatic species)</p> <p><b>Biota</b> (plant and animal species)</p> <p><b>Socioeconomics</b> (return on fisheries, cooperatives operation, status of fishfarm workers and owners)</p>	<p><b>Water quality improvement</b> (extent of reduction in BOD, COD, nutrients and heavy metals)</p> <p><b>Carbon sequestration</b> (carbon stored in various forms in resource recovery process)</p> <p><b>Fish production trends</b></p> <p><b>Trends in invasive species</b></p> <p><b>Trends in number and population of species</b></p>	<p><b>Land use and Land cover change</b> in fish farm areas</p> <p><b>Sewage quality</b> (presence of heavy metals and toxic chemicals)</p> <p><b>Spread of fish invasives</b></p> <p><b>Bioaccumulation</b> in fish</p> <p><b>Disease</b> (related to sewage fed fisheries operations) incidence in fish farm workers and owners</p>
<b>East Kolkata Wetlands</b>	<p><b>Physical setting</b> (area, boundary, connectivity)</p> <p><b>Water regime</b> (sewage flows, inflow -outflow balance, surface-groundwater interactions, inundation regimes, quality, regulation)</p>	<p><b>Ecological character change</b> (change in ecosystem components, processes and services – can also be derived based on assessment of indicators related to ecosystems, habitat, species and / or management)</p> <p><b>Water quality trends</b> (extent of reduction in BOD, COD, nutrients and heavy metals)</p>	<p><b>Land use and Land cover change</b> (with respect to 2000 baseline)</p> <p><b>Bioaccumulation</b> in vegetables and crops</p>

	<p><b>Harvest of produce from wetland region</b> (vegetables, food crops and others)</p> <p><b>Groundwater</b> (level, abstraction and quality)</p> <p><b>Biota</b> (plant and animal species)</p> <p><b>Socioeconomics</b> (return on fisheries, horticulture and agriculture; incomes and assets)</p> <p><b>Sectoral programmes and institutional arrangements</b> for management of land and water resources and biodiversity conservation</p>	<p><b>Soil quality trends</b></p> <p><b>Inundation trends</b></p> <p><b>Groundwater level and quality trends</b></p> <p><b>Air quality trends</b> (PM 2.5, PM 10, temperature, humidity)</p> <p><b>Carbon sequestration</b> (carbon stored in various forms in resource recovery process)</p> <p><b>Trends in key species</b> (waterbirds, marsh mongoose)</p> <p><b>Production trends</b> (vegetables, food crops)</p>	
<b>East Kolkata Wetland Catchments</b>	<p><b>Geology and Geomorphology</b> (Soils, elevation, slope, drainage pattern)</p> <p><b>Climate</b> (Precipitation, Temperature)</p> <p><b>Land use and land cover</b></p> <p><b>Water regimes</b> (river flows, runoff, upstream abstraction)</p>	<p><b>Climate risk and vulnerability</b> (changes in runoff and implications for EKW)</p>	<p><b>Urban planning</b> (water regulating structures and water allocation, sewage treatment infrastructure development, expansion of settlements) and implications for EKW</p>

**Table 13. Monitoring and assessment parameters and indicators**

Parameter	Indicator	Priority	Monitoring Method	Monitoring Frequency
<b>Land Use and Land Cover</b>				
Land use and land cover change within EKW	% area under various land use and cover classes (agriculture, , settlements, wetlands)	High	GIS and Remote Sensing  Radar sensed data	Once every year
Compliance with East Kolkata Wetlands (Conservation and Management) Acts, 2006 and Wetlands (Conservation and Management) Rules, 2017	Violations	High	Mobile based surveillance system operated by communities' wardens  Drone mapping and other technologies for surveillance of land use change	Daily
<b>Hydrological Regimes</b>				
Water flux	Sewage inflow	High	Monitoring stations between Bantala and Ghusighata lock gates as proposed in Map 16	Daily
	Sewage outflow	High		Daily
	Sediment inflow	High		Annual
	Sediment outflow	High		Annual
Water holding capacity	Bathymetry	High	Bathymetric surveys	Once in 5 years
Inundation Regime	Seasonal fluctuation in water spread area	High	Remote sensing	Once in 5 years

Hydrological Regimes				
Sewage quality	Temperature	Medium	Standard procedures of APHA	Monthly
	pH	High		Monthly
	Dissolved Oxygen	High		Monthly
	Specific Conductance	High		Monthly
	Nutrients and Nutrient Cycling (Nitrate, Phosphate, Silicate)	High		Monthly
	Cations and Anions (Calcium, Magnesium, Sulphate, Chloride, Fluoride, Sulphite)	High		Monthly
	Chemical Oxygen Demand	High		Monthly
	Heavy metals (Arsenic, Mercury, Cadmium, Chromium, Lead)	High		Monthly
	Biological oxygen demand	Medium	Standard procedures of APHA	Monthly
	Total Coliform	Medium		Annual
	Faecal coliform	Medium		Annual
	Wetland soils	Texture	Low	Standard procedures of APHA
pH		High	Annual	
Organic carbon		High	Annual	
Available nitrogen		High	Annual	
Available phosphorus		High	Annual	
Available calcium carbonate		Medium	Annual	
Heavy metals (Arsenic, Mercury, Cadmium, Chromium, Lead)		High	Annual	
Ground water	Water level	High	Methodology approved by Groundwater Estimation Committee (1997)	Seasonal (Summer, Monsoon, Post-monsoon, Winter)
	Conductivity	Medium		Annual
	Total hardness	Medium		Annual
	Chloride	Medium		Annual
	Fluoride	High		Annual
	Arsenic	High		Annual
	Heavy metals	High		Annual

Hydrological Regimes				
Water abstraction	Water abstracted for irrigation	Medium	Survey	Annual
	Water abstracted for domestic use	Medium		
Air quality	Suspended Particulate Matter (PM 2.5, PM 10)	High	Standard procedures laid by CPCB for air quality monitoring	Daily
	Temperature and Relative Humidity	High		
Ecosystem Processes and Biodiversity				
Flora	Phytoplankton (diversity and abundance)	Medium	Taxonomic studies, Standard procedures in Central Inland Fisheries Research Institute Bulletin No. 10	Seasonal  (Pre monsoon and Post monsoon, if required)
	Periphyton	Medium		
	Macrophytes (diversity and abundance)	High		
	Species invasion	High	Habitat Sampling and Remote sensing (using high resolution data)	Once in 2 years
	Primary production	High	Standard procedures in Central Inland Fisheries Research Institute Bulletin No. 10	Seasonal
	Bioaccumulation in vegetables and food grains	High	Environmental monitoring protocols of ICAR - CIFRI /Standard procedures of APHA	Bi-annual
Fauna	Zooplankton (diversity and abundance)	Medium	Taxonomic studies, Standard procedures in Central Inland Fisheries Research Institute Bulletin No. 10	Seasonal
	Aquatic macro-invertebrates	Medium	Taxonomic studies, Standard procedures in Central Inland Fisheries Research Institute Bulletin No. 10	Seasonal
	Aquatic Insects	Medium	Taxonomic studies, Standard procedures in Central Inland Fisheries Research Institute Bulletin No. 10	Seasonal
	Fish diversity	High	Taxonomic studies	Once in 5 years
	Amphibians, Reptiles, Mammals	High	Taxonomic studies	Once in 5 years
	Fish catch and effort (number of harvest cycles, catch)	High	Standard procedures in Central Inland Fisheries Research Institute Bulletin No. 10	Monthly



Ecosystem Processes and Biodiversity				
Fauna	Bioaccumulation in fish	High	Environmental monitoring protocols of ICAR - CIFRI / Standard procedures of APHA	One every year
	Water Bird population and diversity	High	Census and Taxonomic studies	Annual
	Avian disease	Medium	Surveillance	Annual
	Feasibility study to ascertain kind of certification possible for fish produced in EKW	Medium	Private standards and certification in fisheries and aquaculture, FAO FISHERIES AND AQUACULTURE TECHNICAL PAPER 553	Once in 5 years
Socioeconomics and livelihoods				
Community dependence on wetland ecosystem services	Fish catch and vegetable production	High	Socioeconomic survey	Monthly
	Performance of cooperatives (Capitalization, infrastructure, catch processed, membership)	High		Annual
	% contribution of fisheries and horticulture to income and employment	High		Bi-annual
	Number of tourists visiting wetland and direct and indirect spending	High		
Livelihood status of wetland dependent communities	Physical capital, financial capital, social capital, human capital indicators of livelihood systems	Medium	Socioeconomic survey	Bi-annual
	Instances of diseases related to working in Sewage -Fed Fisheries	High		
	Number of reported instances of conflicts	Medium		

### 6.3 Assessing management effectiveness

East Kolkata Wetlands are dynamic ecosystems and so are their management needs. Management plans, which are developed based on assumptions known to managers, need to be periodically assessed to make sure that the set goals and objectives are being achieved.

The effectiveness of management towards achieving the overarching objective of maintenance of ecological character can be greatly enhanced if following questions are periodically reflected upon:

- What is the current status of the EKW?
- Is the management achieving the goal of maintenance of ecological character?
- What are the current and future threats?
- Are adequate resources available for implementing management, and if not, how can they be improved?
- Are management processes adequate, effective and efficient?
- What other steps can be taken to improve management?

The Contracting Parties to the Ramsar Convention adopted the R-METT (Ramsar Site Management Effectiveness Tracking Tool) to assist Ramsar site managers in assessing effectiveness of management in achieving wetland wise use outcomes. The assessment looks into the following aspects:

- **Context** of management (wetland ecological character, threats and risks of adverse change).
- **Management planning** that defines how the management goals and objectives have been defined.
- **Inputs** including human, technical and financial resources applied to implement management actions.
- **Process** of management plan implementation.
- **Outputs** (tangible and intangible) that result from implementation of management actions.
- **Outcomes** with respect to the objectives defined by the management plan

It is proposed that management effectiveness assessments for EKW is done at least once in five years so that management action plans are revised and updated to reflect the condition of wetlands as well as ability of management to prevent adverse change in ecological character. A baseline assessment is proposed to be done at inception of the management plan.

## 6.4 Infrastructure and human resources requirements

Implementing the monitoring strategy as outlined in the previous sections requires the following physical and human infrastructure support:

- Remote Sensing and GIS unit with advanced capabilities of remote sensing image processing, preparation of maps and development and maintenance of spatial datasets
- Ecological monitoring laboratory with capabilities for analysis of chemical, physical and biological properties of water and soil
- Mobile based citizen reporting system for recording and reporting illegal encroachments
- Database system for storing and retrieving monitoring and assessment data. The monitoring data would be stored along with metadata, as per the quality control procedures suggested in the following sections.
- Network of hydro-meteorological and water quality stations for hydro-biological monitoring

Deployment of the aforementioned resources can be done in a cost-effective manner by applying the lessons and expertise of the existing infrastructure created by the state government for management of EKW. Need based training programmes will also be conducted to upgrade skills of the concerned state government departments and agencies.

## 6.5 Reporting

Reporting constitutes an important element of wetland monitoring programme. The intended user group, format, style and peer review requirement need to be set in the initial phases of set up of the monitoring programme.

Periodic reports, for example as a part of the annual report of the EKWMA should aim to provide a summary overview of the outcomes of monitoring.

Special publications, for example wetland atlases constituting thematic maps on various parameters are intended to inform stakeholders on wetland status and trends.

Outcomes of specific assessments, for example ecological character status and trends, economic valuation, environmental flows etc. could be made available in the form of technical report series, with an extended summary for general readership. As the monitoring programs get sophisticated over a period of time, real time monitoring options through use of satellite-based data communication techniques can be explored.

## 6.6 Quality control

Quality control in monitoring systems is required to ensure the scientific validity of sampling, laboratory analysis, data analysis and reporting. They also play a critical role in preventing introduction of random and systematic errors in data collection, analysis and reporting.

It is recommended that a Quality Management and Assurance Plan is developed for the monitoring programme. The plan should determine, *inter alia*:

**Specification of objectives** for sampling programme

**Data quality objectives:** maximum amount of uncertainty that can be tolerated to ensure that the data is fit for intended use

**Sampling programme design:** Statistical robustness of sampling frame; means to ensure that samples are representative of environment; sample recording; procedures for minimizing environmental impact

**Documentation:** Procedures for field sample record keeping and methods documentation

**Sample processing validity** (especially for water quality and biological components)

**Data quality control methods:** processes for quality control samples, duplicates and replicates,

**Performance audit procedures,** including data and systems audit

## 6.7 Review and adaptation

A periodic review of the monitoring programme is required to determine the extent to which the objectives, particularly support to management is achieved, and monitoring system remains relevant for the wetland state (particularly in the light of new and emerging threats). The review process should also aim at increasing the sophistication of the monitoring system to be able to assess complex landscape scale processes affecting the ecological character of wetland and related management.

Review process should include documentation on the way wetland inventory, assessment and monitoring information is being used to support management planning and policy goals. Review should also include identification of appropriate mechanisms to ensure that wetland monitoring is continued in the event of a funding shortfall.

## 7 Action Plan

---

Activities to meet the eight objectives have been clustered under four components: Institutions and Governance, Water management and Pollution Abatement, Species and Habitats and Sustainable Resource Development.

**Table 14. Management plan components**

Component	Objectives
Component 1 – Institutions and Governance	<p>Objective 1. Land use and land cover of the wetland is maintained in line with regulatory requirements under Wetlands (Conservation and Management) Rules, 2017 and East Kolkata Wetlands (Conservation and Management) Act, 2006</p> <p>Objective 6. Individual and collective capacity and opportunities for stakeholders and wetland communities to participate in wetland management and contribute to wetland wise use is enhanced.</p> <p>Objective 7. Systematic wetlands inventory, assessment and monitoring system is used to inform management decisions and assess effectiveness.</p> <p>Objective 8. Integration of multiple values of wetlands in sectoral developmental planning is enhanced.</p>
Component 2. Water management and pollution abatement	<p>Objective 2. Sewage quantity and quality received within the wetland is efficiently treated applying traditional waste recovery practices.</p>
Component 3. Species and habitats	<p>Objective 3. Maintaining the biodiversity within East Kolkata Wetlands.</p> <p>Objective 4. Species invasion threats to fisheries is reduced.</p>
Component 4. Sustainable Resource Development and Livelihoods	<p>Objective 5. Sustainable livelihood with risk reduction.</p>




Figure 30. Management interventions proposed at East Kolkata Wetlands

## Component I. Institutions and Governance

### I.1 EKWMA Reorganization and Strengthening

EKWMA requires an efficient and results-oriented institutional structure, and gradual shift from an enforcement to a strategic role in wetland management. A reorganised set up is proposed to include following units:

- Planning and Design unit: Responsible for strategic planning for conservation and wise use, including assessing adherence to the various policy mandates and regulatory provisions.
- Enforcement unit: Responsible for ensuring compliance with extant regulation, and coordinating with line departments on implementation of penal provisions. This unit will closely liaise with police, district administrations of South 24 Parganas and North 24 Parganas, KMC and BMC.
- Wetland monitoring unit: Responsible for systematic wetland inventory, assessment and monitoring. The unit can function out of Institute of Environmental Studies and Wetland Management which has existing infrastructure to support ecological and hydrological monitoring of the wetland.
- Community engagement and communications unit: Responsible for liaison with community and civil society organisations, and ensuring integration of community concerns within management actions. This unit will be also responsible for external communications, based on a communications plan.

The following activities are also proposed to be taken with due approval of the Governing Body of EKWMA:

- Sustainable financing mechanism: A gradual shift to financing through appropriate allocations within the state budgets would need to be made so that core wetland monitoring and management functions are not interrupted if the financing of projects is not available. EKWMA may also render research and consultancy support to other states on wetland management issues, which could be an additional source of revenue.
- Policy coherence: Development programmes around EKW need to be based on a single and coherent policy, which should guide various agencies in the design of various programmes. A mechanism for ensuring that EKWMA is consulted before implementation of developmental programmes is proposed to be developed.

### I.2 Wetland demarcation

It is proposed to demarcate wetland boundary by placing geo-tagged pillars all along the wetland periphery. Demarcation will be done using geo-tagged boundary pillars (approximately 1600 pillars are required to be placed at 500 m interval).

Activities to be carried out are as follows:

- a) Completion of field reconnaissance survey
- b) Placement of geo-tagged pillars

### 1.3 Management zoning


For the ease of management, and also ensuring compliance with the extant regulatory regimes, it is proposed to segment the wetland into four major zones (Map 17), while recognizing that the interdependence between these zones and the overarching goal of securing wise use of the wetlands. It must be accepted that the East Kolkata Wetlands have its unique character where the entire eco-system is dependent on the sewage water reaching the area and thus the wetland area is interspersed in the other zones as well. Naturally, our zoning plan will take this unique aspect into consideration. Following are the broad zones:

- Fishery zone: Wetland area that is subjected to use for fish cultivation using sewage water and subject to multiple land ownership. The zone can be used additionally by providing regulating services for absorbing storm water flows during monsoon, act as a flood storage and help recharge ground water.
- Agriculture zone: A part of the wetland permanently being used for cultivation of rice and other crops. The zone can additionally be managed by using open spaces for storing excess flood water and allow infiltration of stormwater. The zone has the potential to attract wintering waterbirds if organically farmed. Rainwater harvesting in agricultural fields to enhance availability of freshwater.
- Horticulture zone: Part of the solid waste dumping area on the western periphery of the wetland which has been converted as a waste recycling region for cultivating vegetables. The zone can be maintained as an area for practicing horticulture. Dhapa dumpsite of KMC has been existing at this zone of EKW much prior to the declaration of Ramsar site. KMC is conscious of its responsibility in maintaining a healthy environment. However, resource recovery system such as composting, biomethanation, material recovery facility (MRF) for recycling, sanitary land filling is recommended in the zone.
- Nature protection zone: A part of the wetland where biodiversity value is the highest, and habitat use by migratory species. The zone can act as a buffer for species when moving between rural and urban areas. The zone also has the potential to be used as a site for recreation and ecotourism, within the framework of existing provisions of Wetlands (Conservation and Management) Rules, 2017, and other extant regulation.

The zoning plan will guide management interventions. Following steps are proposed:

- Finalization of zoning plan in consultation with stakeholders
- Preparation of zonal management plans
- Implementation of zonal management plans within the framework of action plan as described in this document.





**Map 17. Proposed management zones of East Kolkata Wetlands**

## 1.4 Wetlands Inventory, Assessment and Monitoring System

An integrated wetland inventory, assessment and monitoring system is proposed to be set up to address the overall information needs of wetland management and to provide robust decision support system for the same. Specific objectives and a detailed framework have been outlined in Chapter 5 of the management plan. The following activities are proposed:

### *1.4.1 Establishment of wetland monitoring and research centre*

To support monitoring and research activities related to EKW, a robust and upgraded laboratory is required to be established. However, the laboratory may be housed within the Institute of Environment Studies and Wetland Management, an Institute under the Environment Department which works in close coordination with the EKWMA.

### *1.4.2 Development of database management system*

A database system for storing, retrieving and analysing the WIAMS is proposed to be set up in a GIS environment. This will include:

- development of data quality management and assurance plan including specification of data collection objectives, data quality objectives, sampling programme design, data and metadata documentation procedure, data quality control methods and performance audit procedures;
- development of GIS based database management system

### *1.4.3 Wetland monitoring and evaluation*

Wetland monitoring and inventory protocols for land use and land cover, hydrological regimes, ecosystem processes and biodiversity and socio-economics and livelihoods as proposed in Section 5.2 will be implemented.

### *1.4.4 Surveillance system*

A mobile-based surveillance system for recording infringements to Wetlands (Conservation and Management) Rules, 2017 and East Kolkata Wetlands (Conservation and Management) Act, 2006 is proposed to be developed. The app-based system will enable „community wardens“ to transmit information on violations in EKW to EKWMA through geotagged pictures and related data. Drone mapping and other sophisticated technologies such as installation of CCTV will also be used for surveillance of land use change.

### *1.4.5 Ecosystem Health Report Card*

It is proposed to develop an Ecosystem Health Report Card, and publish annually to assess and communicate wetland monitoring information to decision-makers and stakeholders. The health report card summarizes indicators along major indices (water quality, catchment status, biodiversity status) which represent various ecosystem features of the wetland, and are reported against respective thresholds set in line with management goals.

Following activities are to be taken:

- Convening a methodology workshop for EKW wetlands monitoring unit for firming up variables to be used for assessment and data requirements
- Development of Ecosystem Health Report Card
- Report card publication
- Stakeholder dissemination workshop

## 1.5 Research

Following specific research studies are proposed to be commissioned to address the knowledge gaps in assessing status and trends in ecological character:

- Carbon and GHG Flux assessment – to assess the role of EKW in sequestering carbon and GHG, and this integrate role of these wetlands in climate change mitigation strategies
- Nutrient budget - to assess the quantity of nutrients entering EKW, its uptake in resource recovery practices and discharge downstream
- Bioaccumulation studies – to assess the chain of heavy metal contamination in EKW, quantity of toxic metals accumulating in fish, vegetables and crops cultivated in EKW and possible remedial measures
- Multiple values assessment – to assess multiple values communities living in and around EKW associate with the wetland, the underlying reasons and the ways in which this value can be orientated towards behaviour change for wetland wise use

## 1.6 Capacity Development

To support integrated management of EKW, the human capacity of EKWMA and concerned line departments as well as stakeholders will be enhanced by training workshops on:

- Integrated wetlands management
- Wetlands inventory assessment and monitoring
- Land use planning for multiple wetlands use
- Wetlands and climate change mitigation and adaptation
- Integrating wetlands wise use in developmental planning
- Conducting excursions (National and International), developing course module relating to wetland/environment management

## 1.7 Communication, Education, Participation and Public Awareness

Stakeholder engagement in wetland management will be promoted through creating awareness on biodiversity and ecosystem services values, management strategies adopted and opportunities for participation. Specific activities to be undertaken include:

### 1.7.1 Signage

Signage indicating EKW as a Ramsar Site, is proposed to be placed at all major entry points of the wetlands complex, namely along Basanti Road, EM Bypass and Bantala Lock Gate. Signage are also proposed in the 7 Panchayat Offices and school premises inside wetlands.

At the 19th meeting of the Ramsar Standing Committee, 29 October-1 November 1996, the members adopted a decision that defines recommended wording for signs at all Ramsar Sites, when translated into the local languages of the sites. The decision reads as follows: The Contracting Parties should endeavor to place descriptive signs at all Ramsar Sites, and these signs should include the Ramsar logo, as well as the following suggested text (amended for EKW):

THIS SITE, COVERING 12,500 HECTARES, IS DESIGNATED BY THE  
GOVERNMENT OF INDIA AS A WETLANDS OF INTERNATIONAL IMPORTANCE  
ESTABLISHED UNDER THE CONVENTION ON WETLANDS, THE

INTERNATIONAL TREATY SIGNED IN RAMSAR (IRAN) IN 1971 TO PROMOTE THE CONSERVATION AND SUSTAINABLE USE OF WETLAND AREAS WORLDWIDE.

The protection and management of this site is under the responsibility of: EAST KOLKATA WETLANDS MANAGEMENT AUTHORITY, DEPARTMENT OF ENVIRONMENT, GOVERNMENT OF WEST BENGAL.

#### *1.7.2 Webpage*

The webpage [www.ekwma.in](http://www.ekwma.in) shall be periodically updated with information, datasets and communication products on EKW. The website will be an important public interface of EKWMA and Government of West Bengal on issues related to EKW.

#### *1.7.3 Community Participation*

To enable participation of wetland communities in management, following actions are envisaged:

##### *1.7.3.1 Establishment of a Community Advisory Group*

The Community Advisory Group shall function as a wetland community platform to engage with EKWMA and other line departments on wetland management issues. Specifically, the group shall:

- Articulate community views and suggestions on various management interventions
- Support implementation of various management activities on ground
- Promote awareness on wetland values and functions within community stakeholders
- Support adoption of new technologies aimed at reducing livelihood vulnerabilities and advancing wise use
- Work with EKWMA to promote behaviour change within communities towards wise use of wetlands

The platform meetings shall be convened at least once every six months, and decisions taken shall be placed in the meetings of the EKWMA.

##### *1.7.3.2 Community wardens*

The Community Advisory Group shall be encouraged to appoint community wardens to work as a para-network of wetland monitoring group. The community wardens will be responsible for collecting information on various aspects of wetlands features, and through a mobile-app system, transmit the same to the EKWMA. The community wardens will also be engaged in the awareness and outreach programmes.

#### *1.7.4 Resource material*

Resource material on following themes are proposed to be published:

- East Kolkata Wetlands Ramsar Site
- Traditional Waste Recovery Practice and Wetland Wise Use
- Environmental monitoring system in EKW

- Citizen participation in wetlands management
- Biological diversity of EKW
- People of EKW

#### *1.7.5 Newsletter*

An annual newsletter highlighting progress made in management plan implementation and key emerging issues related to EKW is proposed to be published annually in English and Bengali, and disseminated to all stakeholders.

#### *1.7.6 Workshops and public events*

Public events are proposed to be organized on the eve of World Wetlands Day (Feb 2), World Environment Day (June 5) and International Day for Biological Diversity (May 22) as a means of reaching out to public on the issues of wetland conservation and wise use. Public events on specific issues, as pollution control or water management are also proposed to be organized as a means of engaging with stakeholders.

### **1.8 Monitoring Plan Review**

A mid-term and end-term review of management plan implementation is proposed to assess the extent to which stipulated objectives have been achieved with a high degree of resource efficiency and in participation with stakeholders. A committee on wetlands is proposed to be formed by EKWMA to carry out the evaluation, specifically looking at the following elements:

- Degree to which wetland ecological character is being maintained as a result of management being applied
- Implementation quality, timeliness and resourcing of activities
- Quality and comprehensiveness of wetlands monitoring
- Effectiveness of management being applied, in terms of design, activities, outcomes and impacts
- Quality of stakeholder engagement in implementation of various activities and discharging wetland management functions
- Changes in external environment, requiring adaptation in management plan


## Component 2. Water Management and Pollution Abatement

### 2.1 Dredging of channels

14 highly silted canals measuring 43.8 km are proposed to be dredged to restore their natural flows and enable flow of sewage to different fish farms (Map 18). The desilting will be carried to restore the natural profiles. Other canals may be added as per requirements. EKWMA may request concerned state government departments such as Department of Irrigation & Waterways, Department of Water Resources Investigation and Development for additional budgetary support.

**Table 15. Canals to be desilted in East Kolkata Wetlands**

	District	Canal Name	Length (km)
1	South 24 - Parganas	Bidyadhari -1	7.97
2	South 24 - Parganas	Bidyadhari -2	7.58
3	South 24 - Parganas	Bidyadhari -3	0.89
4	South 24 - Parganas	Khanaberia Khal	0.3
5	South 24 - Parganas	Kharki Khal	2.71
6	South 24 - Parganas	District Board Khal Phase I	2.53
7	South 24 - Parganas	Tihuria Khal	4.45
8	South 24 - Parganas	Chacharia Khal	2.89
9	South 24 - Parganas	Hanakhali Khal	0.99
10	South 24 - Parganas	District Board Khal Phase 2	2.89
11	South 24 - Parganas	Goalbati Khal	1.48
12	North 24 - Parganas	Paran Chaprasi Khal	4.67
13	North 24 - Parganas	Naobhanga Khal (part)	3.38
14	North 24 - Parganas	Ghosher Khal (part)	1.1



**Map 18. Canals proposed to be desilted in East Kolkata Wetlands**

## 2.2 Waste Segregation at EKW inlet

It is proposed to construct a solid waste segregator near Bantala lockgate to regulate flow of solid waste. The exact location of the waste segregator would be decided in consultation with Department of I&W and KMC.

## 2.3 Constructed wetlands at Kulti outfall

The wastewater discharged into Kulti River is not always up to the standards set by Pollution Control Board for discharge into rivers. In order to further augment waste water treatment beyond EKW region, it is proposed to build a system of constructed wetlands prior to discharge into River Kulti. The design of constructed wetlands will be based on EKW, but will use more aquatic vegetation beds through which sewage from DWF would be passed through.

The location of constructed wetlands and design will be finalized in the first year of management plan implementation, and the wetland development work taken up in the second and third year of the management plan, subject to availability of land.

## 2.4 Solid waste management

A plastic waste recycling unit is proposed to be established in the Dhapa region. Recovered plastic may be used to make plastic boards or road construction material. The waste-collectors will form an integral part of the solid waste management system.

## 2.5 Treating Dhapa Leachate

A 300 KLD effluent treatment plant to treat the leachate of Dhapa has been constructed by KMC. The EKWMA will coordinate with KMC to ensure that the effluent is treated to the desired level.

## 2.6 Zero-plastic zone

EKW may be designated a zero-plastic zone. A management framework for this purpose, including components on regulation and behaviour change will be drawn up, in consultation with stakeholders. Screens are proposed to be placed at all inlets to prevent transport of plastic in the canal system of the wetland.



## Component 3. Conservation of Species and Habitats

### 3.1 Asian Waterbird Census

Regular monitoring of waterbird population following standard protocols as the Asian Waterbird Census at all the major congregation sites within and around EKW shall be maintained in a coordinated manner so as to understand comprehensively the significance of the wetlands for waterbirds (both resident and migratory) and to plan and monitor habitat management strategies and actions.

The management plan will support mid-winter counts of waterbird as per Asian Waterbird Census protocol. The census may be carried out in collaboration with NGOs such as Prakriti Samsad, with prior experience in conducting such census. Census programmes will include training of prospective census participants and local waterbird enthusiasts.

Data collected during this monitoring work will include collection of detailed information on habitat structure and seasonal abundance and changes in different prey items (aquatic flora and fauna). Information collected should enable realistic population estimates of the different waterbirds and trends to be developed and provide guidance for management activities.

### 3.2 Bird health

Knowledge of the health of resident and migratory species that inhabit the wetlands is critical to understanding the risk and potential of transmission of avian diseases to other birds and animals, domestic poultry and people. Specific studies need to be undertaken to establish a baseline of the common and potentially harmful avian diseases, including the highly pathogenic strains of avian influenza that have a zoonotic potential.

Under the aegis of management plan, a training on wild bird disease will be provided to staff of Wildlife Department, EKWMA and select NGOs.

### 3.3 Culture of indigenous fish species

Culture of indigenous fish species which are often avoided in culture ponds but are in great demand by the local people, is proposed to be carried out in Goltala bheri which is owned by the government. EKWMA may also invite proposals from the willing fishery owners (govt./private/cooperatives etc.) for technical and financial support from EKWMA for culture of indigenous fish species. The seeds for indigenous fish species such as Koi (*Anabas testudineus*), Magur (*Clarius batrachus*) and Pangus (*Pangasius pangasius*) and others can be produced in the four hatcheries already proposed for the enhancement of fish yield.

### 3.4 Managing invasion of Suckermouth Catfish

Eradication of Suckermouth Catfish is regarded difficult, potentially time consuming and not feasible economically and hence efforts should be focused to contain the current established populations and preventing future introductions. So, to prevent new introduction, EKW farmers are sieving sewage water through multilayer screen before taking it to their ponds and also practicing intensive netting to limit upsurge of existing population. It is proposed to undertake further research on the invasion pathways, and undertake screening at various points along the DWF to prevent further spread of this fish invasive.

### 3.5 Rehabilitation of aquatic vegetation

It is proposed to revegetate peripheral areas of Nalban, Goltala, Captain Bheri and some identified locations with Phragmites, Typha, Shola and other native species to improvise habitat of marsh mongoose, amphibian and reptilian species which prefer impounded water of submerged vegetation for breeding.

There should be complete ban on removal of vegetation from the bheri designated for biodiversity conservation and allow natural regeneration through protective measures. The underground rhizomes, turions and other vegetative parts should be used for propagation of these plant species. It is recommended that economically important plant species should be grown on the banks of bheri.

### 3.6 Communication and education facilities

#### 3.6.1 Interpretation centre

A wetland interpretation centre is proposed to be constructed at Bantala sedimentation tank (defunct) area near Bantala lockgate on the land belonging to KMC and Irrigation and Waterways Department subject to their approval. A 12-hectare land for this purpose has been identified by the KMC, and will be transferred to the EKWMA for this purpose.

The following facilities are proposed to be developed in the interpretation complex:

**Exhibits** including posters, models, flying patterns hanging from ceiling, wetland birds interactive panel and ecosystem food chain.

**Self-learning devices** including touch screen kiosks and maps for self-guided walks.

**Viewing Gallery** comprising panels highlighting the ecological, socioeconomic and cultural aspects. Desks fitted with adequate displays, bird identifications books and wooden benches should be constructed along the gallery.

**Watchtower** to enable visitors to have a view of the EKW landscape. The watchtower would be equipped with binocular facilities.

**Cycling tracks**

**Waste recycling model of East Kolkata Wetlands** indicating treatment of wastes through natural processes and functions of wetlands.

**Children's Play area** consisting of open dioramas and floorings with underwater paintings should be developed as a special section to cater to the young visitors. The area should have several innovative environment-oriented games and interactive food chain, fish trap games, jigsaw puzzles etc.

**Auditorium** having audio-visual facilities for screening documentaries and arranging talks / workshops / meetings.

**Souvenir Shop** for visitors having wetland products, wetland biodiversity replicas, reading materials, photographs, maps for the visitors to take away on payment basis as memorabilia from the visit.

### *3.6.2 Bheri showcase*

It is proposed to develop a showcasing of wastewater utilization in bheri to the visitors. Local communities would be trained to guide visitors about the process of wastewater recovery system and values and function of this special ecosystem. East Kolkata Wetlands Management Authority will develop some models, aquarium, herbariums and boards to describe their operations.

Four to six bheri have been proposed for this purpose - Nalban Bheri, Goltala Bheri, Captian bheri, Charcharia Bheri, Jhograshisa and Gompota Bheri.

### *3.6.3 Guided boat rides*

Restoration of hydrological regimes and the consequent enhancement in water spread area would provide an opportunity for boating for the nature lovers. Guided boat rides shall be arranged for nature lovers to help them explore various aspects of EKW. While cruising through wetlands, the tourists will enjoy the beautiful view of birds as well. Local community groups shall be trained to take up interpretation activities. Guided boat rides are proposed at Nalban Bheri, Goltala Bheri and Charcharia Bheri. Six boats are proposed to be used for these activities.

## Component 4. Sustainable Resource Development

### 4.1 Sustainable Fisheries Development

#### *4.1.1 Bheri desiltation*

It is assessed that production in nearly 30% of bheri under cooperatives are lower due to excessive silt deposition. Under the management plan, a one-time desiltation of bheri under cooperatives would be supported, so that an optimal water column (up to 0.8 m) and natural gradient of water flow can be maintained. The excavated silt will be used to strengthen embankments or used as manure in agricultural farms. The selection of bheri will be done in consultation with the CAG, and would be based on a commitment with the cooperative to maintain the bheri in the good condition to support fish production.

The activities to be undertaken are:

- Meeting of CAG to prepare a list of bheri needing desiltation
- Preparation of estimate of desiltation to be carried out
- Desiltation of the ponds
- Transport of silt to embankments or agricultural farms

#### *4.1.2 Establishment of new hatcheries*

It is proposed to construct 4 hatcheries for production of seeds of Indian major carps and air breathing catfishes in Dhapa Manpur, Tardha Kapasati, Kharki and Kantipota for enhancement of fish yield from East Kolkata Wetlands. The proposed hatcheries with a production capacity of 0.15 million fry / unit for air breathing fishes and 1 million / unit for freshwater fish are proposed to be operated through fish cooperatives.

The revenue generated through sale of fish seeds would be used for operation and maintenance of the hatchery units.

The activities to be undertaken are:

- Formation of SHGs for management of hatcheries and defining rules for management of infrastructure and sharing of usufruct
- Construction of hatchery including brood tank, larval rearing tank, feed culture tank, and generator shed
- Maintenance and hatchery operation including brood stock maintenance, breeding using inducing agents, and larval rearing

#### *4.1.3 Bheri water quality monitoring*

As a part of the wetland monitoring programme under Component I, it is proposed to cover up to 30% of bheri into a water quality monitoring programme. As a part of this programme, each bheri operator will:

- Receive a training on water quality monitoring, covering measurement of parameters and their thresholds
- Receive a water quality testing kit

The water quality data will be uploaded on the monitoring platform through a specially designed app for the purpose. The EKWMA will collate the data and share an overview with the bheri operators on a periodic basis.

## 4.2 Sustainable Agriculture and Horticulture Development

### 4.2.1 Promoting rain water harvesting

Rainwater harvesting can lead to a better utilization of the agricultural land by sustainable intensification and diversification of crops. Several farms have ponds that have become derelict owing to silt deposition.

The management plan shall support desiltation and renovation of the farm ponds with a proper engineering layout up to 1m depth, so that rain water can be utilized in cultivation of low-irrigation rabi crops such as chilli, watermelon, cucurbits, mung bean etc.

### 4.2.2 Diversification of cropping pattern

#### 4.2.2.1 Agriculture

On a demonstration basis, the management plan will support crop diversification in agriculture farms so as to retain agricultural productivity and enhance economic returns to the farmers. Upto 100 farmers will be provided support for cultivation of latest varieties of mung bean, chili and other suitable crops in rabi season.

#### 4.2.2.2 Horticulture

On a demonstration basis, up to 100 horticulture farmers will be provided training in:

- a) Cultivation of fruit crops such as Guava, Ber (Kul) and Sapota (clusters around Bhangor I and II are suited)
- b) Cultivation of high-value vegetables such as green and yellow capsicum and ornamental cabbage
- c) Floriculture (Jasmine, Marigold and Sunflower)
- d) Medicinal plants (Holy Basil, Ghritokumari and others)

Farmers will be provided training for each of these techniques, and the cost of inputs as well as marketing support will be provided, with support of Krishi Vigyan Kendra and Extension Units of Agriculture Department. Identification of beneficiaries will be done by CAG.

## 4.3 Community infrastructure

### 4.3.1 Community Health camps

Health camps for wetlands communities will be run in collaboration with the Department of Health and Family Welfare to address the health hazards.

### 4.3.2 Safe drinking water

Arsenic filters would be provided to the wetland communities to reduce the risk of contracting water borne diseases.

## 8 Budget and Financing

---

### 8.1 Budget

Management plan implementation will entail a budget of Rs. 110.7198 Crore. Of this, the component on institutions and governance is allocated 31.72%. This is followed by allocation of 26.80% for implementing actions under component for sustainable resource development. The components on water management and pollution abatement and conservation of species and habitats have been allocated 21.98% and 19.50% of the budget respectively.

**Table 16. Summary of budget**

S. No	Management Plan Component	Budget (In Lakhs)
1	Institutions and Governance	3511.95
2	Water Management and Pollution Abatement	2433.53
3	Conservation of Species and Habitat	2159.00
4	Sustainable resource development	2967.50
	<b>Grand Total</b>	<b>11071.98</b>

EKWMA shall have the authority to change/revise the indicative allocations in this budget proposal, depending upon the circumstances and considering the "needs" of the wetlands management.

The implementation of management plan is proposed to be through funds provided by the MoEFCC (under NPCA), Government of West Bengal (in the form of state's share) and funds leveraged by building convergence with ongoing schemes of other departments. Table 17 provides an overview of analysis of convergence funding and Table 18 gives detailed activity wise budget.

**Table 17. Analysis of convergence funding**

Management Plan Component	Activity	Total Budget	Funds from Central Government Scheme (Scheme name)	Funds from State Government	Funds from other donors (Project and donor name)	Funds from private sector (Name of the agency)	Funds available from convergence sources	Funds required to be raised
		(a)	(b)	(c)	(d)	(e)	(f)= (b)+(c)+(d)+(e)	(g)=(a)-(f)
<b>Institutions and Governance</b>		<b>3511.95</b>	<b>0.00</b>	<b>3171.95</b>	<b>0.00</b>	<b>0.00</b>	<b>3171.95</b>	<b>340.00</b>
	EKWMA Reorganization and Strengthening	35.00	0.00	35.00	0.00	0.00	35.00	0.00
	Wetland demarcation	64.45	0.00	64.45	0.00	0.00	64.45	0.00
	Management zoning	52	0.00	52	0.00	0.00	52.00	0.00
	Wetlands Inventory, Assessment and Monitoring System	2410.5	0.00	2070.5	0.00	0.00	2070.50	340.00
	Research	232.5	0.00	232.5	0.00	0.00	232.50	0.00
	Capacity Development	57.5	0.00	57.5	0.00	0.00	57.50	0.00
	Communication and Outreach	610	0.00	610	0.00	0.00	610.00	0.00
	Management Plan Implementation Review	50	0.00	50	0.00	0.00	50.00	0.00
<b>Water Management and Pollution Abatement</b>		<b>2433.53</b>	<b>0.00</b>	<b>663.03</b>	<b>0.00</b>	<b>0.00</b>	<b>663.03</b>	<b>1770.50</b>
	Dredging of channels	266.53	0.00	266.53	0.00	0.00	266.53	0.00
	Waste segregation unit at Bantala	50.00	0.00	50.00	0.00	0.00	50.00	0.00
	Constructed wetlands at Kutli outfall	65.00	0.00	0.00	0.00	0.00	0.00	65.00
	Plastic waste recycling unit	125.00	0.00	125.00	0.00	0.00	125.00	0.00
	Treating Dhapa leachate and solid waste management	1905.50	0.00	200.00	0.00	0.00	200.00	1705.50
	Establishing plastic free zone	21.50	0.00	21.50	0.00	0.00	21.50	0.00
<b>Conservation of Species and Habitat</b>		<b>2159.00</b>	<b>0.00</b>	<b>126.50</b>	<b>0.00</b>	<b>0.00</b>	<b>126.50</b>	<b>2032.50</b>
	Asian waterbird census	20.00	0.00	20.00	0.00	0.00	20.00	0.00
	Bird health conservation	3.00	0.00	3.00	0.00	0.00	3.00	0.00
	Culture of indigenous fish species	20.00	0.00	20.00	0.00	0.00	20.00	0.00
	Management of invasive species	30.00	0.00	30.00	0.00	0.00	30.00	0.00
	Rehabilitation of aquatic vegetation	10.00	0.00	10.00	0.00	0.00	10.00	0.00
	Communication and education facilities	2076.00	0.00	43.50	0.00	0.00	43.50	2032.50
<b>Sustainable resource development</b>		<b>2967.50</b>	<b>0.00</b>	<b>467.50</b>	<b>0.00</b>	<b>0.00</b>	<b>467.50</b>	<b>2500.00</b>
	Sustainable fisheries Development	2637.50	0.00	137.50	0.00	0.00	137.50	2500.00
	Sustainable agriculture and horticulture development	105.00	0.00	105.00	0.00	0.00	105.00	0.00
	Community infrastructure	225.00	0.00	225.00	0.00	0.00	225.00	0.00
<b>Grand Total</b>		<b>11071.98</b>	<b>0.00</b>	<b>4428.98</b>	<b>0.00</b>	<b>0.00</b>	<b>4428.98</b>	<b>6643.00</b>

**Table 18. Detailed activity wise budget**

CGS/ GOWB	Core/Non Core	Activity	Sub activity	Physical Target	Unit	Rate	Amount (Rs Lakh)	Year 1		Year 2		Year 3		Year 4		Year 5			
								Physical Target	Amount	Physical Target	Amount	Physical Target	Amount	Physical Target	Amount	Physical Target	Amount		
<b>Component I: Institutions and Governance</b>								<b>3511.95</b>	<b>1,006.30</b>		<b>2,153.72</b>		<b>181.92</b>		<b>72.50</b>		<b>97.50</b>		
<b>1.1 EKWMA Reorganization and Strengthening</b>																			
GOWB	C		1.1.1 Reorganization proposal		1 study	30,00,000	30				30.00			-		-	-		
GOWB	NC		1.1.2 Stakeholder workshop		1 workshop	5,00,000	5			5.00				-		-	-		
<b>1.2 Wetland demarcation</b>																			
GOWB	C		1.2.1 Field survey for identification of pillar points		1600	1,225	19.6	800.00	9.80	800.00	9.80								
GOWB	C		1.2.2 Placement of pillars		1600 pillars	1,553	24.848			800.00	12.42	800.00	12.42			-	-		
GOWB	NC		1.2.3 Replacement and maintenance			LS	20								10.00		10.00		
<b>1.3 Management zoning</b>																			
GOWB	NC		1.3.1 Stakeholder workshop		1 workshop	2,00,000	2	1.00	2.00					-		-	-		
GOWB	C		1.3.2 Preparation of zonal land use management plans including inventory of land ownership		1 plan	50,00,000	50		20.00	1.00	30.00			-		-	-		
<b>1.4 Wetlands Inventory, Assessment and Monitoring System</b>																			
<b>1.4.1 Establishment of wetland monitoring and research centre</b>																			
CGS	NC		Upgradation of Laboratory and wetland monitoring centre			LS	200		100.00		100.00								
CGS	NC		Research associates (staff cost)		4 persons / annu	24,00,000	120	4.00	24.00	4.00	24.00	4.00	24.00	4.00	24.00	4.00	24.00		
<b>1.4.2 Development of wetlands database management system</b>																			
CGS	C		Database design and development		1 Consultancy	LS	20		20.00										
CGS	NC		Maintenance																
<b>1.4.3 Wetlands monitoring and evaluation</b>																			
GOWB	NC		Surveillance system including helpline																
GOWB	NC		System design		1 consultancy	1000000	10	1.00	10										
GOWB	C		Infrastructure procurement (CCTV, Drones) (including O&M)				2000		500		1500								
GOWB	C		App design		1 consultancy	1000000	10	1.00	10										
GOWB	NC		Travel and Contingencies		per annum	600000	30		6		6		6.00		6.00		6.00		
GOWB	NC		Annual monitoring report publication		5 publication	200000	10	1.00	2	1.00	2	1.00	2.00	1.00	2.00	1.00	2.00		
<b>1.4.4 Ecosystem Health Report Card</b>																			
GOWB	NC		Methods workshop		1 workshop	1,00,000	1	1.00	1.00										
GOWB	C		Report preparation and publication		3 Report Cards	1,50,000	4.5	1.00	1.50			1.00	1.50			1.00	1.50		
GOWB	NC		Stakeholder workshop		2 workshops	2,50,000	5					1.00	2.50			1.00	2.50		
<b>1.5 Research</b>																			
<b>1.5.1 Carbon and GHG flux assessment and nutrient budget</b>																			
GOWB	C		Inception workshop		1 workshop	1,50,000	1.5	1.00	1.50										
GOWB	C		Study		1 study	40,00,000	40		10.00		20.00		10.00						
GOWB	NC		Result sharing workshop		1 workshop	2,50,000	2.5					1.00	2.50						
GOWB	C		Publication		1 publication	2,50,000	2.5					1.00	2.50						
<b>1.5.2 Hydrological assessment (volume of sewage inflow and study on canals and waterbodies management)</b>																			
GOWB	C		Inception workshop		1 workshop	1,50,000	1.5	1.00	1.50										
GOWB	C		Study		1 study	40,00,000	40		10.00		20.00		10.00						
GOWB	NC		Result sharing workshop		1 workshop	2,50,000	2.5					1.00	2.50						
GOWB	C		Publication		1 publication	2,50,000	2.5					1.00	2.50						



Core/Non Core	Activity	Sub activity	Physical Target	Unit	Rate	Amount (Rs Lakh)	Year 1		Year 2		Year 3		Year 4		Year 5	
							Physical Target	Amount	Physical Target	Amount	Physical Target	Amount	Physical Target	Amount	Physical Target	Amount
		1.5.3 Bioaccumulation of heavy metals studies and locating pollution pathways and illegal industrial units														
C		Inception workshop	1	workshop	1,50,000	1.5	1.00	1.50								
C		Study	1	study	40,00,000	40		10.00	20.00		10.00					
NC		Result sharing workshop	1	workshop	2,50,000	2.5				1.00	2.50					
C		Publication	1	publication	2,50,000	2.5				1.00	2.50					
		1.5.4 Assessment of production of fish and fish disease and feasibility study of certification possible for fish produced in EKW, Assessment of agriculture (paddy), horticulture, medicinal Plants, floriculture and uses														
C		Inception workshop	2	workshop	1,50,000	3	2.00	3.00								
C		Study	2	study	40,00,000	80		20.00	40.00		20.00					
NC		Result sharing workshop	2	workshop	2,50,000	5				2.00	5.00					
C		Publication	2	publication	2,50,000	5				2.00	5.00					
		<b>1.6 Capacity Development</b>														
NC		1.6.1 Training Needs Assessment	1	study	10,00,000	10	1.00	10.00								
NC		1.6.2 Training workshops														
		1.6.2.1 For EKWMA and concerned state government agencies														
NC		Integrated wetlands management	1	training	5,00,000	5	1.00	5.00								
NC		Wetlands inventory, assessment and monitoring	1	training	5,00,000	5			1.00	5.00						
NC		Wetland and climate change adaptation	1	training	5,00,000	5				1.00	5.00					
NC		Integrating wetlands in developmental planning	1	training	5,00,000	5						1.00	5.00			
NC		1.6.2.2 For NGOs and CBOs	5	trainings	2,50,000	12.5	1.00	2.50	1.00	2.50	1.00	2.50	1.00	2.50	1.00	2.50
NC		1.6.2.3 For communities	10	trainings	1,50,000	15	2.00	3.00	2.00	3.00	2.00	3.00	2.00	3.00	2.00	3.00
		<b>1.7 Communication and Outreach</b>														
C		1.7.1 Signage	50	units	10,00,000	500	20.00	200.00	30.00	300.00						
C		1.7.2 Webpage	1	consultancy	10,00,000	10	1.00	10.00								
C		1.7.3 Community engagement meetings	20	meetings	50,000	10	4.00	2.00	4.00	2.00	4.00	2.00	4.00	2.00	4.00	2.00
C		1.7.4 Resource material	10	publications	4,00,000	40			3.00	12.00	4.00	16.00	2.00	8.00	1.00	4.00
C		1.7.5 Newsletter	5	publications	4,00,000	20	1.00	4.00	1.00	4.00	1.00	4.00	1.00	4.00	1.00	4.00
C		1.7.6 Public Events	15	events	2,00,000	30	3.00	6.00	3.00	6.00	3.00	6.00	3.00	6.00	3.00	6.00
		<b>1.8 Management Plan Implementation Review</b>														
NC		1.8.1 Mid-term review	1	assessment	20,00,000	20					1.00	20.00				
NC		1.8.2 End-term review	1	assessment	30,00,000	30									1.00	30.00
		<b>Component 2: Water Management and Pollution Abatement</b>				<b>2433.53</b>	<b>14.70</b>	<b>801.72</b>	<b>942.27</b>	<b>613.34</b>	<b>61.50</b>					
		<b>2.1 Dredging of channels</b>														
C		2.1.1 Field surveys and preparation of estimates	44	km	6,000	2.64	20.00	1.20	24.00	1.44						
C		2.1.2 Dredging	44	km	599748.002	263.89			20.00	119.95	24.00	143.94				
		<b>2.2 Waste segregation unit at Bantala</b>														
C		2.2.2 Installation of waste segregation unit	1	unit	50,00,000	50			-	1.00	50.00					
		<b>2.3 Constructed wetlands at Kutli outfall</b>														
C		2.3.1 Design and Estimation			LS	5				5.00						
C		2.3.2 Creation of constructed wetlands	20	ha	3,00,000	60					20.00	60.00				

Core/Non Core	Activity	Sub activity	Physical Target	Unit	Rate	Amount (Rs Lakh)	Year 1		Year 2		Year 3		Year 4		Year 5	
							Physical Target	Amount	Physical Target	Amount	Physical Target	Amount	Physical Target	Amount	Physical Target	Amount
		<b>2.4 Plastic waste recycling unit</b>														
C		2.4.1 Design and Estimation			LS	5				5.00						
C		2.4.2 Installation of waste recycling unit			LS	120					120.00					
		<b>2.5 Treating Dhapa leachate and solid waste management</b>														
C		2.5.1 Anaerobic biological treatment				50		10.00		10.00		10.00		10.00		10.00
C		2.5.2 Solid Waste Management and Biomining	236500	lakh tonne	700	1655.5			78,833.00	551.83	78,833.00	551.83	78,834.00	551.84		
C		2.5.3 Solid Waste Management at GP level	4	GP	50,00,000	200			1.00	50.00	1.00	50.00	1.00	50.00	1.00	50.00
		<b>2.6 Establishing plastic free zone</b>														
NC		2.6.1 Sensitization workshop	15	workshops	50,000	7.5	3.00	1.50	3.00	1.50	3.00	1.50	3.00	1.50	3.00	1.50
NC		2.6.2 Plastic waste collection units	10	units	60,000	6			5.00	3.00	5.00	3.00				
NC		2.6.3 Outreach material	20	publications	40,000	8	5.00	2.00	10.00	4.00	5.00	2.00				
		<b>Component 3: Conservation of Species and Habitat</b>				<b>2159.00</b>		<b>16.50</b>		<b>1077.00</b>		<b>1048.50</b>		<b>13.00</b>		<b>4.00</b>
		<b>3.1 Asian waterbird census</b>														
NC		3.1.1 Training	5	trainings	1,50,000	7.5	1.00	1.50	1.00	1.50	1.00	1.50	1.00	1.50	1.00	1.50
C		3.1.2 Annual Census	5	census	2,00,000	10	1.00	2.00	1.00	2.00	1.00	2.00	1.00	2.00	1.00	2.00
C		3.1.3 Reporting	5	reports	50,000	2.5	1.00	0.50	1.00	0.50	1.00	0.50	1.00	0.50	1.00	0.50
		<b>3.2 Bird health conservation</b>														
NC		3.2.1 Training workshops for wild bird disease management	2	workshops	1,50,000	3			1.00	1.50			1.00	1.50		
C		3.2.2 Establishing baseline														
C		3.2.3 Surveillance and reporting														
		<b>3.3 Culture of indigenous fish species</b>														
C		3.3.1 Feasibility assessment	1	study	10,00,000	10	1.00	5.00		5.00		-		-		-
NC		3.3.2 Training for selective breeding	2	trainings	2,50,000	5			1.00	2.50	1.00	2.50		-		-
C		3.3.3 Establishing fish stock				5						5.00		-		-
C		3.3.4 Surveillance and reporting														
		<b>3.4 Management of invasive species</b>														
C		3.4.1 Study on species pathways	1	study	15,00,000	15	1.00	7.50		7.50		-		-		-
NC		3.4.2 Screening of pathways and eradication				15				5.00		5.00		5.00		-
		<b>3.5 Rehabilitation of aquatic vegetation</b>														
C		3.5.1 Site assessment	1	study		5					1.00	5.00		-		-
C		3.5.2 Vegetation regeneration / reestablishment	10	ha	50,000	5					5.00	2.50	5.00	2.50		-
		<b>3.6 Communication and education facilities</b>														
C		3.6.1 Interpretation centre	10,000	m2	20,000	2000				1,000.00		1,000.00		-		-
C		Exhibits												-		-
C		Viewing gallery												-		-
NC		Watchtower	3	units	2,50,000	7.5					3.00	7.50		-		-
C		Nature trails	2	units	12,50,000	25			1.00	12.50	1.00	12.50		-		-
NC		Waste recycling model of EKW												-		-
NC		Children play area												-		-
C		Auditorium												-		-
NC		Souvenir shop												-		-
		3.6.2 Bheri showcase												-		-
NC		Training of community members as guides	3	trainings	1,50,000	4.5			2.00	3.00	1.00	1.50		-		-
NC		Establishing bheri models	3	bheri	2,00,000	6			2.00	4.00	1.00	2.00		-		-
		3.6.3 Guided boat rides														
NC		Purchase of boats	6	boat	5,00,000	30				6.00	30.00			-		-
NC		Training of boatmen	3	trainings	1,00,000	3			2.00	2.00	1.00	1.00		-		-

Core/Non Core	Activity	Sub activity	Physical Target	Unit	Rate	Amount (Rs Lakh)	Year 1		Year 2		Year 3		Year 4		Year 5	
							Physical Target	Amount	Physical Target	Amount	Physical Target	Amount	Physical Target	Amount	Physical Target	Amount
<b>Component 4: Sustainable resource development</b>						<b>2,967.50</b>		<b>542.50</b>		<b>1,643.00</b>		<b>620.50</b>		<b>92.00</b>		<b>69.50</b>
		<b>4.1 Sustainable fisheries Development</b>														
		4.1.1 Bheri desiltation														
C		Meeting of CAG to prepare a list of bheri needing desiltation	Costed within community consultation meetings													
C		Desiltation estimate preparation														
C		Bheri Desiltation	25	bheri	1,00,00,000	2500	5.00	500.00	15.00	1,500.00	5.00	500.00	-	-	-	-
C		Transportation of silt to embankments or agricultural farms						-	-	-	-	-	-	-	-	-
		4.1.2 Establishment of new hatchery and fish feed marketing unit														
C		Formation of SHG														
NC		Training workshops	4	trainings	1,50,000	6	-	-	4.00	6.00	-	-	-	-	-	-
NC		Fish hatchery establishment and fish feed marketing units	4	units	10,00,000	40	-	-	-	-	2.00	20.00	2.00	20.00	-	-
NC		Operation and maintenance	4	units	6,00,000	24	-	-	-	-	-	-	2.00	12.00	2.00	12.00
		4.1.3 Bheri water quality and fish monitoring														
NC		Training on water quality monitoring and fish disease management	5	trainings	1,50,000	7.5	-	-	3.00	4.50	2.00	3.00	-	-	-	-
NC		Provision of water quality kits	500	kits	10,000	50	-	-	300.00	30.00	200.00	20.00	-	-	-	-
NC		App for data collection and management	1	app		10	1.00	10.00	-	-	-	-	-	-	-	-
		<b>4.2 Sustainable agriculture and horticulture development</b>														
		4.2.1 Promoting rain water harvesting														
NC		Training	10	trainings	1,50,000	15	-	-	5.00	7.50	5.00	7.50	-	-	-	-
NC		Unit establishment	10	clusters	2,50,000	25	-	-	5.00	12.50	5.00	12.50	-	-	-	-
NC		Handholding support				5	-	-	-	-	-	-	2.50	2.50	2.50	2.50
		4.2.2 Diversification of cropping pattern														
NC		Training on cultivation of crops	10	trainings	1,50,000	15	5.00	7.50	5.00	7.50	-	-	-	-	-	-
NC		Demonstration plots	10	ha	4,00,000	40	5.00	20.00	5.00	20.00	-	-	2.50	2.50	2.50	2.50
NC		Handholding support				5	-	-	-	-	-	-	-	-	-	-
		<b>4.3 Community infrastructure</b>														
C		4.3.1 Health camps	5	camps	5,00,000	25	1.00	5.00	1.00	5.00	1.00	5.00	1.00	5.00	1.00	5.00
C		4.3.2 Safe drinking water, sanitation and nutrition management	100	units	2,00,000	200	-	-	25.00	50.00	25.00	50.00	25.00	50.00	25.00	50.00
<b>Grand Total</b>						<b>11,071.98</b>		<b>1,580.00</b>		<b>5,675.44</b>		<b>2,793.19</b>		<b>790.84</b>		<b>232.50</b>

Activity	Funds required from Central Government Scheme (CGS) (Rs. in Lakh)	%	Funds required from State Government Scheme (GOWB) (Rs. in Lakh)	%	Amount (Rs. in lakh)
Core	6315.50	95.07	3852.98	86.99	10168.48
Non Core	327.50	4.93	576.00	13.01	903.50
<b>Total</b>	<b>6643.00</b>	<b>100.00</b>	<b>4428.98</b>	<b>100.00</b>	<b>11,071.98</b>

## 8.2 Financing

Yearwise breakup of requirement of funds in terms of core and non-core activities as required from NPCA and Government of West Bengal are provided in Table 19 and 20 respectively.

**Table 19. Yearwise breakup of requirement of funds required from NPCA**

	Core/ Non Core	Activity	Sub activity	Year 1	Year 2	Year 3	Year 4	Year 5	Amount (Rs Lakh)
				Amount	Amount	Amount	Amount	Amount	
		<b>Component 1: Institutions and Governance</b>		<b>144.00</b>	<b>124.00</b>	<b>24.00</b>	<b>24.00</b>	<b>24.00</b>	<b>340</b>
		1.4	Wetlands Inventory, Assessment and Monitoring System						
		1.4.1	Establishment of wetland monitoring and research centre						
<b>CGS</b>	<b>NC</b>		Upgradation of Laboratory and wetland monitoring centre	100.00	100.00	-	-	-	200
<b>CGS</b>	<b>NC</b>		Research associates (staff cost)	24.00	24.00	24.00	24.00	24.00	120
		1.4.2	Development of wetlands database management system						
<b>CGS</b>	<b>C</b>		Database design and development	20.00	-	-	-	-	20
<b>CGS</b>	<b>NC</b>		Maintenance						
		<b>Component 2: Water Management and Pollution Abatement</b>		<b>10</b>	<b>566.83</b>	<b>621.83</b>	<b>561.84</b>	<b>10</b>	<b>1770.5</b>
		2.3	Constructed wetlands at Kutli outfall						
<b>CGS</b>	<b>C</b>	2.3.1	Design and Estimation	-	5.00	-	-	-	5
<b>CGS</b>	<b>C</b>	2.3.2	Creation of constructed wetlands	-	-	60.00	-	-	60
		2.5	Treating Dhapa leachate and solid waste management						
<b>CGS</b>	<b>C</b>	2.5.1	Anaerobic biological treatment	10.00	10.00	10.00	10.00	10.00	50
<b>CGS</b>	<b>C</b>	2.5.2	Solid Waste Management and Biomining	-	551.83	551.83	551.84	-	1655.5

	Core/ Non Core	Activity	Sub activity	Year 1	Year 2	Year 3	Year 4	Year 5	Amount (Rs Lakh)
	<b>Component 3: Conservation of Species and Habitat</b>			-	1,012.50	1,020.00	-	-	2032.5
		3.6	Communication and education facilities						
CGS	C		3.6.1 Interpretation centre	-	1,000.00	1,000.00	-	-	2000
CGS	C		Exhibits	-	-	-	-	-	
CGS	C		Viewing gallery	-	-	-	-	-	
CGS	NC		Watchtower	-	-	7.50	-	-	7.5
CGS	C		Nature trails	-	12.50	12.50			25
CGS	NC		Waste recycling model of EKW	-	-	-	-	-	
CGS	NC		Children play area	-	-	-	-	-	
CGS	C		Auditorium	-	-	-	-	-	
CGS	NC		Souvenir shop	-	-	-	-	-	
	<b>Component 4: Sustainable resource development</b>			500.00	1,500.00	500.00	-	-	2,500.00
		4.1	Sustainable fisheries Development						
			4.1.1 Bheri desiltation						
CGS	C		Meeting of CAG to prepare a list of bheri needing desiltation						
CGS	C		Desiltation estimate preparation						
CGS	C		Bheri Desiltation	500.00	1,500.00	500.00	-	-	2500
CGS	C		Transportation of silt to embankments or agricultural farms	-	-	-	-	-	
		<b>Grand Total</b>		<b>654.00</b>	<b>3,203.33</b>	<b>2,165.83</b>	<b>585.84</b>	<b>34.00</b>	<b>6,643.00</b>

**Table 20. Yearwise breakup of requirement of funds required from Government of West Bengal**

GOWB	Core/ Non Core	Activity	Sub activity	Year 1	Year 2	Year 3	Year 4	Year 5	Amount (Rs Lakh)
				Amount	Amount	Amount	Amount	Amount	
<b>Component I: Institutions and Governance</b>				<b>862.30</b>	<b>2,029.72</b>	<b>157.92</b>	<b>48.50</b>	<b>73.50</b>	<b>3,171.948</b>
		<b>1.1</b>	<b>EKWMA Reorganization and Strengthening</b>						
GOWB	C		1.1.1 Reorganization proposal		30.00	-	-	-	30.00
GOWB	NC		1.1.2 Stakeholder workshop		5.00	-	-	-	5.00
		<b>1.2</b>	<b>Wetland demarcation</b>						
GOWB	C		1.2.1 Field survey for identification of pillar points	9.80	9.80				19.60
GOWB	C		1.2.2 Placement of pillars	-	12.42	12.42	-	-	24.848
GOWB	NC		1.2.3 Replacement and maintenance				10.00	10.00	20.00
		<b>1.3</b>	<b>Management zoning</b>						
GOWB	NC		1.3.1 Stakeholder workshop	2.00	-	-	-	-	2.00
GOWB	C		1.3.2 Preparation of zonal land use management plans including inventory of land ownership	20.00	30.00	-	-	-	50.00
		<b>1.4</b>	<b>Wetlands Inventory, Assessment and Monitoring System</b>						
			1.4.3 Wetlands monitoring and evaluation						
GOWB	NC		Surveillance system including helpline						
GOWB	NC		System design	10					10.00
GOWB	C		Infrastructure procurement (CCTV, Drones) (including O&M)	500	1500				2,000.00
GOWB	C		App design	10					10.00
GOWB	NC		Travel and Contingencies	6	6	6.00	6.00	6.00	30.00
GOWB	NC		Annual monitoring report publication	2	2	2.00	2.00	2.00	10.00
			1.4.4 Ecosystem Health Report Card						
GOWB	NC		Methods workshop	1					1.00
GOWB	C		Report preparation and publication	1.5		1.50		1.50	4.50
GOWB	NC		Stakeholder workshop			2.50		2.50	5.00
		<b>1.5</b>	<b>Research</b>						
			1.5.1 Carbon and GHG flux assessment and nutrient budget						
GOWB	C		Inception workshop	1.50					1.50
GOWB	C		Study	10.00	20.00	10.00			40.00
GOWB	NC		Result sharing workshop			2.50			2.50
GOWB	C		Publication			2.50			2.50
			1.5.2 Hydrological assessment (volume of sewage inflow and study on canals and waterbodies management)						
GOWB	C		Inception workshop	1.50					1.50
GOWB	C		Study	10.00	20.00	10.00			40.00
GOWB	NC		Result sharing workshop			2.50			2.50
GOWB	C		Publication			2.50			2.50

	Core/ Non Core	Activity	Sub activity	Year 1	Year 2	Year 3	Year 4	Year 5	Amount (Rs Lakh)
			1.5.3 Bioaccumulation of heavy metals studies and locating pollution pathways and illegal industrial units						
GOWB	C		Inception workshop	1.50					1.50
GOWB	C		Study	10.00	20.00	10.00			40.00
GOWB	NC		Result sharing workshop			2.50			2.50
GOWB	C		Publication			2.50			2.50
			1.5.4 Assessment of production of fish and fish disease and feasibility study of certification possible for fish produced in EKW, Assessment of agriculture (paddy), horticulture, medicinal Plants, floriculture and uses						
GOWB	C		Inception workshop	3.00					3.00
GOWB	C		Study	20.00	40.00	20.00			80.00
GOWB	NC		Result sharing workshop			5.00			5.00
GOWB	C		Publication			5.00			5.00
			<b>1.6 Capacity Development</b>						
GOWB	NC		1.6.1 Training Needs Assessment	10.00					10.00
GOWB	NC		1.6.2 Training workshops					-	
			1.6.2.1 For EKVMA and concerned state government agencies						
GOWB	NC		Integrated wetlands management	5.00					5.00
GOWB	NC		Wetlands inventory, assessment and monitoring		5.00				5.00
GOWB	NC		Wetland and climate change adaptation			5.00			5.00
GOWB	NC		Integrating wetlands in developmental planning				5.00		5.00
GOWB	NC		1.6.2.2 For NGOs and CBOs	2.50	2.50	2.50	2.50	2.50	12.50
GOWB	NC		1.6.2.3 For communities	3.00	3.00	3.00	3.00	3.00	15.00
			<b>1.7 Communication and Outreach</b>						
GOWB	C		1.7.1 Signage	200.00	300.00				500.00
GOWB	C		1.7.2 Webpage	10.00	-				10.00
GOWB	C		1.7.3 Community engagement meetings	2.00	2.00	2.00	2.00	2.00	10.00
GOWB	C		1.7.4 Resource material	-	12.00	16.00	8.00	4.00	40.00
GOWB	C		1.7.5 Newsletter	4.00	4.00	4.00	4.00	4.00	20.00
GOWB	C		1.7.6 Public Events	6.00	6.00	6.00	6.00	6.00	30.00
			<b>1.8 Management Plan Implementation Review</b>						
GOWB	NC		1.8.1 Mid-term review			20.00			20.00
GOWB	NC		1.8.2 End-term review					30.00	30.00

	Core/ Non Core	Activity	Sub activity	Year 1	Year 2	Year 3	Year 4	Year 5	Amount (Rs Lakh)
				Amount	Amount	Amount	Amount	Amount	
<b>Component 2: Water Management and Pollution Abatement</b>				<b>4.7</b>	<b>234.89</b>	<b>320.44</b>	<b>51.5</b>	<b>51.5</b>	<b>663.03</b>
<b>GOWB</b>		<b>2.1</b>	<b>Dredging of channels</b>						
	<b>C</b>		2.1.1 Field surveys and preparation of estimates	1.20	1.44				2.64
<b>GOWB</b>	<b>C</b>		2.1.2 Dredging		119.95	143.94			263.89
<b>GOWB</b>		<b>2.2</b>	<b>Waste segregation unit at Bantala</b>						
	<b>C</b>		2.2.2 Installation of waste segregation unit	-	50.00	-	-	-	50.00
<b>GOWB</b>		<b>2.4</b>	<b>Plastic waste recycling unit</b>	-	-	-	-	-	
<b>GOWB</b>	<b>C</b>		2.4.1 Design and Estimation		5.00				5.00
	<b>C</b>		2.4.2 Installation of waste recycling unit			120.00			120.00
<b>GOWB</b>		<b>2.5</b>	<b>Treating Dhapa leachate and solid waste management</b>						
	<b>C</b>		2.5.3 Solid Waste Management at GP level		50.00	50.00	50.00	50.00	200.00
<b>GOWB</b>		<b>2.6</b>	<b>Establishing plastic free zone</b>						
<b>GOWB</b>	<b>NC</b>		2.6.1 Sensitization workshop	1.50	1.50	1.50	1.50	1.50	7.50
	<b>NC</b>		2.6.2 Plastic waste collection units		3.00	3.00			6.00
	<b>NC</b>		2.6.3 Outreach material	2.00	4.00	2.00			8.00
<b>Component 3: Conservation of Species and Habitat</b>				<b>16.50</b>	<b>64.50</b>	<b>28.50</b>	<b>13.00</b>	<b>4.00</b>	<b>126.50</b>
<b>GOWB</b>		<b>3.1</b>	<b>Asian waterbird census</b>						
<b>GOWB</b>	<b>NC</b>		3.1.1 Training	1.50	1.50	1.50	1.50	1.50	7.50
	<b>C</b>		3.1.2 Annual Census	2.00	2.00	2.00	2.00	2.00	10.00
	<b>C</b>		3.1.3 Reporting	0.50	0.50	0.50	0.50	0.50	2.50
<b>GOWB</b>		<b>3.2</b>	<b>Bird health conservation</b>						
<b>GOWB</b>	<b>NC</b>		3.2.1 Training workshops for wild bird disease management	-	1.50		1.50		3.00
	<b>C</b>		3.2.2 Establishing baseline	-	-		-		
	<b>C</b>		3.2.3 Surveillance and reporting	-	-		-		
<b>GOWB</b>		<b>3.3</b>	<b>Culture of indigenous fish species</b>						
<b>GOWB</b>	<b>C</b>		3.3.1 Feasibility assessment	5.00	5.00	-	-	-	10.00
<b>GOWB</b>	<b>NC</b>		3.3.2 Training for selective breeding	-	2.50	2.50	-	-	5.00
	<b>C</b>		3.3.3 Establishing fish stock	-	-	5.00	-	-	5.00
	<b>C</b>		3.3.4 Surveillance and reporting						



	Core/ Non Core	Activity	Sub activity	Year 1	Year 2	Year 3	Year 4	Year 5	Amount (Rs Lakh)
GOWB									
GOWB		3.4	Management of invasive species						
	C	3.4.1	Study on species pathways	7.50	7.50	-	-	-	15.00
	NC	3.4.2	Screening of pathways and eradication	-	5.00	5.00	5.00	-	15.00
GOWB									
GOWB		3.5	Rehabilitation of aquatic vegetation						
	C	3.5.1	Site assessment	-	-	5.00	-	-	5.00
	C	3.5.2	Vegetation regeneration / reestablishment	-	-	2.50	2.50	-	5.00
GOWB									
GOWB		3.6.2	Bhery showcase	-	-	-	-	-	
	NC		Training of community members as guides	-	3.00	1.50	-	-	4.50
	NC		Establishing bhery models	-	4.00	2.00	-	-	6.00
GOWB									
GOWB		3.6.3	Guided boat rides						
	NC		Purchase of boats	-	30.00	-	-	-	30.00
	NC		Training of boatmen	-	2.00	1.00	-	-	3.00
			<b>Component 4: Sustainable resource development</b>	<b>42.50</b>	<b>143.00</b>	<b>120.50</b>	<b>92.00</b>	<b>69.50</b>	<b>467.50</b>
GOWB		4.1	Sustainable fisheries Development						
GOWB		4.1.2	Establishment of new hatchery and fish feed marketing unit						
GOWB	C		Formation of SHG						
GOWB	NC		Training workshops	-	6.00	-	-	-	6.00
	NC		Fish hatchery establishment and fish feed marketing units	-		20.00	20.00	-	40.00
	NC		Operation and maintenance	-		-	12.00	12.00	24.00
GOWB									
GOWB		4.1.3	Bheri water quality and fish monitoring						
GOWB	NC		Training on water quality monitoring and fish disease management	-	4.50	3.00	-	-	7.50
	NC		Provision of water quality kits	-	30.00	20.00	-	-	50.00
	NC		App for data collection and management	10.00	-	-	-	-	10.00
GOWB		4.2	Sustainable agriculture and horticulture development						
GOWB		4.2.1	Promoting rain water harvesting						
GOWB	NC		Training	-	7.50	7.50	-	-	15.00
	NC		Unit establishment	-	12.50	12.50	-	-	25.00
	NC		Handholding support	-	-	-	2.50	2.50	5.00
GOWB									
GOWB		4.2.2	Diversification of cropping pattern						
GOWB	NC		Training on cultivation of crops	7.50	7.50	-	-	-	15.00
	NC		Demonstration plots	20.00	20.00				40.00
	NC		Handholding support			2.50	2.50		5.00
GOWB									
		4.3	Community infrastructure						
GOWB	C	4.3.1	Health camps	5.00	5.00	5.00	5.00	5.00	25.00
	C	4.3.2	Safe drinking water, sanitation and nutrition management	-	50.00	50.00	50.00	50.00	200.00
		<b>Grand Total</b>		<b>926.00</b>	<b>2,472.11</b>	<b>627.36</b>	<b>205.00</b>	<b>198.50</b>	<b>4,428.98</b>

## 8.3 Phasing of Activities

**Table 21. Phasing of activities**

Activity	Sub activity	Year 1		Year 2		Year 3		Year 4		Year 5	
		1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half
<b>Component I: Institutions and Governance</b>											
<b>1.1 EKWMA Reorganization and Strengthening</b>											
	1.1.1										
	1.1.1										
	1.1.2										
<b>1.2 Wetland demarcation</b>											
	1.2.1										
	1.2.1										
	1.2.2										
	1.2.3										
<b>1.3 Management zoning</b>											
	1.3.1										
	1.3.1										
	1.3.2										
	1.3.2										
<b>1.4 Wetlands Inventory, Assessment and Monitoring System</b>											
	1.4.1										
	1.4.1										
	1.4.1										
	1.4.2										
	1.4.2										
	1.4.2										
	1.4.3										
	1.4.3										
	1.4.3										
	1.4.3										
	1.4.3										
	1.4.3										
	1.4.3										
	1.4.3										
	1.4.3										
	1.4.4										
	1.4.4										
	1.4.4										
	1.4.4										
	1.4.4										

Activity	Sub activity	Year 1		Year 2		Year 3		Year 4		Year 5	
		1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half
<b>I.5 Research</b>											
1.5.1	Carbon and GHG flux assessment and nutrient budget										
	Inception workshop										
	Study										
	Result sharing workshop										
	Publication										
1.5.2	Hydrological assessment (volume of sewage inflow and study on canals and waterbodies management)										
	Inception workshop										
	Study										
	Result sharing workshop										
	Publication										
1.5.3	Bioaccumulation of heavy metals studies and locating pollution pathways and illegal industrial units										
	Inception workshop										
	Study										
	Result sharing workshop										
	Publication										
1.5.4	Assessment of production of fish and fish disease and feasibility study of certification possible for fish produced in EKW, Assessment of agriculture (paddy), horticulture, medicinal Plants, floriculture and uses										
	Inception workshop										
	Study										
	Result sharing workshop										
	Publication										
<b>I.6 Capacity Development</b>											
1.6.1	Training Needs Assessment										
1.6.2	Training workshops										
1.6.2.1	For EKWMA and concerned state government agencies										
	Integrated wetlands management										
	Wetlands inventory, assessment and monitoring										
	Wetland and climate change adaptation										
	Integrating wetlands in developmental planning										
1.6.2.2	For NGOs and CBOs										
1.6.2.3	For communities										
<b>I.7 Communication and Outreach</b>											
1.7.1	Signage										
1.7.2	Webpage										
1.7.3	Community engagement meetings										
1.7.4	Resource material										
1.7.5	Newsletter										
1.7.6	Public Events										
<b>I.8 Management Plan Implementation Review</b>											
1.8.1	Mid-term review										
1.8.2	End-term review										

Activity	Sub activity	Year 1		Year 2		Year 3		Year 4		Year 5	
		1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half
<b>Component 2: Water Management and Pollution Abatement</b>											
<b>2.1 Dredging of channels</b>											
	2.1.1										
	2.1.2										
<b>2.2 Waste segregation unit at Bantala</b>											
	2.2.2										
<b>2.3 Constructed wetlands at Kutli outfall</b>											
	2.3.1										
	2.3.2										
<b>2.4 Plastic waste recycling unit</b>											
	2.4.1										
	2.4.2										
<b>2.5 Treating Dhapa leachate and solid waste management</b>											
	2.5.1										
	2.5.2										
	2.5.3										
<b>2.6 Establishing plastic free zone</b>											
	2.6.1										
	2.6.2										
	2.6.3										
<b>Component 3: Conservation of Species and Habitat</b>											
<b>3.1 Asian waterbird census</b>											
	3.1.1										
	3.1.2										
	3.1.3										
<b>3.2 Bird health conservation</b>											
	3.2.1										
	3.2.2										
	3.2.3										
<b>3.3 Culture of indigenous fish species</b>											
	3.3.1										
	3.3.2										
	3.3.3										
	3.3.4										
<b>3.4 Management of invasive species</b>											
	3.4.1										
	3.4.2										
<b>3.5 Rehabilitation of aquatic vegetation</b>											
	3.5.1										
	3.5.2										

Activity	Sub activity	Year 1		Year 2		Year 3		Year 4		Year 5	
		1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half	1 st Half	2 nd Half
<b>3.6 Communication and education facilities</b>											
3.6.1	Interpretation centre										
	Exhibits										
	Viewing gallery										
	Watchtower										
	Nature trails										
	Waste recycling model of EKW										
	Children play area										
	Auditorium										
	Souvenir shop										
3.6.2	Bhery showcase										
	Training of community members as guides										
	Establishing bhery models										
3.6.3	Guided boat rides										
	Purchase of boats										
	Training of boatmen										
<b>Component 4: Sustainable resource development</b>											
<b>4.1 Sustainable fisheries Development</b>											
4.1.1	Bheri desiltation										
	Meeting of CAG to prepare a list of bheri needing desiltation										
	Desiltation estimate preparation										
	Bheri Desiltation										
	Transportation of silt to embankments or agricultural farms										
4.1.2	Establishment of new hatchery and fish feed marketing unit										
	Formation of SHG										
	Training workshops										
	Fish hatchery establishment and fish feed marketing units										
	Operation and maintenance										
4.1.3	Bheri water quality and fish monitoring										
	Training on water quality monitoring and fish disease management										
	Provision of water quality kits										
	App for data collection and management										
<b>4.2 Sustainable agriculture and horticulture development</b>											
4.2.1	Promoting rain water harvesting										
	Training										
	Unit establishment										
	Handholding support										
4.2.2	Diversification of cropping pattern										
	Training on cultivation of crops										
	Demonstration plots										
	Handholding support										
<b>4.3 Community infrastructure</b>											
4.3.1	Health camps										
4.3.2	Safe drinking water, sanitation and nutrition management										

## 8.4 Financing Arrangements

The management plan may be financed under the National Mission of Clean Ganga (NMCG) under the relevant scheme guidelines. Additional sources of funds may be the National Plan for Conservation of Aquatic Ecosystems of the MoEFCC, which requires that the state government provides 40% of matching funds.

## References

---

- Alam, M, Alam, M.M, Curray, J.R., Chowdhury, M.L.R and Gani, M.R. (2003): 'An overview of the sedimentary geology of the Bengal Basin in the regional tectonic framework and basin fill history', *Sedimentary Geology*, 155, 179-208.
- Allison, M.A., Khan, S.R., Goodbred Jr., S.L., Kuehl, S.A. (2003). Stratigraphic evolution of the late Holocene Ganges-Brahmaputra lower delta plain. *Sedimentary Geology*, 155, 317–342.
- Bagchi, K. (1944): *The Ganges Delta*, University of Calcutta, Calcutta.
- Bandyopadhyay, S. (2007). Evolution of the Ganga Brahmaputra delta: A review. *Geographical Review of India*, 69(3), 235–268.
- Banerjee M, Sen PK, Dastidar AG 1984. "On the depositional condition of the Holocene sediments of Bengal basin with remarks on environmental condition of the soft grey clay deposition in Calcutta". *Indian Geotech Conf* 84(1):63–69
- Banerjee, Sarmila and Dey, Debanjana 2017. "Eco-system Complementarities and Urban Encroachment: A SWOT Analysis of the East Kolkata Wetlands, India," *Cities and the Environment (CATE)*: Vol. 10: Iss. 1, Article 2.
- Barui NC, Chanda S 1992. "Late Quaternary pollen analysis in relation to palaeoecology, biostratigraphy and dating of Calcutta peat". *Indian Natl Sci Acad* 58B (4):191–200
- Bhadra, T, Mukhopadhyay A, and Hazra Sugata. 2017. "Identification of river discontinuity using geo-informatics to improve freshwater flow & ecosystem services in Indian Sundarban delta". *Springer Remote Sensing/Photogrammetry*, 137–152. doi:10.1007/978-3-319-46010-9\_10
- Biswas B 1969. "Unpublished records of Zoological Survey of India". In Ghosh A.K. 1990. *Biological Resources of wetlands of East Calcutta*. *Indian Journal of Landscape System and Ecological Studies*. 13(1): 10-23.
- Biswas KP 1927. "Flora of the Salt Lakes, Calcutta", *Journal of Department of Science, University of Calcutta*, 8.
- Brown, S. And Nicholls, R.J. (2015): Subsidence and human influences in mega deltas: The case of the Ganges-Brahmaputra-Meghna, *Science of the Total Environment* 527-528, 362-374.
- Bunting, S W, Kundu, N, and Mukherjee, M. 2002. "Situation Analysis: Production Systems and Natural Resource Management in PU Kolkata." Working Paper. Stirling, UK: Institute of Aquaculture.
- Bunting, Stuart W., Jules Pretty, and Peter Edwards. 2010. "Wastewater-Fed Aquaculture in the East Kolkata Wetlands, India: Anachronism or Archetype for Resilient Ecocultures? Wastewater-Fed Aquaculture in the EKW." *Reviews in Aquaculture* 2 (3): 138–53. <https://doi.org/10.1111/j.1753-5131.2010.01031.x>.
- CGWB (Central Ground Water Board), 2004. "Hydrogeology and ground water development of North-24 Parganas District, West Bengal". Kolkata: Government of India, CGWB Eastern Region, Technical report, series D.
- Chakraborty G., and Das Gupta D 2019. "From conflict to co-production a multi stakeholder analysis in preserving East Kolkata Wetlands". *Forum for policy dialogue on water conflicts in India*
- Chakraborty, Gorky, and Dhruba Das Gupta. 2019. "From Conflict to Co-Production A Multi-Stakeholder Analysis in Preserving the East Kolkata Wetlands." Pune: Forum for Policy
- Chandra, K., Raghunathan, C. and Mao, A.A. 2020. *Biodiversity Profile of East Kolkata Wetlands*, 1- 326 Jointly published by the Director, Zool. Surv. India, Kolkata and East Kolkata Wetlands Management Authority, Department of Environment, Govt. of West Bengal
- Chatterji GC, Deshmukh DS, Sinha SC, Niyogi BN 1959. "Some recent observations on the depth of sediment in the Ganga valley below Varanasi. *Ganga Valley: a symposium*". ONGC, Dehradun
- CPCB. 2019. "Summary of the Detailed Study Taken by East Kolkata Wetlands Management Authority, Institute of Environmental Studies & Wetland Management and Central Pollution Control Board, Eastern Regional Directorate, Kolkata." Kolkata: Central Pollution Control Board.

- Curry, J.R. (1994): 'Sediment volume and mass beneath the Bay of Bengal'. *Earth and Planetary Science Letters*, 125, 371-383.
- Curry, J.R., Emmel, F.J, Moore, D.G.(2003). The Bengal Fan: morphology, geometry, stratigraphy, history and process; *Marine and Petroleum Geology* 19, 1191-1223.
- Das A., and Mandal A., 2018. "Waste systems: Its utility and analyses in aquaculture". In: Jana B., Mandal R., Jayasankar P. (eds) *Wastewater Management Through Aquaculture*. Springer, Singapore
- Dasgupta, R. 1973. "Contribution of botany of a portion of Salt Lakes, West Bengal", *Ind.Mus. Bull.*, vol. 1.
- Dasgupta, S., Gosain, A.K., Rao, S., Roy, S., Sarraf, M., 2013. "A megacity in a changing climate: the case of Kolkata". *Climatic Change* 116, 747–766. <https://doi.org/10.1007/s10584-012-0516-3>
- Dasgupta, S. and Panigrahi, A., 2014. Studies on The Effect Of Aquatic Pollution On Ichthyofaunal Diversity Of The East Kolkata Wetlands. *International Journal of Research in Applied, Natural and Social Sciences*, 2(4), pp.145-152.
- De, M., Bhunia, S. and Sengupta, T. 1989. "A Preliminary Account on Major Wetland Fauna of Calcutta and Surroundings". *Ecology* 3 (9): 5-11
- Dey, Debanjana, and Sarmila Banerjee. 2018. "How Expensive Is the Decay of East Kolkata Wetlands? An Estimation of Opportunity Cost for Kolkata." In *Sustainable Urbanization in India*, edited by Jenia Mukherjee, 181–205. Exploring Urban Change in South Asia. Singapore: Springer Singapore. [https://doi.org/10.1007/978-981-10-4932-3\\_10](https://doi.org/10.1007/978-981-10-4932-3_10).
- Dutta, Joystu, Arpita Saha, and Abhijit Mitra. 2016. "Impact of Acidification on Heavy Metal Levels in a Bheri of East Kolkata Wetlands (EKW), a Ramsar Site in the Indian Sub-Continent." *International Journal of Development Research* 7 (7): 15452–58.
- EKWMA. 2016. "Birds of East Kolkata Wetlands." East Kolkata Wetlands Management Authority.
- EKWMA, 2020. Pictorial Guide Amphibians, Reptiles and Mammals of East Kolkata Wetlands. East Kolkata Wetlands Management Authority and Zoological Survey of India.
- Finlayson, C Max. 2012. "Forty Years of Wetland Conservation and Wise Use: EDITORIAL." *Aquatic Conservation: Marine and Freshwater Ecosystems* 22 (2): 139–43. <https://doi.org/10.1002/aqc.2233>.
- Finlayson, C Max, Nick Davidson, Dave Pritchard, G Randy Milton, and Heather Mackay. 2011. "The Ramsar Convention and Ecosystem-Based Approaches to the Wise Use and Sustainable Development of Wetlands." *Journal of International Wildlife Law & Policy* 14 (3–4): 176–98.
- Ghosh, D. and S. Sen, 1987. "Ecological History of Calcutta's Wetland Conversion". *Environ. Conserv.*, 14: 219-226. DOI: 10.1017/s0376892900016416
- Ghosh, D., 2005. "Ecology and Traditional Wetland Practice: Lessons from Wastewater Utilisation in the East Calcutta Wetlands". 1st Edn., *Worldview*, Kolkata, pp: 120.
- Ghosh, Dhrubajyoti. 1999. "Wastewater Utilisation in East Calcutta Wetlands." *Occasional Paper*. Nieuwehaven, The Netherlands: IRC. 2005. *Ecology and Traditional Wetland Practice: Lessons from Wastewater Utilisation in the East Calcutta Wetlands*. Kolkata: *Worldview*.
- Ghosh, S. K., and Ghosh D. 2003. "Community based rehabilitation of wetlands in West Bengal, India", S. B. Ray et al. (ed.), *Contemporary Studies in Natural Resource Management in India*, Forest Studies Series, Inter-India Publication, New Delhi.
- Ghosh, S., 2018. "Wastewater-Fed Aquaculture in East Kolkata Wetlands: State of the Art and Measures to Protect Biodiversity", in: Jana, B.B., Mandal, R.N., Jayasankar, P. (Eds.), *Wastewater Management Through Aquaculture*. Springer Singapore, Singapore, pp. 119–137. [https://doi.org/10.1007/978-981-10-7248-2\\_6](https://doi.org/10.1007/978-981-10-7248-2_6)
- Hait AK, Das HK, Ghosh S, Ray AK, Chanda S 1996. "Environmental variations in Late Quaternary sequence of Kolaghat, West Bengal, India". *Curr Sci* 70(12):1089–1093
- Hussan, Ajmal, J.K. Sundaray, R.N. Mandal, Farhana Hoque, Arabinda Das, P.P. Chakrabarti, and S. Adhikari. 2019. "Invasion of Non-Indigenous Suckermouth Armoured Catfish of the Genus *Pterygoplichthys* (Loricariidae) in the East Kolkata Wetlands: Stakeholders' Perception." *Indian Journal of Fisheries* 66 (2). <https://doi.org/10.21077/ijf.2019.66.2.86267-05>.
- Indiawris.gov.in. 2020. India-WRIS. [online] Available at: <<https://indiawris.gov.in/wris/#/Aquifer>> [Accessed 20 July 2020].



- Jana S. 2018. "Socioeconomic Impacts and Cost-Benefit Analysis of Wastewater-Fed Aquaculture". In: Jana B., Mandal R., Jayasankar P. (eds) *Wastewater Management Through Aquaculture*. Springer, Singapore
- Joystu, Dutta. 2017. "Bioaccumulation of Toxic Heavy Metals in the Edible Fishes of Eastern Kolkata Wetlands (EKW), the Designated Ramsar Site of West Bengal, India." *International Journal of Aquaculture and Fishery Sciences*, March, 018–021. <https://doi.org/10.17352/2455-8400.000023>.
- Khatun, A., Pal, S., Mukherjee, A., Samanta, P., Mondal, S., Kole, D., Chandra, P. and Ghosh, A., 2016. Evaluation of metal contamination and phytoremediation potential of aquatic macrophytes of East Kolkata Wetlands, India. *Environmental Health and Toxicology*, 31, p.e2016021.
- KMC Drainage Report, 2020. Report on S&D Department for Chief Secretary's Environmental Monitoring Cell. 1-3
- Kumar, Bhupander, Kurunthachalam Senthil Kumar, Mahalakshmi Priya, Debapriya Mukhopadhyay, and Rita Shah. 2010. "Distribution, Partitioning, Bioaccumulation of Trace Elements in Water, Sediment and Fish from Sewage Fed Fish Ponds in Eastern Kolkata, India." *Toxicological & Environmental Chemistry* 92 (2): 243–60. <https://doi.org/10.1080/02772240902942394>.
- Kumar, Ritesh. 2018. "East Kolkata Wetlands and the Regulation of Water Quality." In *The Wetland Book*, edited by C. Max Finlayson, Mark Everard, Kenneth Irvine, Robert J. McInnes, Beth A. Middleton, Anne A. van Dam, and Nick C. Davidson, 1293–99. Dordrecht: Springer Netherlands. [https://doi.org/10.1007/978-90-481-9659-3\\_222](https://doi.org/10.1007/978-90-481-9659-3_222).
- Kundu, N., & Chakraborty, A. 2017. "Dependence on Ecosystem Goods and Services: A Case Study on East Kolkata Wetlands, West Bengal, India". *Wetland Science*, 381–405. doi:10.1007/978-81-322-3715-0\_20
- Mahapatra, B K, and W S Lakra. 2014. "Ornamental Fishes of East Kolkata Wetland, West Bengal, India." *International Journal of Scientific Research* 3 (12): 406–8.
- Mallick, Jayanta Kumar. 2009. "View of Endemic Marsh Mongoose *Herpestes Palustris* (Carnivora\_ Herpestidae) of East Kolkata Wetlands, India: A Status Report." *Journal of Threatened Taxa* 1 (4).
- Mondal, Biswajit, Gour Dolui, Malay Pramanik, Santu Maity, Sumantra Sarathi Biswas, and Raghunath Pal. 2017. "Urban Expansion and Wetland Shrinkage Estimation Using a GIS-Based Model in the East Kolkata Wetland, India." *Ecological Indicators* 83 (December): 62–73. <https://doi.org/10.1016/j.ecolind.2017.07.037>.
- Mukherjee, A. and Bardhan, S., 2019. "An Assessment of Direct and Indirect Emission Reduction Potential of Natural Wetland Systems of Kolkata, India". *International Journal of Applied Engineering Research*, 14(8), p.1923.
- Mukherjee, J., & Ghosh, A. 2015. "Water justice city profile: Kolkata, India". [https://www.bartlett.ucl.ac.uk/dpu/water-justice/outputs/Kolkata\\_report.pdf](https://www.bartlett.ucl.ac.uk/dpu/water-justice/outputs/Kolkata_report.pdf). Mukherjee, Jenia. 2020. "Untamed Practices." In *Blue Infrastructures*, by Jenia Mukherjee, 85–123. *Exploring Urban Change in South Asia*. Singapore: Springer Singapore. [https://doi.org/10.1007/978-981-15-3951-0\\_4](https://doi.org/10.1007/978-981-15-3951-0_4).
- Mukherjee, V., Das, A., Akhand, A., & Gupta, G. 2013. "Toxicity and profitability of rice cultivation under wastewater irrigation: the case of the East Calcutta Wetlands". *Ecological Economics*, 93, 292–300. doi: 10.1016/j.ecolecon.2013.06.010
- Nandi, S, C Srivastava, and K M Agarwal. 2013. "Heavy Metal Concentration in Surface Water of East Kolkata Wetland." *Indian Journal of Public Health Research & Development* 4 (4): 227–30.
- Nath, Susanta, and Madhumita Bhoumik. 2013. "Levels of Toxic Metals in Edible Fish from a Wetland of India." *International Journal of Environmental Sciences* 3 (5): 1509–15.
- Niyogi, S. and Ray, S., 2017. "Attempt to Grab EKW Land Foiled" | *Kolkata News - Times Of India*.
- Oldham.(1870): President's address. *Proceedings, Asiatic Society of Bengal for February, 1870, Calcutta*.
- Parihar, Seema Mehra, Soma Sarkar, Amitava Dutta, Shilpi Sharma, and Tanushree Dutta. 2013. "Characterizing Wetland Dynamics: A Post-Classification Change Detection Analysis of the East Kolkata Wetlands Using Open Source Satellite Data." *Geocarto International* 28 (3): 273–87. <https://doi.org/10.1080/10106049.2012.705337>.

- Raychaudhuri, S., Mishra, M., Nandy, P., & Thakur, A. R. 2008. "Waste management: A case study of ongoing traditional practices at east Calcutta Wetland". *American Journal of Agricultural and Biological Science*, 3(1), 315–320.  
<https://doi.org/10.3844/ajabssp.2008.315.320>
- Rennell, J. (1780): *A Bengal Atlas*, London. (edited by Kalyan Rudra (2016) and published by Sahitya Samsad, Kolkata.
- Rudra, K. (2008): *Banglar Nadikatha* (in Bengali). Sahitya Samsad, Kolkata.
- Rudra, K. (2012): *Atlas of Changing River Courses in West Bengal*, Sea Explorers' Institute, Kolkata.
- Rudra, K. (1999): 'The hypothesis of easterly flight of the Ganga water: Fact or Fiction', *Indian Journal of Geography and Environment*, 4, Deptt. of Geography, Vidyasagar University, 52-55
- Rudra, K. (2014): Changing river courses in the western part of the Ganga–Brahmaputra delta. *Geomorphology*, 227, 87–100.
- Rudra, K. (2015): Ref: Changing river courses in the western part of the Ganga–Brahmaputra, *Geomorphology* 250 (2015) 454–458.
- Rudra, K. (2018): *Rivers of the Ganga-Brahmaputra-Meghna delta / A Fluvial Account of Bengal*. Springer.
- Sahu, P, and P K Sikdar. 2011. "Threat of Land Subsidence in and around Kolkata City and East Kolkata Wetlands, West Bengal, India." *Journal of Earth System Science* 120 (3): 435–46. <https://doi.org/10.1007/s12040-011-0077-2>.
- Sahu, P., Sikdar, P.K., 2008. "Hydrochemical framework of the aquifer in and around East Kolkata Wetlands, West Bengal, India". *Environ Geol* 55, 823–835.  
<https://doi.org/10.1007/s00254-007-1034-x>
- Sarkar, Subhasis, Phani Bhusan Ghosh, Alok Kumar Sil, and Tapan Saha. 2011. "Heavy Metal Pollution Assessment through Comparison of Different Indices in Sewage-Fed Fishery Pond Sediments at East Kolkata Wetland, India." *Environmental Earth Sciences* 63 (5): 915–24. <https://doi.org/10.1007/s12665-010-0760-7>.
- Sen PK, Banerjee M 1990. "Palynoplankton stratigraphy and environmental changes during Holocene in Bengal basin, India". *Rec Palaeobot Palynol* 62:25–35
- Sikdar PK 2000. "Geology of the Quaternary Aquifers of the Twin City of Calcutta–Howrah". *J Geol Soc India* 56:169–181
- Sikdar PK, Mondal S, Saha L, Sarkar SS, Banerjee S 2002. "Environmental impact assessment of a proposed info-tech complex in East Calcutta wetlands". *Environmentalist* 22(4): 241–260
- Subhayan Dutta and Swati Chakraborty 2017 *The Effect of Pollution on Hydrological Parameters Analysis of East Kolkata Wetland Area*. *International Journal of Development Research*. Vol. 07, Issue, 09, pp.15452-1545
- Stackhouse, P., 2020. NASA POWER | Data Access Viewer. [online] Power.larc.nasa.gov. Available at: <<https://power.larc.nasa.gov/data-access-viewer/>> [Accessed 15 July 2020].
- Vicziany, M., Chattopadhyay, D., Bhattacharyya, S., 2017. "Food from Sewage: Fish from the East Kolkata Wetlands and the Limits of Traditional Knowledge". *South Asia: Journal of South Asian Studies* 40, 619–644. <https://doi.org/10.1080/00856401.2017.1341038>
- Vishwakarma, A., Kunte, A., Mishra, D., Panchal, S., and Kundu S., 2014. "Dhapa Dumpsite Environmental and Social Assessment Report". West Bengal Pollution Control Board <http://documents1.worldbank.org/curated/en/383421497427377147/pdf/SFG3430-VI-EA-P091031-Box402914B-PUBLIC-Disclosed-6-14-2017.pdf>.
- Vivekananda Mukherjee and Gautam Gupta 2011. *Toxicity and Profitability of Rice Cultivation under Waste-Water Irrigation: The Case of the East Calcutta Wetlands*. SANDEE Working Papers, ISSN 1893-1891; WP 62–11
- WBPCB, 2019-20. *Analysis Report of Kolkata Sewage Canal from March, 2019 to February, 2020*. West Bengal Pollution Control Board.

## Annexes

---

### AI Brief Document (as per Wetlands (Conservation and Management) Rules, 2017)

Brief Document for the East Kolkata Wetlands (Ramsar Site No. 1208)

State / Union Territory: West Bengal

Name and address of person(s) compiling this information: East Kolkata Wetlands Management Authority, Pranisampad Bhavan, 5th floor, LB-2, Sector III, Salt Lake, Kolkata- 700 106

#### **Section I: Identification, Location and Jurisdiction**

I.1 Name of the Wetland (Alternative names, including in local language should be given in parenthesis after official name)

#### **East Calcutta Wetlands/ East Kolkata Wetlands**

I.2 Name of the Village(s) , Tehsil(s), Municipal area (s)

**List of mouzas falling within the East Kolkata Wetlands is enclosed as Annex A 3**

I.3 Name of the District(s) in which wetland complex is located:

#### **South 24 Parganas and North 24 Parganas**

I.4 Geographical coordinates (Latitude and Longitude, to degree, minutes and second)

Latitude: From 22° 25' N to 22° 35' N

Longitude: From 88° 20' E to 88° 35' E

I.5 Name of the Department / Agency which has jurisdiction over the wetland / wetlands complex

**East Kolkata Wetlands Management Authority, Department of Environment, Government of West Bengal**

## Section 2: Site Characteristics

2.1 Area of wetland / wetlands category (ha): **12500**

2.2 Wetland type (Please tick appropriate categories and sub-categories)

Category	Subcategory
<input checked="" type="checkbox"/> Natural (Inland)	<input type="checkbox"/> Permanent lakes <input type="checkbox"/> Seasonal/ intermittent lakes <input type="checkbox"/> Permanent streams/ creeks <input type="checkbox"/> Seasonal/ intermittent streams/ creeks <input type="checkbox"/> Oxbow <input type="checkbox"/> River floodplain <input checked="" type="checkbox"/> Permanent freshwater marshes <input type="checkbox"/> Seasonal/ intermittent freshwater marshes <input type="checkbox"/> Shrub-dominated wetlands <input type="checkbox"/> Tree-dominated wetlands <input type="checkbox"/> Geothermal wetlands <input type="checkbox"/> Karst and other subterranean hydrological systems
<input type="checkbox"/> Natural (Coastal)	<input type="checkbox"/> Coastal lagoon <input type="checkbox"/> Estuary <input type="checkbox"/> Intertidal mud, sand or salt flats <input type="checkbox"/> Mangroves <input type="checkbox"/> Coral reefs
<input checked="" type="checkbox"/> Human-made	<input checked="" type="checkbox"/> Aquaculture Pond <input type="checkbox"/> Tank <input type="checkbox"/> Saltpan <input type="checkbox"/> Dam / Reservoir

2.3 Depth (m) : Average \_\_\_\_\_ | \_\_\_\_\_ Maximum \_\_\_\_3\_\_\_\_\_

2.4 Elevation (m above mean sea level) 2 -6.5 amsl

2.5 Water regimes

a) Main source of water (tick all applicable)

Rainfall

Groundwater

Catchment runoff

- Direct / indirect inflow from surface water  
 Others, please specify **sewage water**
- b) Water permanence  
 Mostly permanent     Mostly intermittent
- c) Destination of water from wetland  
 Feeds groundwater     To downstream catchment     To river     To sea
- d) Water pH  
 Acid (< 5.5)     Circumneutral (5.5 – 7.4)     Alkaline (> 7.4)     Not known
- e) Water salinity  
 Fresh (< 0.5 g/l)     Brackish (0.5 – 30 g/l)     Euhaline (30- 40 g/l)     Hypersaline (>40g/l)  
 Not known
- f) Nutrient in water  
 Eutrophic     Mesotrophic     Oligotrophic     Not known

## 2.6 Climatic setting

- a) Annual Rainfall /Snowfall(mm): 1600 mm (rainfall)  
b) Temperature (°C): Minimum 10 Maximum 40  
c) Humidity (%): Minimum 70 Maximum 90

2.7 Area of zone of influence (in ha): 162500 (direct catchment area)

2.8 Major land use within zone of influence (provide as approximate % of catchment area)

Forests	_____ %
Plantation	_____ %
Agriculture	_____ 59_ %

2.9 Map of wetland complex and zone of influence

(To be enclosed as Annexure I and Annexure II to this proposal)

## Section 3: Biodiversity

3.1 Notable plant species present in wetland

\_\_\_\_\_ At least 93 species of plants have been identified in the wetlands \_\_\_\_\_

3.2 Notable animal species present in wetland

\_\_\_\_\_ As per the Indian Wildlife Protection Act, 1972, the East Kolkata Wetlands is a habitat of Schedule I species viz. *Varanus salvator*, *Varanus flavescens*, *Lissemys punctata* and Schedule II species viz. *Paradoxurus hermaphrodites*, *Canis aureus*, *Herpestes auro-punctatus*. Marsh mongoose (*Herpestes auro-punctatus*) is endemic to the region and is also included in the schedule II of Indian Wildlife Protection Act, 1972\_

3.3 Species of conservation significance (rare, endangered, threatened, endemic species)

\_\_\_\_\_ East Kolkata Wetlands (EKW) are home to endangered and vulnerable species such as *Clarias magur*, *Pangasianodon hypophthalmus*, *Cyprinus carpio*, *Oreochromis mossambicus* and *Wallago attu* (Near Threatened) to mention a few \_\_\_\_\_

3.4 Major plant invasive alien species

\_\_\_\_\_ Some of the major invasive species are *Ageratum conyzoides*, *Alternanthera paronychioides*, *Argemone Mexicana* and *Parthenium hysterophorus* \_\_\_\_\_

3.5 Major animal invasive alien species

\_\_\_\_\_ NA \_\_\_\_\_

**Section 4: Ecosystem services**

Importance	Relevant for the site (please tick yes or no)	If Yes, Details (upto 50 words for each category)
Source of drinking water for people living and around	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	
Source of water for agriculture	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	EKW have an intricate network of canals through which sewage water is let into the bhery. This sewage water is extensively being used for irrigation
Fisheries	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	EKW comprises a large number of sewage-fed shallow pisciculture ponds fishponds
Cultivation of aquatic food plants	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Significant aquatic food plants are rice ( <i>Oryza Sativa</i> ), Lalshak ( <i>Amaranthus gangeticus</i> ), Noteyshak ( <i>Amaranthus viridis</i> )

For buffalo wallowing and use of domesticated animals	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Insignificant
Medicinal plants	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Medicinal plants (tulsi, ghritokumari) have been introduced for crop diversification and enhancing returns to farmers
Is a recreational site	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	
Buffering communities from extreme events as floods and storms	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Storm Weather Flow (S.W.F.) Channel and Dry Weather Flow (D.W.F.) Channel carry storm weather discharges and dry weather discharges, pumped out from the different pumping stations through EKW since Kolkata is sloped towards east
Groundwater recharge	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	
Water purification	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	The sewage is fed to the fishponds on regulated basis. The sewerage goes to the wetland gets purified with the help of solar energy and creates food for the fishes by the growth of planktons

Acts as a sink for sediments	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Sewage water carries sediment load. The fishponds of EKW act as Waste Stabilization Ponds where sediments are trapped at the bottom of the pond floor
Has significant cultural and religious values	<input type="checkbox"/> Yes <input type="checkbox"/> No	Not assayed
Is a site for recreation and tourism	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	A variety of migratory birds visit the wetlands. Water body centric recreation sites are being developed here. Nalban is a good example.
Supports noteworthy plants species	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Please refer to Annex A4.
Supports noteworthy animal species	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Please refer to Annex A4.
Site of high congregation of migratory water birds	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Please refer to Annex A4.
Supports life cycle of fish or amphibians	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Please refer to Annex A4.
Mining	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	
Any other, please list		



## Section 5: Pre-Existing Rights and Privileges

Nature of right and privilege	Relevant for the site (please tick yes or no)	Does this negatively impact the wetland's ecological health?	Brief description (upto 50 words for each category)
Community Fishing (without any lease or permission from government department)	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> Not assessed	A number of informal fishermen's co-operative operate bhery
Fishing under lease from government department	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> Not assessed	Fisheries management through bhery are under three forms of ownership: a) government; b) cooperatives, and c) private. Bhery ownership by cooperatives has gradually reduced (from 27% to 6% during 2002-15)
Harvest of plants (without any lease or permission from government department)	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not assessed	
Harvest of plants under lease from government department	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not assessed	
Agriculture or horticulture within wetland	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> Not assessed	Vegetable production from East Kolkata Wetlands meets nearly 20% of the vegetable requirement of Kolkata city. Proximity to the city, availability of water makes horticulture a household activity in East Kolkata Wetlands, farmers even rent small plots for raising vegetables.

Grazing	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not assessed	
Religious practices	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not assessed	
Withdrawal of water for domestic use	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not assessed	For domestic purposes water from fishponds is used.
Withdrawal of water for agriculture or fisheries	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> Not assessed	Rice is cultivated using effluent water from the bhery, covering an area of 2637 ha. Rice cultivation is one of the major livelihoods within the wetland followed by cultivation of vegetables.
Bathing or wallowing of domestic animals	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> Not assessed	Insignificant
Plying of boats	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> Not assessed	Non-motorised boats are used mainly for recreation and fishing purposes.
Any other, please list here	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not assessed	

## Section 6: Present and Potential Threats

Threat	Degree	Present or Potential	Additional information, if any
Changes in water inflow and outflow	<input type="checkbox"/> High <input type="checkbox"/> Medium <input type="checkbox"/> Low	<input type="checkbox"/> Present <input type="checkbox"/> Potential	
Pollution	<input type="checkbox"/> High <input checked="" type="checkbox"/> Medium <input type="checkbox"/> Low	<input checked="" type="checkbox"/> Present <input type="checkbox"/> Potential	Area under solid waste has doubled. Threats due to pollution and discharge of leachate remain challenges needing to be addressed.
Unsustainable harvest of biological resources	<input type="checkbox"/> High <input type="checkbox"/> Medium <input type="checkbox"/> Low	<input type="checkbox"/> Present <input type="checkbox"/> Potential	
Mining	<input type="checkbox"/> High <input type="checkbox"/> Medium <input type="checkbox"/> Low	<input type="checkbox"/> Present <input type="checkbox"/> Potential	
Siltation	<input type="checkbox"/> High <input checked="" type="checkbox"/> Medium <input type="checkbox"/> Low	<input checked="" type="checkbox"/> Present <input type="checkbox"/> Potential	Flow regulation has led to accumulation of sediments and siltation in the wetland catchment. The fishponds of EKW also act as Waste Stabilization Ponds where sediments coming from sewage water are trapped at the bottom of the pond
Encroachment	<input checked="" type="checkbox"/> High <input type="checkbox"/> Medium <input type="checkbox"/> Low	<input checked="" type="checkbox"/> Present <input type="checkbox"/> Potential	The EKW, being adjacent to the metropolis, is subjected to considerable

			pressure for real estate projects. Expansion of settlements inside the wetland boundary poses a risk of reducing the inundated area
Spread of invasive species	<input type="checkbox"/> High <input type="checkbox"/> Medium <input checked="" type="checkbox"/> Low	<input checked="" type="checkbox"/> Present <input type="checkbox"/> Potential	
Any other, please list	<input type="checkbox"/> High <input type="checkbox"/> Medium <input type="checkbox"/> Low	<input type="checkbox"/> Present <input type="checkbox"/> Potential	

**Section 7: Activities Proposed to be prohibited (other than those listed in Rule 4(2) of Wetlands Rules)**

Activity	Prohibited within wetlands or zone of influence	Details of specific area wherein activity is prohibited	Name of department / agency responsible for regulation	Additional information, if any
	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence			

**Section 8: Activities Proposed to be regulated**

Activity	Place a tick mark if relevant	Regulation within wetlands or zone of influence	Level of regulation (in terms of people, restricted area or any other)	Name of department / agency responsible for regulation	Additional information, if any
Withdrawal of water / impoundment/diversion or any other hydrological intervention	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	Water allocation for human and ecological purposes through assessing and implementing environmental flows as a basis for operation of all the hydraulic structures in the upstream and downstream reaches.		

Harvesting of resources (living / non-living)	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence			
Grazing	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence			
Discharge of treated sewage/ effluent / wastewater	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	Regulating discharge of industrial effluents as per standards		
Construction of boat jetties, and facilities for temporary use, as pontoon bridges	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence			
Aquaculture, agriculture and horticulture activities within the wetland boundaries.	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	For the ease of management, and also ensuring compliance with the extent regulatory regimes, it is proposed to segment the wetland into four major zones- fishery zone, agriculture zone, horticulture zone, nature protection zone		

Trekking and Tourism	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence			
----------------------	--------------------------	--	--	--	--

**Section 9: Activities Proposed to be permitted**

Activity	Place a tick mark if relevant	Within wetlands or zone of influence	Additional information, if any
Ecotourism	<b>X</b>	<b>X</b> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	Construction and operationalization of an interpretation centre with facilities. Specific training programmes for various target groups are proposed to be undertaken as an integral part of the activities. Signages, communication and transport facilities be developed for complete tourist education and recreation experience.
	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	
	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	

	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	
	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	
	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	
	<input type="checkbox"/>	<input type="checkbox"/> Wetland / Wetlands complex boundary <input type="checkbox"/> Zone of influence	

### Section 10: Listing of Available Scientific Resources Used

East Kolkata Wetlands Newsletter (2010), Volume I

Integrated Management Plan of East Kolkata Wetlands (Management Action Plan 2021-2026)


### CHECKLIST

- Responsible agency has been clearly identified and details of contact person included
- Wetland/ wetlands complex boundary has been delineated using GIS and firmed up by adequate ground truthing
- Wetland/ wetlands complex map has been provided at required scale
- Zone of influence has been delineated and included in wetland map or a separate map
- Wetland zone of influence is sufficient to manage all activities
- Site's importance have been listed, and for major categories, justification is provided
- Site's biodiversity values are listed, and for major categories, justification is provided


- List of pre-existing rights and privileges is provided
- Consistency or inconsistency of pre-existing rights and privileges is indicated to be best of available knowledge
- Threats to site are listed, and for major categories details are provided
- Activities prohibited, beyond those already listed in Rule 4(2) have been mentioned
- List of activities to be regulated within wetlands and zone of influence is provided
- List of activities to be permitted is provided



## Annexure I- Map of wetland complex



## Annexure II Zone of influence of East Kolkata Wetlands



## A2 Wetland Health Card

Features	Indicator	Desired Value							Comments
			A	B	C	D	E	Score	
Area	% Wetland converted to non-wetland use since 2000	0%	0%	1-5%	6-10%	11-20%	More than 20%	B	
Hydrology and Catchment	Ratio of number of natural inlets choked and diverted to total number of natural inlets	<0.2	0-0.2	0.3-0.4	0.4-0.6	0.7-0.8	More than 0.8	A	
	Ratio of number of natural outlets choked and diverted to total number of natural outlets	<0.2	0-0.2	0.3-0.4	0.4-0.6	0.7-0.8	More than 0.8	A	
	Biological Oxygen Demand	Between 3-6 mg/l	80-100% sample meet the criteria	60-80% sample meet the criteria	40-60% sample meet the criteria	20-40% sample meet the criteria	Less than 20% sample meet the criteria	D	

Biodiversity	% Wetland area covered by invasive macrophytes.	<10%	<10%	11-20%	21-30%	31-40%	More than 40%	B	
	Annual waterbird count as a proportion of average count of last 5 years	0.7	More than 0.7	0.6-0.7	0.5-0.6	0.4-0.5	less than 0.5		
Governance	Clearly demarcated wetlands map	Wetland map prepared and approved by SWA	Wetlands map prepared and approved by SWA	Wetlands map prepared and under consideration of SWA	Wetlands map prepared but not placed under SWA	Wetland map under preparation	Wetlands map not prepared	A	
	Wetland Management plan	Management plan prepared and approved by State Wetlands Authority	Management plan prepared and approved by SWA	Management plan prepared and submitted to SWA	Management plan prepared, not submitted to SWA	Management plan under preparation	No management plan	C	
	Wetland's notification	Wetland notified under extent regulation	Final notification under extent regulation	Draft notification	Regulation under process	Regulation planned, process initiated	No regulation	A	

	Frequency	Assigned weight	
Number of indicators in Rank A	4	1	4
Number of indicators in Rank B	2	0.8	1.6
Number of indicators in Rank C	1	0.6	0.6
Number of indicators in Rank D	1	0.4	0.4
Number of indicators in Rank E	0	0.2	0
Number of indicators not calculated/ not known	1	0	0
		Total	6.6
Wetland Score			0.83
Grade			<b>B-</b>

### A3 List of Mouzas

<b>District</b>	<b>Panchayet/</b>	<b>Police</b>	<b>Mouza</b>	<b>J.L.</b>	<b>Status</b>	
24- Parganas (South)	Kolkata Municipal Corporation	Pragati Maidan/ Anandapur	Dhapa	2	Part mouza	
			Chowbaga	3	Full	
		Pragati Maidan	Bonchtala	4	Part mouza	
			Dhalenda	8	Full	
		Pragati Maidan/ Anandapur	Paschim Chowbaga	9	Full mouza	
	Anandapur	Nonadanga	10	Part mouza		
	Kheyadaha - 2 Gram Panchayat	Narendrapur	Chak Kolar Khal	1	Full	
			Karimpur	2	Full	
			Jagatipota	3	Full	
			Mukundapur	4	Full	
			Atghara	5	Full	
			Ranabhutia	6	Full	
			Kantipota	7	Full	
			Bhagabanpur	8	Full	
			Kharki	9	Full	
			Deara	10	Full	
			Kheyadaha - 1 Gram Panchayat	Kheadaha	11	Full
				Khodahati	12	Full
				Goalpota	13	Full
				Kumapukuria	14	Full
				Tardaha	15	Full
				Tihuria	16	Full
				Nayabad	17	Full
	Pratapnagar Gram Panchayat	Sonarpur	Samukpota	91	Full	
			Pratapnagar	92	Full	
			Garal	93	Full	
	Banamghata Gram Panchayat	Kolkata Leather Complex	Dakshin Dhapa	1	Full	
			Kochpukur (Erstwhile Dhapa	2	Part mouza	
			Hatgachha	4	Full	
			Hadia	5	Full	
Beonta - 2 Gram Panchayat			Dharmatala	6	Full	
			Kulberia	7	Full	
Beonta - 1 Gram			Beonta	27	Full	
Tardaha Gram			Tardaha Kapashati	38	Full	
Kolkata Municipal Corporation	Purba Jadabpur	Kalikapur	20	Part mouza		
24- Parganas (North)	Bidhannagar Municipal Corporation	South Bidhannagar	Dhapa Manpur	1	Part mouza	
		Electronics Complex	Thakdari	19	Part mouza	

## A4 List of Species

Data Source – Chandra, Raghunathan, and Mao 2020. Biodiversity Profile of East Kolkata Wetlands. Jointly published by the Director, Zool. Surv. India, Kolkata and East Kolkata Wetlands Management Authority, Department of Environment, Govt. of West Bengal

### I. List of Protozoa: Free-living Ciliates

Taxa
Phylum CILIOPHORA Class GYMNOSTOMATEA Order SPATHIDIIDA
Family LACRYMARIIDAE 1. Lacrymaria olor (Müller, 1786) Bory, 1824 Order HAPTORIDA
Family DIDINIIDAE 2. Didinium nasutum (Müller, 1773) Stein, 1859 Order PLEUROSTOMATIDA
Family LITONOTIDAE 3. Litonotus lamella (Müller, 1773) Schewiakoff, 1886 4. Litonotus fasciola (Ehrenberg, 1838) Wrzesniowski, 1870 Class CYRTOPHORIA Order CYRTOPHORIDA
Family CHILODONELLIDAE 5. Chilodonella uncinata (Ehrenberg, 1838) Strand, 1928 6. Trithigmostoma steini (Blochmann, 1895) Foissner, 1988 (Synonym- Chilodontopsis bengalensis Ghosh, 1921) Class PROTOSTOMATEA Order PROSTOMATIDA
Family COLEPIDAE 7. Coleps hirtus (Müller, 1786) Nitzsch, 1827 Class LITOSTOMATEA Order DILEPTIDA
Family DILEPTIDAE 8. Dileptus anatinus Golińska, 1971 Class HYPOTRICHEA Order EUPLOTIDA Family EUPLOTIDAE 9. Aspidisca lynceus Müller, 1773 Family GASTROCIRRHIDAE 10. Euplotoides woodruffi Gaw, 1939 Order STICHOTRICHIDA
Family HYPOTRICHIDIIDAE 11. Hypotrichidium conicum Ilowaisky, 1921 Order UROSTYLIDA
Family UROSTYLIDAE 12. Urostyla grandis Ehrenberg, 1830 Order OXYTRICHIDA

<p>Family OXYTRICHIDAE  13. <i>Oxytricha granulifera</i> Foissner &amp; Adam, 1983  14. <i>Stylonychia mytilus</i> (Müller, 1773) Ehrenberg, 1830  Class OLIGOTRICHEA  Order HALTERIIDA</p>
<p>Family HALTERIIDAE  15. <i>Halteria grandinella</i> (Müller, 1773) Dujardin, 1841  Class OLIGOHYMENOPHOREA  Order PENICULIDA</p>
<p>Family PARAMECIIDAE  16. <i>Paramecium caudatum</i> Ehrenberg, 1834  Order HYMENOSTOMATIDA</p>
<p>Family FRONTONIIDAE  17. <i>Frontonia</i> sp.  Order PERITRICHIDA</p>
<p>Family VORTICELLIDAE  18. <i>Vorticella</i> sp.  Class HETEROTRICHEA  Order HETEROTRICHIDA</p>
<p>Family STENTORIDAE  19. <i>Stentor coeruleus</i> (Pallas, 1766) Ehrenberg, 1830  Family SPIROSTOMIDAE  20. <i>Spirostomum teres</i> Claparède &amp; Lachmann, 1858  Class NASSOPHOREA  Order NASSULIDA</p>
<p>Family LEPTOPHARYNIDAE  21. <i>Leptopharynx costatus</i> Mermoud, 1914  Order SYNHYMENIIDA</p>
<p>Family ORTHODONELLIDAE  22. <i>Orthodonella gutta</i> (Cohn, 1866) Kahl, 1931 (Synonym- <i>Trachelius gutta</i> Sahrhage, 1915)</p>

## 2. List of Protozoa: Free-living Testate Amoebae

<b>Taxa</b>
<p>Phylum AMOEBOZOA Luhe 1913  Class TESTACEA LOBOSEA de Saedeller, 1934  Order ARCELLINIDA Kent, 1880</p>
<p>Family ARCELLIDE Ehrenberg 1832  1. <i>Arcella arenaria</i> Greef, 1866  2. <i>Arcella discoides</i> Ehrenberg, 1843  3. <i>Arcella vulgaris</i> Ehrenberg, 1830</p>
<p>Family MICROCORYCIIDAE DeSaedeleer, 1934  4. <i>Diploclamys leidy</i> Greeff, 1888</p>

Family MICROCHLAMYIIDAE Ogden, 1985

5. *Pyxidicula operculata* (Agardh, 1827) Ehrenberg, 1838

6. *Pyxidicula invisitata* Awerinzew, 1906

Family CENTROPYXIDAE Jung, 1942

7. *Centropyxis aculeata* Stein, 1857

8. *Centropyxis sylvatica* DeLandre, 1929

9. *Centropyxis cassis* (Wallich, 1864) DeLandre, 1929

10. *Centropyxis ecornis* (Ehrenberg, 1841) Leidy, 1879

11. *Centropyxis minuta* DeLandre, 1929

12. *Centropyxis spinosa* (Cash and Hopkinson, 1905) DeLandre, 1929

Family TRIGINOPYXIDAE Loeblich and Tappan, 1964

13. *Cyclopyxis arcelloides* (Penard, 1902) DeLandre, 1929

14. *Cyclopyxis eurystoma* (DeLandre, 1929) DeLandre, 1929

15. *Triginopyxis arcula* (Leidy, 1879) Penard, 1912

Family PLAGIOPYXIDAE Bonnet and Thomas, 1960

16. *Bullinularia indica* (Penard, 1907) DeLandre, 1953

17. *Plagiopyxis declivis* Bonnet and Thomas, 1955

Family DIFFLUGIIDAE Wallich, 1864

18. *Diffiugia acuminata* DeLandre, 1929

19. *Diffiugia accutissima* DeLandre, 1931

20. *Diffiugia corona* Wallich 1864

21. *Diffiugia curvicaulis* Penard, 1899

22. *Diffiugia elegans* Penard, 1890

23. *Diffiugia globulosa* (Dujardin, 1837) Penard, 1902

24. *Diffiugialobostoma* Leidy, 1879

25. *Diffiugia bryophila* (Penard, 1902) Jung, 1942

26. *Diffiugia oblonga* Ehrenberg, 1838

27. *Diffiugia urceolata* Carter, 1864

Family HELEOPERIDAE Jung, 1942

28. *Heleopera sylvatica* Penard, 1890

Family NEBELIDAE Taranek, 1882

29. *Quadrullella symmetrica* (Wallich, 1864) Schulze, 1875

Family LESQUEREUSIIDAE Jung, 1942

30. *Lesqueruesia modesta* Rhumbler, 1895

31. *Lesqueruesia spiralis* (Ehrenberg, 1840)

Phylum RHIZARIA Cavalier-Smith, 2002

Class TESTACEA FILOSEA de Saedeleer, 1934

Order EUGLYPHIDA Copeland, 1956

Family EUGLYPHIDAE Wallich, 1864

32. *Euglypha ciliata* Ehrenberg, 1848

33. *Euglypha compressa* Carter, 1890

34. *Euglypha laevis* Perty 1849

35. *Euglypha rotunda* Wailes and Penard 1911

36. *Euglypha strigosa* (Ehrenberg, 1848)

37. *Euglypha tuberculata* Dujardin, 1841

38. *Tracheleuglypha dentata* (Vejdowsky, 1832) DeLandre, 1928



Family Assulinidae Lara et al., 2007 39. <i>Placocista spinosa</i> (Carter, 1865) Leidy, 1879
Family Cyphoderiidae deSaedeleer, 1934 40. <i>Cyphoderia ampulla</i> (Ehrenberg, 1840) Leidy, 1878
Family TRINEMATIDAE Hoogenraad et de Groot, 1940 41. <i>Trinema enchelys</i> (Ehrenberg, 1838) Leidy, 1878 42. <i>Trinema lineare</i> Penard, 1890

### 3. List of Rotifers

Taxa
Phylum Rotifera Class Eurotatoria Subclass Monogononta Order Ploima
Family: Brachionidae 1. <i>Anuraeopsis fissa</i> Gosse, 1851 2. <i>Brachionus angularis</i> Gosse, 1851 3. <i>B. bidentatus</i> Anderson, 1889 4. <i>B. calyciflorus</i> Pallas, 1766 5. <i>B. caudatus</i> Barrois and Daday, 1894 6. <i>B. ahlstromi</i> Lindeman, 1939 7. <i>B. diversicornis</i> Daday, 1883 8. <i>B. falcatus</i> Zacharias, 1898 9. <i>B. forficula</i> Wierzejski, 1891 10. <i>B. plicatilis</i> Muller, 1786 11. <i>B. quadridentatus</i> Hermann, 1786 12. <i>B. rubens</i> Ehrenberg, 1838 13. <i>B. urceolaris</i> Muller, 1773 14. <i>Brachionus</i> sp. 15. <i>Keratella tropica</i> (Apstein, 1907) 16. <i>Keratella</i> sp. 17. <i>Platyonus patulus</i> (Muller, 1786)
Family: Lecanidae 19. <i>L. leontina</i> (Turner, 1892) 20. <i>L. papuana</i> (Murray, 1913) 21. <i>L. signifera</i> (Jennings, 1896) 22. <i>Lecane (Monostyla) decipiens</i> (Murray, 1913) 23. <i>L. (Monostyla) sp.</i>
Family: Lepadellidae 24. <i>Lepadella</i> sp.
Family: Trichocercidae 25. <i>Trichocerca (Diurella) similis</i> (Wierzejski, 1893) 26. <i>Trichocerca (Diurella) weberi</i> (Jennings, 1903)

<p>Family: Asplanchnidae  28. <i>A. intermedia</i> Hudson, 1886  29. <i>Asplanchna</i> sp.  Order: Flosculariacea</p>
<p>Family: Filinidae  30. <i>Filinia longiseta</i> (Ehrenberg, 1834)  31. <i>Filinia opoliensis</i> (Zacharias, 1898)  32. <i>Filinia</i> sp.</p>
<p>Family: Testudinellidae  33. <i>Testudinella patina</i> (Hermann, 1783)</p>
<p>Family: Scaridiidae  34. <i>Scaridium longicaudum</i> (Muller, 1786)  Subclass: Bdelloidea</p>
<p>Family: Philodinidae  35. <i>Rotaria</i> sp.  36. <i>Philodina citrina</i> Ehrenberg, 1830  37. <i>Philodina</i> sp.</p>

#### 4. List of Soil Nematodes

<b>Taxa</b>
<p>Phylum NEMATODA Rudolphi, 1808 (Lankester, 1877)  Order DORYLAIMIDA Pearse, 1942  Suborder DORYLAIMINA Pearse, 1936  Superfamily DORYLAIMOIDEA De Man, 1976  Family DORYLAIMIDAE De Man, 1976  Subfamily DORYLAIMINAE De Man, 1976  Genus DORYLAIMUS Dujardin, 1845  1. <i>Dorylaimus bengalensis</i> Sen, Chatterjee &amp; Manna, 2011</p>
<p>Subfamily LAIMYDORINAE Andr�ssy, 1969  Genus LAIMYDORUS Siddiqi, 1969  2. <i>Laimydorus baldus</i> Baqri &amp; Jana, 1982  3. <i>Laimydorus istvani</i> Sen, Chatterjee &amp; Manna, 2012  4. <i>Laimydorus siddiqii</i> Baqri &amp; Jana, 1982</p>
<p>Subfamily THORNENEMATINAE Siddiqi, 1969  Genus INDODORYLAIMUS Ali &amp; Prabha, 1974  5. <i>Indodorylaimus baqrii</i> Sen, Chatterjee &amp; Manna, 2012  6. <i>Indodorylaimus asaccatus</i> Sen, Chatterjee &amp; Manna, 2012</p>
<p>Genus SICAGUTTUR Siddiqi, 1971  7. <i>Sicaguttur sartum</i> Siddiqi, 1971</p>
<p>Genus COOMANSINEMOIDES Sen, Chatterjee &amp; Manna, 2012  8. <i>Coomansinemoides wasimi</i> Sen, Chatterjee &amp; Manna, 2012</p>
<p>Family APORCELAIMIDAE Heyns, 1965  Subfamily APORCELAIMINAE Heyns, 1965  Genus APORCELAIMELLUS Heyns, 1965  9. <i>Aporcelaimellus amylovorus</i> (Thorne &amp; Swanger, 1936) Heyns, 1965  10. <i>Aporcelaimellus baqrii</i> Ahmad &amp; Jairajpuri, 1982  11. <i>Aporcelaimellus chauhani</i> Baqri &amp; Khera, 1975  12. <i>Aporcelaimellus coomansi</i> Baqri &amp; Khera, 1975  13. <i>Aporcelaimellus indicus</i> Baqri &amp; Jairajpuri, 1968  14. <i>Aporcelaimellus papillatus</i> (Bastian, 1865) Baqri &amp; Khera, 1975</p>

15. <i>Aporcelaimellus tritici</i> (Bastian, 1865) Andrassy, 1986 Family QUDSIANEMATIDAE Jairajpuri, 1965 Subfamily DISCOLAIMINAE Siddiqi, 1969 Genus DISCOLAIMUS Cobb, 1913
16. <i>Discolaimus tenax</i> Siddiqi, 1964 Genus DISCOLAIMIUM Thorne, 1939
17. <i>Discolaimium mazhari</i> Baqri & Jairajpuri, 1968 Genus DISCOLAIMOIDES Heyns, 1963
18. <i>Discolaimoides bulbiferus</i> (Cobb, 1906) Heyns, 1963 Superfamily LONGIDOROIDEA Thorne, 1935 Family XIPHINEMATIDAE Dalmasso, 1969 Subfamily XIPHINEMATINAE Dalmasso, 1969 Genus XIPHINEMA Cobb, 1913
19. <i>Xiphinema americanum</i> Cobb, 1913 20. <i>Xiphinema inaequale</i> Khan & Ahmad, 1976 21. <i>Xiphinema manasiae</i> Sen, Chatterjee & Manna, 2010
Superfamily BELONDIROIDEA Thorne, 1939 Family BELONDIRIDAE Thorne, 1939 Subfamily BELONDIRINAE Thorne, 1939 Genus AXONCHIUM Cobb, 1920 Genus AXONCHIUM Cobb, 1920
22. <i>Axonchium (Axonchium) coomansi</i> Sen, Chatterjee & Manna, 2011
Superfamily TYLENCHOLAIMOIDEA Filipjev, 1934 Family MYDONOMIDAE Thorne, 1964 Subfamily MYDONOMINAE Thorne, 1964 Genus DORYLAIMOIDES Thorne & Swanger, 1936 Subgenus DORYLAIMOIDES Thorne & Swanger, 1936
23. <i>Dorylaimoides (Dorylaimoides) geraldii</i> Sen, Chatterjee & Manna, 2012 Subgenus Longidorylaimoides Jairajpuri and Ahmad, 1992
24. <i>Dorylaimoides (Longidorylaimoides) parvus</i> Thorne & Swanger, 1936 Family LEPTONCHIDAE Thorne, 1935 Subfamily TYLEPTINAE Thorne, 1935 Genus TYLEPTUS Thorne, 1939
25. <i>Tyleptus projectus</i> Thorne, 1939
Suborder NYGOLAIMINA Ahmad & Jairajpuri, 1979 Superfamily NYGOLAIMOIDEA Thorne, 1935 Family NYGOLAIMIDAE Thorne, 1935 Subfamily NYGOLAIMINAE Thorne, 1935 Genus AQUATIDES Heyns, 1968
26. <i>Aquatides heynsi</i> Sen, 2017 Genus LAEVIDES Heyns, 1968
27. <i>Laevides laevis</i> (Thorne, 1939) Thorne, 1974 Family NYGELLIDAE Andrassy, 1958 Subfamily NYGELLIBNAE Andrassy, 1958 Genus NYGELLUS Thorne, 1939
28. <i>Nygellus shamimi</i> Sen, 2015 Family AETHOLAIMIDAE Jairajpuri, 1965 Subfamily AETHOLAIMINAE Jairajpuri, 1965 Genus AETHOLAIMUS Williams, 1962
29. <i>Aetholaimus indicus</i> Jairajpuri, 1965 Order TYLENCHIDA Thorne, 1949 Suborder TYLENCHINA Chitwood in Chitwood and Chitwood, 1950 Infraorder TYLENCHATA Siddiqi, 2000 Superfamily TYLENCHOIDEA Örley, 1880 Family BELONOLAIMIDAE Whitehead, 1960 Subfamily TELOTYLENCHINAE Siddiqi, 1960

Genus TYLENCHORHYNCHUS Cobb, 1913 30. <i>Tylenchorhynchus mashhoodi</i> Siddiqi & Basir, 1959 31. <i>Tylenchorhynchus swarupi</i> Singh & Khera, 1978 Genus TROPHURUS Loof, 1956 32. <i>Trophurus clavicaudatus</i> Sen, Chatterjee & Manna, 2012
Family HOPLOLAIMIDAE Filipjev, 1934 Subfamily HOPLOLAIMINAE Filipjev, 1934 Genus HOPLOLAIMUS Daday, 1905 33. <i>Hoplolaimus indicus</i> Sher, 1963
Genus SCUTELLONEMA Andrassy, 1958 34. <i>Scutellonema bengalensis</i> Sen, 2019
Family PRATYLENCHIDAE Thorne, 1949 Genus HIRSCHMANNIELLA Luc & Goodey, 1963 35. <i>Hirschmanniella oryzae</i> (van Breda De Hann, 1902) Luc & Goodey, 1963 36. <i>Hirschmanniella gracilis</i> (de Mann, 1880) Luc & Goodey, 1963

## 5. List of Soil Mites

Taxa
Family: HYPOCHTHONIDAE Berlese, 1910 Genus <i>Nothrolhmannia</i> Balogh, 1968 1. <i>Nothrolhmannia calcarata</i> Balogh, 1968
Family LOHMANIIDAE Berlese, 1916 Genus <i>Papillacarus</i> Kunst, 1959 2. <i>Papillacarus hirsutus</i> Aoki, 1961 Genus <i>Javacarus</i> Balogh, 1961 3. <i>Javacarus (Javacarus) kuehnelti</i> Balogh, 1961 Genus <i>Annectacarus</i> Grandjean, 1950 4. <i>Annectacarus</i> sp.
Family ORIBOTRITIIDAE Balogh, 1943 Genus <i>Indotritia</i> Jacot, 1929 5. <i>Indotritia</i> sp.
Family PHTHIRACARIDAE Perty, 1841 Genus <i>Hoplophorella</i> Berlese, 1923 6. <i>Hoplophorella (Hoplophorella) vitrina</i> Berlese, 1913 7. <i>Hoplophorella lanceoseta</i> (Balogh & Mahunka, 1981)
Family NOTHRIDAE Berlese, 1896 Genus <i>Nothrus</i> Koch, 1836 8. <i>Nothrus</i> sp.
Family BASILOBELBIDAE Balogh, 1961 Genus <i>Basilobelba</i> Balogh, 1958 9. <i>Basilobelba</i> sp.
Family NIPPOBODIDAE Aoki, 1959 Genus <i>Nippobodes</i> Aoki, 1959 10. <i>Nippobodes</i> sp.
Family DAMPFIELLIDAE Balogh, 1961 Genus <i>Dampella</i> Sellnick, 1931 11. <i>Dampella prostrata</i> Aoki, 1965
Family OPPIIDAE Sellnick, 1937 Genus <i>Lasiobelba</i> Aoki, 1959 12. <i>Lasiobelba (Lasiobelba) kuehnelti</i> Csiszar, 1961 Genus <i>Brachioppiella</i> Hammer, 1962 13. <i>Brachioppiella (Brachioppiella) periculosa</i> Hammer, 1962 Genus <i>Multioppia</i> Hammer, 1961

<p>14. <i>Multioppia stellifera</i> Hammer, 1961 Genus <i>Amerioppia</i> Hammer, 1961</p> <p>15. <i>Amerioppia</i> sp. Family ARCEREMAEIDAE Balogh, 1972 Genus <i>Arceremaeus</i> Hammer, 1961</p> <p>16. <i>Arceremaeus incaensis</i> Hammer, 1961* Family TECTOCEPHEIDAE Grandjean, 1954 Genus <i>Tectocephus</i> Berlese, 1896</p> <p>17. <i>Tectocephus</i> sp.</p> <p>18. <i>Tectocephus velatus velatus</i> Michael, 1880 Family AUSTRACHIPTERIIDAE Luxton, 1985 Genus <i>Lamellobates</i> Hammer, 1958</p> <p>19. <i>Lamellobates molecula</i> Berlese, 1916 Genus <i>Hypozetes</i> Balogh, 1959 <i>Hypozetes laysanensis</i> Aoki, 1964</p> <p>Family ORIBATELLIDAE Jacot, 1925 Genus <i>Cuspidozetes</i> Hammer, 1962</p> <p>20. <i>Cuspidozetes armatus</i> Hammer, 1962* Genus <i>Lamellobates</i> Hammer, 1958 <i>Lamellobates molecula</i> Berlese, 1916 Family DRYMOBATIDAE J. &amp; P. Balogh, 1984 Genus <i>Rykella</i> Balogh, 1962</p> <p>21. <i>Rykella insignis</i> Balogh, 1962 Family MOCHLOZETIDAE Grandjean, 1960 Genus <i>Mochlozetes</i> Grandjean, 1930</p> <p>22. <i>Mochlozetes penetrabilis</i> Grandjean, 1930* Family ORIBATULIDAE Thor, 1929 Genus <i>Oribatula</i> Berlese, 1896</p> <p>23. <i>Oribatula lata</i> Hammer, 1961 Family SCHELORIBATIDAE Jacot, 1935 Genus <i>Monoschelobates</i> Balogh &amp; Mahunka, 1969</p> <p>24. <i>Monoschelobates parvus</i> Balogh &amp; Mahunka, 1969 Genus <i>Scheloribates</i> Berlese, 1908</p> <p>25. <i>Scheloribates (Scheloribates) curvialatus</i> Hammer, 1961</p> <p>26. <i>Scheloribates thermophilus</i> Hammer, 1961</p> <p>27. <i>Scheloribates (Scheloribates) huancayensis</i> Hammer, 1961</p> <p>28. <i>Scheloribates (Scheloribates) pallidulus</i> Koch, 1841</p> <p>29. <i>Scheloribates pauliensis</i> Perez-Inigo &amp; Baggio, 1980</p> <p>30. <i>Scheloribates elagantulus</i> Hammer, 1961</p> <p>31. <i>Scheloribates</i> sp. Genus <i>Perscheloribates</i> Hammer, 1973</p> <p>32. <i>Perscheloribates (Perscheloribates) albialatus</i> Hammer, 1961 Family PROTORIBATIDAE J. and P. Balogh, 1984 Genus <i>Protoribates</i> Berlese, 1908</p> <p>33. <i>Protoribates (Protoribates) magnus</i> Aoki, 1982</p> <p>34. <i>Protoribates (Protoribates) capucinus</i> Berlese, 1908 Genus <i>Vilhenabates</i> Balogh, 1963</p> <p>35. <i>Vilhenabates minutus</i> Balogh, 1958*</p> <p>36. <i>Vilhenabates (Phalacrozetes) sinatus</i> Aoki, 1965 Genus <i>Setoxylobates</i> Balogh and Mahunka, 1967</p> <p>37. <i>Setoxylobates foveolatus</i> Balogh and Mahunka, 1967</p> <p>Family HAPLOZETIDAE Grandjean, 1936 Genus <i>Indoribates (Haplozetes)</i> Willman, 1935</p> <p>38. <i>Indoribates</i> sp. Genus <i>Trachyoribates</i> Berlese, 1908</p> <p>39. <i>Trachyoribates ovulum</i> Berlese, 1908</p> <p>40. <i>Trachyoribates pinguis</i> Balogh &amp; Mahunka, 1978</p> <p>41. <i>Trachyoribates</i> sp.</p>
--

Genus <i>Pilobatella</i> Balogh & Mahunka, 1967
42. <i>Pilobatella punctata</i> Balogh & Mahunka, 1967
Family GALUMNIDAE Jacot, 1925
Genus <i>Galumna</i> Heyden, 1826
43. <i>Galumna abellifera</i> Hammer, 1958
44. <i>Galumna australis</i> (Berlese, 1914)
45. <i>Galumna pallida</i> Hammer, 1958
46. <i>Galumna</i> sp.
Genus <i>Pergalumna</i> Grandjean, 1936
47. <i>Pergalumna numerosa</i> Sellnick, 1923
48. <i>Pergalumna magnipora</i> Hammer, 1961
Genus <i>Cryptogalumna</i> Grandjean, 1957
49. <i>Cryptogalumna cryptodonta</i> Grandjean, 1957
Genus <i>Psammogalumna</i> Balogh, 1943
50. <i>Psammogalumna hungarica</i> Sellnick, 1925

## 6. List of Spiders

Taxa
Family: ARANEIDAE
1. <i>Araneus mitificus</i> (Simon, 1886)
2. <i>Argiope</i> sp.
3. <i>Argiope catenulate</i> (Doleschall, 1859)
4. <i>Cyclosa neilensis</i> Tikader, 1977
5. <i>Eriovixia</i> sp.
Family: CLUBIONIDAE
6. <i>Clubiona filicata</i> O. Pickard-Cambridge, 1874
Family: HERSILIIDAE
7. <i>Hersilia savignyi</i> Lucas, 1836
Family: LYCOSIDAE
8. <i>Lycosa bistrata</i> Gravely, 1924
9. <i>Pardosa pseudoannulata</i> (Bösenberg & Strand, 1906)
10. <i>Pardosa sumatrana</i> (Thorell, 1890)
11. <i>Trochosa punctipes</i> (Gravely, 1924)
12. <i>Trochosa</i> sp.
13. <i>Wadicosa quadrifera</i> (Gravely, 1924)
Family: OXYOPIDAE
14. <i>Oxyopes hindostanicus</i> Pocock, 1901
Family: SALTICIDAE
15. <i>Hyllus semicupreus</i> (Simon, 1885)
16. <i>Menemerus bivittatus</i> (Dufour, 1831)
17. <i>Myrmarachne melanocephala</i> MacLeay, 1839
18. <i>Phaeacius lancearius</i> (Thorell, 1895)
19. <i>Plexippus paykulli</i> (Audouin, 1826)
20. <i>Rhene rubrigerax</i> (Thorell, 1887)
21. <i>Rhene</i> sp.
Family: TETRAGNATHIDAE
22. <i>Guizygiella</i> sp.
23. <i>Guizygiella melanocrania</i> (Thorell, 1887)
24. <i>Tetragnatha javana</i> (Thorell, 1890)
25. <i>Tetragnatha keyserlingi</i> Simon, 1890
26. <i>Tetragnatha nitens</i> (Audouin, 1826)
27. <i>Tetragnatha</i> sp.
28. <i>Tetragnatha vermiformis</i> Emerton, 1884

Family: THERIDIIDAE
29. <i>Argyrodes argentatus</i> O. Pickard-Cambridge, 1880
Family: THOMISIDAE
30. <i>Camaricus formosus</i> Thorell, 1887
31. <i>Runcinia insecta</i> (L. Koch, 1875)
32. <i>Thomisus</i> sp.

## 7. List of Crustacea: Cladocerans

Taxa
Phylum Arthropoda Super Class Crustacea Class Branchiopoda Super Order Cladocera Order Ctenopoda
Family Sididae 1 <i>Diaphanosoma excisum</i> Sars, 1885 2 <i>D. sarsi</i> Richard, 1894 3 <i>Pseudosida bidentata</i> Herrick, 1884 Order Anomopoda
Family Daphnidae 4 <i>Ceriodaphnia cornuta</i> Sars, 1885 5 <i>Ceriodaphnia</i> sp. 6 <i>Daphnia</i> sp. 7 <i>Simocephalus vetulus</i> (O.F. Muller, 1776)
Family Moinidae 8 <i>Moina micrura</i> Kurz, 1874 9 <i>Moina</i> sp. 10 <i>Moinadaphnia macleayi</i> (King, 1853) 11 <i>Moinodaphnia</i> sp.
Family Bosminidae 12 <i>Bosmina longirostris</i> (Muller, 1785) 13 <i>Bosmina</i> sp.
Family Macrothricidae 14 <i>Macrothrix goeldii</i> Richard, 1897
Family Chydoridae Subfamily Chydorinae 15 <i>Chydorus sphaericus</i> (O.F. Muller, 1776) 16 <i>Coronatella rectangula</i> (G.O. Sars, 1862) 17 <i>Pleuroxus siilis</i> Vavra, 1900
Subfamily Aloninae 18 <i>Alona affinis</i> (Leydig, 1860) 19 <i>Alona quadrangularis</i> (O.F. Muller, 1776) 20 <i>Alona</i> sp. 21 <i>Biapertura karua</i> (King, 1853) 22 <i>Notoalona globulosa</i> (Daday, 1898) 23 <i>Kurzia longirostris</i> (Daday, 1898) 24 <i>Kurzia</i> sp. 25 <i>Cladoceran neonates</i>

## 8. List of Crustacea: Ostracods

Taxa
Phylum Arthropoda Superclass Crustacea Class Ostracoda Order Podocopida Suborder Podocopa
Family: Cyprididae 1. <i>Cypris subglobosa</i> Sowerby, 1840 2. <i>Strandesia</i> sp. Subfamily Stenocyprinae 3. <i>Stenocypris major</i> (Baird) 1859

## 9. List of Crustacea: Copepods

Taxa
Phylum Arthropoda Superclass Crustacea Class Copepoda Order Calanidae Subfamily Diaptomidae 1. <i>Heliodiaptomus contortus</i> (Gurney, 1907) 2. <i>Heliodiaptomus viduus</i> (Gurney, 1916)
Family: Pseudodiaptomidae 3. <i>Pseudodiaptomus annandalei</i> Sewell, 1919
Family: Cyclopididae Subfamily Eucyclopiniae 4. <i>Eucyclops</i> sp. 5. <i>Mesocyclops hyalinus</i> (Rehberg, 1880) 6. <i>Mesocyclops leuckarti</i> (Claus, 1857) 7. <i>Microcyclops varicans</i> Sars, 1863 8. <i>Thermocyclops rylovi rylovi</i> (Smirnov, 1928) Order Harpacticoida 9. <i>Euterpina acutifrons</i> (Dana 1847) 10. <i>Cyclopoid nauplii</i> 11. <i>Cyclopoid copepodite</i>



## 10. List of Crustacea: Crabs and Shrimps

<b>Taxa</b>
<p>Kingdom Animalia</p> <p>Phylum Arthropoda</p> <p>Subphylum Crustacea</p> <p>Class Malacostraca</p> <p>Subclass Eumalacostraca</p> <p>Superorder Eucarida Calman, 1904</p> <p>Order Decapoda Latreille, 1802</p> <p>Suborder Dendrobranchiata Spence Bate, 1888</p> <p>Superfamily Penaeoidea Rafinesque, 1815</p> <p>Family Penaeidae Rafinesque, 1815</p> <p>Genus <i>Metapenaeus</i> Wood-Mason in Wood-Mason &amp; Alcock, 1891</p>
<p>1. <i>Metapenaeus brevicornis</i> (H. Milne Edwards 1837 [in Milne Edwards, 1834-1840]) Genus <i>Penaeus</i> Fabricius, 1798</p>
<p>2. <i>Penaeus merguensis</i> de Man, 1888 [in de Man, 1887-1888]</p>
<p>3. <i>Penaeus semisulcatus</i> De Haan, 1844 [in De Haan, 1833-1850]</p>
<p>4. <i>Penaeus indicus</i> H. Milne Edwards, 1837</p> <p>Suborder Pleocymata Burkenroad, 1963</p> <p>Infraorder Anomura Macleay, 1838 (Hermit crabs)</p> <p>Superfamily Paguroidea Latreille, 1802</p> <p>Family Diogenidae Ortman, 1892</p> <p>Genus <i>Clibanarius</i> Dana, 1852</p>
<p>5. <i>Clibanarius padavensis</i> de Man, 1888 [in de Man, 1887-1888]</p> <p>Infraorder Caridea Dana, 1852 (Prawns/Shrimps)</p> <p>Superfamily Atyoidea De Haan, 1849 (in De Haan, 1833-1850)</p> <p>Family Atyidae De Haan, 1849 (in De Haan, 833-1850)</p> <p>Genus <i>Caridina</i> H. Milne Edwards, 1837</p>

(in H. Milne Edwards, 1834-1840)
6. <i>Caridina gracilirostris</i> de Man, 1892
7. <i>Caridina gracilipes</i> De Man, 1892
8. <i>Caridina propinqua</i> De Man, 1908 Superfamily Palaemonoidea Rafinesque, 1815 Family Palaemonidae Rafinesque, 1815 Genus <i>Leptocarpus</i> Holthuis, 1950
9. <i>Leptocarpus fluminicola</i> (Kemp, 1917) Genus <i>Macrobrachium</i> Spence Bate, 1868
10. <i>Macrobrachium lamarrei lamarrei</i> (H.M.Edwards)
11. <i>Macrobrachium malcolmsonii</i> (H. Milne-Edwards, 1844)
12. <i>Macrobrachium rosenbergii</i> (de Man, 1879)
13. <i>Macrobrachium rude</i> (Heller, 1862)
14. <i>Macrobrachium</i> sp. Genus <i>Palaemon</i> Weber, 1795
15. <i>Palaemon styliferus</i> H. Milne Edwards, 1840 [in H. Milne Edwards, 1834-1840- De Man (1908)] Infraorder Brachyura, Latreille, 1802 (Crabs) Section Eubrachyura, Saint Laurent, 1980 Subsection Thoracotremata Guinot, 1977 Superfamily Grapsoidea, MacLeay, 1875 Family <i>Varunidae</i> H. Milne Edwards, 1853 Genus <i>Varuna</i> H. Milne Edwards
16. <i>Varuna litterata</i> (Fabricius, 1798)-De Man (1908) Superfamily Ocypodoidea Rafinesque, 1815 Family Ocypodidae, Rafinesque, 1815 Genus <i>Tubuca</i> Bott, 1973
17. <i>Tubuca acuta</i> (Stimpson, 1858) Subsection Heterotremata Guinot, 1977 Superfamily Pilumnoidea Samouelle, 1819 Family Pilumnidae Samouelle, 1819 Genus <i>Benthopanope</i> Davie, 1989

<p>18. <i>Benthopanope indica</i> (de Man, 1887 [in de Man, 1887-1888])</p> <p>Superfamily Gecarcinucoidea Rathbun, 1904</p> <p>Family Gecarcinucidae Rathbun, 1904</p> <p>Genus <i>Sartoriana</i> Bott, 1969</p>
<p>19. <i>Sartoriana spinigera</i> (Wood-Mason, 1871)</p> <p>Superorder Peracarida Calman, 1904</p> <p>Order Isopoda Latreille, 1817</p> <p>Suborder Cymothoida Wagele, 1989</p> <p>(A parasitic Crustacean)</p> <p>Superfamily Bopyroidea Rafinesque, 1815</p> <p>Family Bopyridae Rafinesque, 1815</p> <p>Genus <i>Probopyrus</i> Giard &amp; Bonnier, 1888</p>
<p>20. <i>Probopyrus prashadi</i> (Chopra, 1923)</p> <p>Superorder Peracarida Calman, 1904</p> <p>Order Isopoda Latreille, 1817</p> <p>Suborder Oniscidea Latreille, 1802</p> <p>Family Porcellionidae, Brant, 1831</p> <p>Genus <i>Porcellio</i> Latreille, 1804</p>
<p>21. <i>Porcellio assamensis</i> Chopra, 1924</p> <p>Order Mysida Boas, 1883 (Opposum Shrimps)</p> <p>Family Mysidae Haworth, 1825</p> <p>Genus <i>Gangemysis</i> Derzhavin, 1924</p>
<p>22. <i>Gangemysis assimilis</i> (W. Tattersall, 1908)</p> <p>Genus <i>Mesopodopsis</i> Czemiavsky, 1882</p>
<p>23. <i>Mesopodopsis orientalis</i> (W. Tattersall, 1908)</p> <p>Order Amphipoda</p> <p>Suborder Gammaridea</p>
<p>24. <i>Gammarid amphipods</i></p>

## 11. List of Apterygotan fauna

Taxa
COLLEMBOLA
Family Neanuridae
1. <i>Protanura carpenteri</i> Mukherjee, 1932
2. <i>Paleonura siva</i> (Yosii, 1966) Cassagnau, 1982
3. <i>Friesea excelsa</i> Denis, 1936
Family Hypogastruridae
4. <i>Ceratophysella indica</i> (Salmon, 1956)
5. <i>Ceratophysella indovaria</i> (Salmon, 1970)
6. <i>Xenylla obscura</i> Imms, 1912
7. <i>Xenylla reducta</i> Prabhoo, 1971
Family Onychiuridae
8. <i>Allonychiurus indicus</i> (Choudhuri & Roy, 1965) Pomorski, 2000
Family Tullbergidae
9. <i>Paratullbergia indica</i> Salmon, 1965
Family Brachystomellidae
10. <i>Brachystomella terrafolia</i> Salmon, 1944
Family Isotomidae
11. <i>Isotomodes dagamae</i> Prabhoo, 1971
12. <i>Hemisotoma thermophila</i> (Axelson, 1900) Bagnall, 1949
13. <i>Ballistura bengalensis</i> Yosii, 1966
14. <i>Isotomurus balteatus</i> (Reuter, 1876) Handschin, 1929
15. <i>Isotomurus palustris</i> (Muller, 1776) Borner, 1906
16. <i>Isotomurus ciliatus</i> Stach, 1947
17. <i>Folsomides purvulus</i> Stach, 1922
Family Entomobryidae
18. <i>Homidia cingula</i> (Borner, 1906) Yosii, 1959
19. <i>Sinella curviseta</i> Brook, 1882
20. <i>Lepidocyrtus (Lanocyrtus) caeruleicornis</i> Bonet, 1930
21. <i>Lepidocyrtus (Lepidocyrtus) curvicollis</i> Bourlet, 1841
22. <i>Lepidocyrtus (Acrocyrtus) heterolepis</i> Yosii, 1959
23. <i>Lepidocyrtus (Acrocyrtus) malayanus</i> Yosii, 1959
24. <i>Lepidocyrtus (Cinctocyrtus) medius</i> Schaeffer, 1898
25. <i>Lepidocyrtus (Ascocyrtus) scaber</i> Ritter, 1911
26. <i>Lepidocyrtus magnificus</i> Carpenter, 1924
27. <i>Dicranocentrus indicus</i> Bonet, 1930
28. <i>Dicranocentrus cercifer</i> (Imms, 1912) Mari Mutt, 1979
29. <i>Alloscopus tetracanthus</i> (Borner, 1906)
30. <i>Seira indica</i> (Ritter, 1911) Yosii, 1966
31. <i>Seira indra</i> (Imms, 1912)
32. <i>Seira lateralis</i> Yossi, 1966
33. <i>Seira punctata</i> (Ritter, 1911)

Family Paronellidae
34. <i>Salina bengalensis</i> Mitra, 1973
35. <i>Salina celebensis</i> (Schaeffer, 1898) Yosii, 1959
36. <i>Salina indica</i> (Imms, 1912) Yosii, 1960
37. <i>Salina striata</i> (Handschin, 1928)
38. <i>Salina yosii</i> Salmon, 1964
39. <i>Dicranocentroides flavescens</i> Yosii, 1966
40. <i>Cyphoderopsis ceylonica</i> Yosii, 1966
41. <i>Cyphoderus javanus</i> Börner, 1906
42. <i>Cyphoderus albinus</i> Nicolet, 1842
Family Neelidae
43. <i>Neelus murinus</i> Folsom, 1896
44. <i>Megalothorax minimus</i> Willem, 1900
Family Arrhopalitidae
45. <i>Pygmarrhopalites habeii</i> (Yosii, 1956)
Family Sminthuridae
46. <i>Sminthurus parvullus</i> Ritter, 1911
47. <i>Sminthurus viridis</i> (Linn. 1758) Bourlet, 1843
48. <i>Sphyrotheca gangetica</i> Yosii, 1966
Family Dicyrtomidae
49. <i>Calvatomina pagoda</i> (Yosii, 1966)
DIPLURA
50. <i>Lepidocampa juradoi bengalensis</i> Rani & Mitra, 1977
ZYGENTOMA
Family Lepismatidae
51. <i>Acrotelsa collaris</i> (Fabricius, 1793)
52. <i>Ctenolepisma longicaudata</i> Escherich, 1905
53. <i>Ctenolepisma ciliata</i> (Dufour, 1831)
54. <i>Ctenolepisma nigra</i> (Oudemans, 1890)

## 12. List of Odonata: Dragonflies and Damselflies

<b>Taxa</b>
Order: Odonata Suborder: Zygoptera
Superfamily: Coenagrionoidea Family: Coenagrionidae
1. <i>Agriocnemis pygmaea</i> (Rambur)
2. <i>Ceriagrion cerinorubellum</i> (Brauer)
3. <i>Ceriagrion coromandelianum</i> (Fabricius)
4. <i>Ischnura aurora</i> (Brauer)
5. <i>Ischnura senegalensis</i> (Rambur)
6. <i>Onychargia atrocyana</i> (Selys)
7. <i>Pseudagrion microcephalum</i> (Rambur)
8. <i>Pseudagrion rubriceps</i> (Selys)
Suborder: Anisoptera Superfamily: Aeshnoidea
Family: Aeshnidae

9. *Anax guttatus* (Burmeister)  
Superfamily: Gomphoidea Family: Gomphidae  
Superfamily: Libelluloidea Family: Libellulidae
10. *Ictinogomphus rapax* (Rambur)  
11. *Brachydiplax chalybea* (Rambur)  
12. *Brachythemis contaminata* (Fabricius)  
13. *Bradinopyga geminata* (Rambur)  
14. *Crocothemis servilia* (Drury)  
15. *Diplacodes trivialis* (Rambur)  
16. *Lathrecista asiatica* (Fabricius)  
17. *Macrodiplax cora* (Brauer)  
18. *Neurothemis tullia* (Drury)  
19. *Orthetrum pruinosum* (Burmeister)  
20. *Orthetrum sabina* (Drury)  
21. *Pantala flavescens* (Fabricius)  
22. *Rhodothemis rufa* (Rambur)  
23. *Rhyothemis variegata* (Linnaeus)  
24. *Trithemis aurora* (Burmeister)  
25. *Trithemis festiva* (Rambur)  
26. *Trithemis pallidinervis* (Kirby)

### 13. List of Orthoptera: Grasshoppers and Crickets

Taxa
1. <i>Acrida exaltata</i> (Walker, 1859) Tribe Phlaeobini Brunner von Wattenwyl, 1893 Genus <i>Phlaeoba</i> Stål, 1860
2. <i>Phlaeoba infumata</i> Brunner von Wattenwyl, 1893
3. <i>Phlaeoba panteli</i> Bolivar, 1902 Subfamily Catantopinae Genus <i>Diabolocatantops</i> Jago, 1984
4. <i>Diabolocatantops innotabilis</i> (Walker, 1870) Genus <i>Choroedocus</i> Bolivar, 1914
5. <i>Choroedocus capensis</i> (Thunberg, 1815)
6. <i>Choroedocus robustus</i> (Serville, 1839) Tribe Oxyrrhepini Tinkham, 1940 Genus <i>Oxyrrhepes</i> Stål, 1873
7. <i>Oxyrrhepes obtusa</i> (Haan, 1842) Genus <i>Pachyacris</i> Uvarov, 1923
8. <i>Pachyacris vinosa</i> (Walker, 1870) Tribe Paraconophymatini Otte, 1995 Genus <i>Paraconophyma</i> Uvarov, 1921
9. <i>Paraconophyma scabra</i> (Walker, 1870) Genus <i>Xenocatantops</i> Dirsh & Uvarov, 1953
10. <i>Xenocatantops humilis</i> (Serville, 1839) Subfamily Eyeprepocnemidinae Genus <i>Eyprepocnemis</i> Fieber, 1853
11. <i>Eyprepocnemis alacris alacris</i> (Serville, 1839)

Genus <i>Eupreponotus</i> Uvarov, 1921
12. <i>Eupreponotus inflatus</i> Uvarov, 1921 Genus <i>Heteracris</i> Walker, 1870
13. <i>Heteracris pulchra</i> (Bolivar, 1902) Genus <i>Tylotropidius</i>
14. <i>Tylotropidius varicornis</i> (Walker, 1870) Subfamily Gomphocerinae Genus <i>Aulacobothrus</i> Bolivar, 1902
15. <i>Aulacobothrus luteipes luteipes</i> (Walker, 1871) Genus <i>Leva</i> Bolivar, 1909
16. <i>Leva indica</i> (Bolivar, 1902) Subfamily Hemiacridinae Tribe Hieroglyphini Genus <i>Hieroglyphus</i> Krauss, 1877
17. <i>Hieroglyphus banian</i> (Fabricius, 1798) Subfamily Oedipodinae Genus <i>Aiolopus</i> Fieber, 1853
18. <i>Aiolopus thalassinus tamulus</i> (Fabricius, 1798) Genus <i>Gastrimargus</i> Saussure, 1884
19. <i>Gastrimargus africanus africanus</i> (Saussure, 1888) Genus <i>Heteropternis</i> Stål, 1873
20. <i>Heteropternis respondens</i> (Walker, 1859) Genus <i>Morphacris</i> Walker, 1870
21. <i>Morphacris fasciata</i> (Thunberg, 1815) Genus <i>Oedaleus</i> Fieber, 1853
22. <i>Oedaleus abruptus</i> (Thunberg, 1815)
23. <i>Oedaleus senegalensis</i> (Krauss, 1877) Genus <i>Trilophidia</i> Stål, 1873
24. <i>Trilophidia annulata</i> (Thunberg, 1815) Genus <i>Dittopternis</i> Saussure, 1884
25. <i>Dittopternis venusta</i> (Walker, 1870) Genus <i>Locusta</i> Linnaeus, 1758
26. <i>Locusta migratoria migratoria</i> (Linnaeus, 1758) Subfamily Coptacridinae Genus <i>Epistaurus</i> Bolivar, 1889
27. <i>Epistaurus sinetyi</i> Bolivar, 1902
28. <i>Eucoptacra saturata</i> (Walker, 1870) Subfamily Oxyinae Genus <i>Gesonula</i> Uvarov, 1940
29. <i>Gesonula punctifrons</i> (Stål, 1861) Genus <i>Oxya</i> Serville, 1831
30. <i>Oxya fuscovittata</i> (Marschall, 1836)
31. <i>Oxya hyla</i> Serville, 1831
32. <i>Oxya nitidula</i> (Walker, 1870)
33. <i>Oxya velox</i> (Fabricius, 1787) Subfamily Tropidopolinae Tribe Tristriini Mishchenko, 1945 Genus <i>Tristria</i> Stål, 1873
34. <i>Tristria pulvinata</i> (Uvarov, 1921) Tribe Tropidopolini Jacobson, 1905 Genus <i>Tropidopola</i> Stål, 1873
35. <i>Tropidopola longicornis</i> (Fieber, 1853) Subfamily Cyrtacanthacridinae Genus <i>Cyrtacanthacris</i> Walker, 1870

36. <i>Cyrtacanthacris tataricatatarica</i> (Linnaeus, 1758) Genus <i>Patanga</i> Uvarov, 1923
37. <i>Patanga succincta</i> (Johansson, 1763) Genus <i>Schistocerca</i> Stål, 1873
38. <i>Schistocerca gregaria</i> (Forskål, 1775) Subfamily Spathosterninae Genus <i>Spathosternum</i> Krauss, 1877
39. <i>Spathosternum prasiniferum prasiniferum</i> (Walker, 1871) Superfamily Pyrgomorphoidea Family Pyrgomorphidae Genus <i>Atractomorpha</i> Saussure, 1862
40. <i>Atractomorpha crenulata</i> (Fabricius, 1793)
41. <i>Atractomorpha psittacina</i> (Haan, 1842) Tribe Tagastini Bolivar, 1905 Genus <i>Tagasta</i> Bolivar, 1905
42. <i>Tagasta indica</i> Bolivar, 1905 Genus <i>Chrotogonus</i> Serville, 1838
43. <i>Chrotogonus (Chrotogonus) trachypterus trachypterus</i> (Blanchard, 1836) Superfamily Tetrigoidea Family Tetrigidae Subfamily Scelimeninae Genus <i>Indoscelimena</i> Günther, 1938
44. <i>Indoscelimena angulata</i> (Hancock, 1915)
45. <i>Indoscelimena flavopicta</i> (Bolivar, 1909)
46. <i>Indoscelimena saussurei</i> (Hancock, 1915) Tribe Thoradontini Kevan, 1966 Genus <i>Thoradonta</i> Hancock, 1909
47. <i>Thoradonta nodulosa</i> (Stål, 1860)
48. <i>Thoradonta spiculoba</i> Hancock, 1912
49. <i>Thoradonta bengalensis</i> Shishodia, 1991 Tribe Criotettigini Kevan, 1966 Genus <i>Criotettix</i> Bolivar, 1887
50. <i>Criotettix bispinosus</i> (Dalman, 1818)
51. <i>Criotettix inornatus</i> (Walker, 1871) Genus <i>Eucriotettix</i> Hebard, 1929
52. <i>Eucriotettix rufescens</i> (Kirby, 1914) Genus <i>Loxilobus</i> Hancock, 1904
53. <i>Loxilobus striatus</i> Hancock, 1915 Subfamily Tetriginae Genus <i>Coptotettix</i> Bolivar, 1887
54. <i>Coptotettix hancockus</i> Shishodia & Varshney, 1987 Genus <i>Euparatettix</i> Hancock, 1904
55. <i>Euparatettix indicus</i> (Bolivar, 1887)
56. <i>Euparatettix personatus</i> (Bolivar, 1887)
57. <i>Euparatettix tricarinatus</i> (Bolivar, 1887) Genus <i>Hedotettix</i> Bolivar, 1887
58. <i>Hedotettix costatus</i> Hancock, 1912
59. <i>Hedotettix gracilis</i> (Haan, 1842) Genus <i>Paratettix</i> Bolivar, 1887
60. <i>Paratettix histricus</i> (Stål, 1861)
61. <i>Paratettix variabilis</i> Bolivar, 1887 Genus <i>Ergatettix</i> Kirby, 1914
62. <i>Ergatettix dorsifera</i> (Walker, 1871)



63. <i>Ergatettix interruptus</i> (Brunner von Wattenwyl, 1893) Suborder ENSIFERA Infraorder Oedischioidea Superfamily Grylloidea Family Gryllidae Subfamily Gryllinae Genus <i>Acheta</i> Fabricius, 1775
64. <i>Acheta domesticus</i> (Linnaeus, 1758) Genus <i>Gryllus</i> Linnaeus, 1758 Subgenus <i>Gryllus</i> Linnaeus, 1758
65. <i>Gryllus (Gryllus) bimaculatus</i> De Geer, 1773 Genus <i>Gryllodes</i> Saussure, 1874
66. <i>Gryllodes sigillatus</i> (Walker, 1869) Genus <i>Modicogryllus</i> Chopard, 1961 Subgenus <i>Modicogryllus</i> Chopard, 1961
67. <i>Modicogryllus (Modicogryllus) confirmatus</i> (Walker, 1859) Genus <i>Plebeiogryllus</i> Randell, 1964
68. <i>Plebeiogryllus guttiventris guttiventris</i> (Walker, 1871) Genus <i>Velarifictorus</i> Randell, 1964 Subgenus <i>Velarifictorus</i> Randell, 1964
69. <i>Velarifictorus (Velarifictorus) aspersus</i> (Walker, 1869) Genus <i>Loxoblemmus</i> Saussure, 1877
70. <i>Loxoblemmus equestris</i> Saussure, 1877 Subfamily Oecanthinae Tribe Oecanthini Genus <i>Oecanthus</i> Serville, 1831
71. <i>Oecanthus rufescence</i> Serville, 1839 Family Trigonidiidae Subfamily Nemobiinae Tribe Nemobiini Saussure, 1877 Genus <i>Dianemobius</i> Vickery, 1973
72. <i>Dianemobius fascipes</i> (Walker, 1869) Genus <i>Nemobius</i> Serville, 1838
73. <i>Nemobius strigipennis</i> (Chopard, 1928) Genus <i>Pteronemobius</i> Jacobson and Bianchi, 1905 Subgenus <i>Pteronemobius</i> Jacobson, 1904
74. <i>Pteronemobius (Pteronemobius) heydenii</i> concolor (Walker, 1871) Genus <i>Polionemobius</i> Gorochov, 1983
75. <i>Polionemobius taprobanensis</i> (Walker, 1869) Subfamily Trigonidiinae Tribe Trigonidiini Genus <i>Amusurgus</i> Brunner von Wattenwyl, 1893 Subgenus <i>Amusurgus</i> Brunner von Wattenwyl, 1893
76. <i>Amusurgus (Amusurgus) fulvus</i> Chopard, 1969
77. <i>Amusurgus (Amusurgus) oedemeroides</i> (Walker, 1871) Genus <i>Natula</i> Gorochov, 1874
78. <i>Natula longipennis</i> (Serville, 1838) Genus <i>Homoeoxipha</i> Saussure, 1874
79. <i>Homoeoxipha lycoides</i> (Walker, 1869) Subfamily Euscyrinae Genus <i>Euscyrtus</i> Guérin-Méneville, 1844 Subgenus <i>Osus</i> Gorochov, 1987
80. <i>Euscyrtus (Osus) concinnus</i> (Haan, 1842)
81. <i>Euscyrtus (Osus) hemelytrus</i> (Haan, 1842) Family Phalangopsidae

Subfamily Cachoplistinae Tribe Cachoplistini Saussure, 1877 Genus <i>Cacoplistes</i> Brunner, 1873 Subgenus <i>Laminogryllus</i> Gorochoy, 2003
82. <i>Cacoplistes (Laminogryllus) rogenhoferi</i> Saussure, 1877 Family Gryllotalpidae Subfamily Gryllotalpinae Genus <i>Gryllotalpa</i> Latreille, 1802
83. <i>Gryllotalpa africana</i> Beauvois, 1805 Superfamily Tettigonioidea Family Tettigoniidae Subfamily Conocephalinae Tribe Copiphorini Karny, 1912 Genus <i>Euconocephalus</i> Karny, 1907
84. <i>Euconocephalus incertus</i> (Walker, 1869)
85. <i>Euconocephalus pallidus</i> (Redtenbacher, 1891) Subfamily Phaneropterinae Genus <i>Trigonocorypha</i> Stål, 1873
86. <i>Trigonocorypha unicolor</i> (Stoll, 1787) Subfamily Pseudophyllinae Genus <i>Sathrophyllia</i> Stål, 1874
87. <i>Sathrophyllia femorata</i> (Fabricius, 1787) Genus <i>Onomarchus</i> Stål, 1874
88. <i>Onomarchus leuconotus</i> (Serville, 1839) Subfamily Pseudophyllinae Tribe Phyllomimini Genus <i>Acanthoprion</i> Pictet & Saussure, 1892
89. <i>Acanthoprion suspectum</i> (Brunner von Wattenwyl, 1895) Tribe Cocconotini Genus <i>Meronicidius</i> Serville, 1831
90. <i>Meronicidius ochraceus</i> (Stoll, 1813) Subfamily Listroscelidinae Tribe Phisidini Jin, Xingbao, 1987 Genus <i>Phisis</i> Stål, 1860
91. <i>Phisis pectinata</i> (Guérin, 1832) Subfamily Meconematinae Genus <i>Xiphidiopsis</i> Redtenbacher, 1891 Subgenus <i>Xiphidiopsis</i> Redtenbacher, 1891
92. <i>Xiphidiopsis (Xiphidiopsis) straminula</i> (Walker, 1871)

#### 14. List of Blattodea: Isoptera- Termites

Taxa
Order BLATTODEA Wattenwyl, 1882
Infraorder ISOPTERA Brulle, 1832
Family RHINOTERMITIDAE Froggatt, 1897
Subfamily COPTOTERMITINAE Holmgren, 1910
Genus <i>Coptotermes</i> Wasmann, 1896
1. <i>Coptotermes ceylonicus</i> Holmgren, 1911
2. <i>Coptotermes kishori</i> Roonwal and Chhotani, 1962
Subfamily HETEROTERMITINAE Froggatt, 1897
Genus <i>Heterotermes</i> Froggatt, 1897
3. <i>Heterotermes indicola</i> (Wasmann, 1902)

Family TERMITIDAE Latreille, 1802 Subfamily MACROTERMITINAE Kemner, 1934 Genus <i>Odontotermes</i> Holmgren, 1910
4. <i>Odontotermes assmuthi</i> Holmgren, 1913
5. <i>Odontotermes feae</i> (Wasmann, 1896) Genus <i>Microtermes</i> Wasmann, 1902
6. <i>Microtermes obesi</i> Holmgren, 1912
7. <i>Microtermes unicolor</i> Snyder, 1933

## 15. List of Terrestrial Bugs (Hemiptera)

<b>Taxa</b>
ORDER HEMIPTERA SUBORDER AUCHENORRHYNCHA INFRAORDER FULGOROMORPHA FAMILY FLATIDAE SUBFAMILY FLATINAE TRIBE NEPHESINI SUBTRIBE CRYPTOFLATINA Genus <i>Melicharia</i> Kirkaldy, 1900
1. <i>Melicharia sinhalana</i> Kirkaldy, 1900
INFRAORDER CICADOMORPHA SUPERFAMILY CERCOPOIDEA FAMILY APHROPHORIDAE SUBFAMILY APHROPHORINAE TRIBE CLOVIINI Genus <i>Clovia</i> Stål, 1866
2. <i>Clovia conifera</i> (Walker, 1851)
3. <i>Clovia puncta</i> (Walker, 1851) TRIBE PTYELINI Genus <i>Ptyelus</i> St. Farg. & Serville, 1825
4. <i>Ptyelus inconspicuus</i> Dist. 1908
5. <i>Ptyelus declaratus</i> Melich, 1903 SUPERFAMILY MEMBRACOIDEA FAMILY MEMBRACIDAE SUBFAMILY CENTROTINAE TRIBE LEPTOCENTRINI Genus <i>Leptocentrus</i> Stål, 1866
6. <i>Leptocentrus taurus</i> Fabr, 1775 TRIBE GARGARINI Genus <i>Gargara</i> Amy. & Serv., 1843
7. <i>Gargara robusta</i> Distant, 1907
8. <i>Gargara mixta</i> Buckt., 1903 FAMILY CICADELIIDAE SUBFAMILY CICADELLINAE TRIBE CICADELLINI Genus <i>Cofana</i> Melichar, 1926
9. <i>Cofana spectra</i> (Distant, 1908) SUBFAMILY DELTOCEPHALINAE TRIBE CHIASMINI Genus <i>Exitianus</i> Ball, 1929
10. <i>Exitianus indicus</i> (Distant, 1908)

11. <i>Exitianus nanus</i> (Distant, 1975) Genus <i>Nephotettix</i> Matsumura, 1902
12. <i>Nephotettix nigropictus</i> (Stål, 1870) TRIBE PENTHIMIINI Genus <i>Penthimia</i> Germar, 1821
13. <i>Penthimia compacta</i> Walker, 1851 TRIBE HECALINI Genus <i>Hecalus</i> Stål, 1864
14. <i>Hecalus porrectus</i> (Walker, 1858) SUBFAMILY IASSINAE TRIBE KRISNINI Genus <i>Krisna</i> Kirkaldy, 1900
15. <i>Krisna strigicollis</i> (Spinola, 1852) SUBFAMILY IDIOCERINAE TRIBE IDIOCERINI Genus <i>Amritodus</i> Anufriev, 1970
16. <i>Amritodus atkinsoni</i> (Lethierry, 1889) Genus <i>Idioscopus</i> Baker, 1915
17. <i>Idioscopus nitidulus</i> (Walker, 1870) SUBORDER HETEROPTERA INFRAORDER CIMICOMORPHA SUPERFAMILY REDUVIOIDEA FAMILY REDUVIIDAE SUBFAMILY HARPACTORINAE Genus 14. <i>Scipinia</i> Stål, 1861
18. <i>Scipinia horrida</i> (Stål, 1859) INFRAORDER PENTATOMORPHA SUPERFAMILY PENTATOMOIDEA FAMILY DINIDORIDAE SUBFAMILY DINIDORINAE Genus <i>Coridius</i> Illiger, 1807
19. <i>Coridius ianus</i> (Fabricius, 1775) FAMILY PENTATOMIDAE SUBFAMILY PENATOMINAE Genus <i>Eysarcoris</i> Hahn, 1834
20. <i>Eysarcoris guttiger</i> (Thunberg, 1783)
21. <i>Eysarcoris montivagus</i> (Distant, 1902)
22. <i>Eysarcoris ventralis</i> (Westwood, 1837) Genus <i>Acrosternum</i> Fieber, 1860
23. <i>Acrosternum graminea</i> (Fabricius, 1787) FAMILY SCUTELLERIDAE Genus <i>Chrysocoris</i> Hahn, 1834
24. <i>Chrysocoris stollii</i> (Wolff., 1801) Genus <i>Fitha</i>
25. <i>Fitha ardens</i> (Walker, 1867) SUPERFAMILY COREOIDEA FAMILY ALYDIDAE SUBFAMILY MICRELYTERINAE Genus 19. <i>Leptocorisa</i> Latreille, 1829
26. <i>Leptocorisa oratoria</i> (Fabr., 1794)
27. <i>Leptocorisa varicornis</i> (Fabricius, 1803) FAMILY COREIDAE SUBFAMILY COREINAE Genus <i>Cletomorpha</i> Mayr, 1866
28. <i>Cletomorpha hastata</i> (Fabricius, 1787) Genus <i>Cletus</i> Stål, 1860

29. <i>Cletus bipunctatus</i> (Westwood, 1842)
30. <i>Cletus punctulatus</i> (Westwood, 1842)
31. <i>Cletus punctiger</i> (Dallas, 1852)
32. <i>Cletus trigonus</i> (Thunberg, 1783)
33. <i>Cletus borealis</i> (Blotze., 1934)
34. <i>Cletus pygophorus</i> Gupta & Singh, 2013 SUPERFAMILY LYGAEOIDEA FAMILY LYGAEIDAE SUBFAMILY LYGAEINAE Genus <i>Spilostethus</i> Stål, 1868
35. <i>Spilostethus hospes</i> (Fabricius, 1794)
36. <i>Spilostethus pandurus</i> <i>militaris</i> (Fabricius, 1763) Genus <i>Graptostethus</i> Stål, 1868
37. <i>Graptostethus argentatus</i> (Fabricius, 1803)
38. <i>Graptostethus quadrisignatus</i> Distant, 1879
39. <i>Graptostethus servus</i> (Fabricius, 1787) SUBFAMILY ORSILLINAE TRIBE NYSIINI Genus <i>Nysius</i> Dallas, 1852
40. <i>Nysius ceylanicus</i> (Motschulsky, 1863) FAMILY GEOCORIDAE SUBFAMILY GEOCORINAE Genus <i>Geocoris</i> Fallen, 1814
41. <i>Geocoris ochropterus</i> Fieber, 1844 FAMILY RHYPAROCHROMIDAE SUBFAMILY RHYPAROCHROMINAE TRIBE RHYPAROCHROMINI Genus <i>Dieuches</i> Dohrn, 1860
42. <i>Dieuchus insignis</i> (Distant, 1904) TRIBE MYODOCHINI Genus <i>Pseudopachybrachius</i> Malipatil, 1978
43. <i>Pseudopachybrachius guttus</i> Malipatil, 1978 Genus <i>Paraecosmetus</i> Malipatil, 1978
44. <i>Paraecosmetus pallicornis</i> (Dallas, 1852) Genus <i>Paromius</i> Fieber, 1860
45. <i>Paromius exiguous</i> (Distant, 1904)

## 16. List of Aquatic and Semi-Aquatic (Heteroptera)

Order HEMIPTERA Linnaeus, 1758
Suborder HETEROPTERA Latreille, 1810
Infraorder NEPOMORPHA Popov, 1968
Superfamily NEPOIDEA Latreille, 1802
Family NEPIDAE Latreille, 1802
Subfamily NEPINAE Latreille, 1802
Genus <i>Laccotrephes</i> Stål, 1866
1. <i>Laccotrephes griseus</i> (Guérin-Méneville, 1844)
2. <i>Laccotrephes maculatus</i> (Fabricius, 1775) Subfamily RANATRINAE Douglas & Scott, 1865 Genus <i>Ranatra</i> Fabricius, 1790
3. <i>Ranatra digitata</i> Hafiz & Pradhan, 1947
4. <i>Ranatra elongate</i> Fabricius, 1790
5. <i>Ranatra filiformis</i> Fabricius, 1790

6. <i>Ranatra varipes varipes</i> Stål, 1861 Family BELOSTOMATIDAE Leach, 1815 Subfamily BELOSTOMATINAE Leach, 1815 Genus <i>Diplonychus</i> Laporte, 1833
7. <i>Diplonychus annulatus</i> (Fabricius, 1781)
8. <i>Diplonychus molestus</i> (Dufour, 1863)
9. <i>Diplonychus rusticus</i> (Fabricius, 1781) Superfamily PLEOIDEA Fieber, 1851 Family PLEIDAE Fieber, 1851 Genus <i>Paraplea</i> Esaki & China, 1928
10. <i>Paraplea frontalis</i> (Fieber, 1844) Superfamily NOTONECTOIDEA Latreille, 1802 Family NOTONECTIDAE Latreille, 1802 Subfamily ANISOPINAE Hutchinson, 1929 Genus <i>Anisops</i> Spinola, 1837
11. <i>Anisops bouvieri</i> Kirkaldy, 1904
12. <i>Anisops breddini</i> Kirkaldy, 1901
13. <i>Anisops kuroiwae</i> Matsumura, 1915
14. <i>Anisops sardeus sardeus</i> Herrich-Shaffer, 1850
15. <i>Anisops tahitiensis</i> undblad, 1934 Subfamily NOTONECTINAE Latreille, 1802 Genus <i>Enithares</i> Spinola, 1837
16. <i>Enithares ciliate</i> (Fabricius, 1798) Genus <i>Nychia</i> Stål, 1860
17. <i>Nychia Sappho</i> Kirkaldy, 1901 Superfamily CORIXOIDEA Leach, 1815 Family CORIXIDAE Leach, 1815 Subfamily CYMATIINAE Walton, 1940 Genus <i>Cymatia</i> Flor, 1860
18. <i>Cymatia apparens</i> (Distant, 1910) Family MICRONECTIDAE Jaczewski, 1924 Genus <i>Micronecta</i> Kirkaldy, 1897 Subgenus <i>Basileonecta</i> Hutchinson, 1940
19. <i>Micronecta (Basileonecta) scutellaris scutellaris</i> (Stål, 1858) Subgenus <i>Pardanecta</i> Wrblewski, 1962
20. <i>Micronecta (Pardanecta) haliploides</i> Horváth, 1904 Subgenus <i>Sigmonecta</i> Wrblewski, 1962
21. <i>Micronecta (Sigmonecta) quadristrigata</i> Breddin, 1905 Infraorder Mesovelia GERROMORPHA Popov, 1971 Superfamily MESOVELIOIDEA Douglas & Scott, 1867 Family MESOVELIIDAE Douglas & Scott, 1867 Subfamily MESOVELIINAE Douglas & Scott, 1867 Genus Mulsant & Rey, 1852
22. <i>Mesovelia easaci</i> Jehamalar & Chandra, 2017
23. <i>Mesovelia vittigera</i> Horváth, 1895 Superfamily HYDROMETROIDEA Billberg, 1820 Family HYDROMETRIDAE Billberg, 1820 Subfamily HYDROMETRINAE Billberg, 1820 Genus <i>Hydrometra</i> Latreille, 1796
24. <i>Hydrometra greeni</i> Kirkaldy, 1898
25. <i>Hydrometra okinawana</i> Drake, 1951* Superfamily GERROIDEA Reuter, 1910 Family VELIIDAE Amyot & Serville, 1843 Division MICROVELIIFORMES Andersen, 1982 Subfamily MICROVELIINAE China & Usinger, 1949 Genus <i>Microvelia</i> Westwood, 1834

Subgenus <i>Dilutovelina</i> Zettel, 2012
26. <i>Microvelia (Dilutovelina) leveillei leveillei</i> (Lethierry, 1877)
Subgenus <i>Picaultia</i> Distant, 1913
27. <i>Microvelia (Picaultia) douglasi</i> Scott, 1874
Family GERRIDAE LEACH, 1815
Subfamily RHAGADOTARSINAE Lundblad, 1933
Genus <i>Rhagadotarsus</i> Breddin, 1905
Subgenus <i>Rhagadotarsus</i> Breddin, 1905
28. <i>Rhagadotarsus (Rhagadotarsus) kraepelini</i> Breddin, 1905
Subfamily GERRINAE Bianchi, 1896
Genus <i>Aquarius</i> Schellenberg, 1800
29. <i>Aquarius adelaidis</i> Dohrn, 1860
Genus <i>Neogerris</i> Matsumura, 1913
30. <i>Neogerris parvulus</i> (Stål, 1859)
Genus <i>Limnogonus</i> Stål, 1868
Subgenus <i>Limnogonus</i> Stål, 1868
31. <i>Limnogonus (Limnogonus) fossarum fossarum</i> (Fabricius, 1775)
32. <i>Limnogonus (Limnogonus) nitidus</i> (Mayr, 1865)

## 17. List of Ants

<b>Taxa</b>
Superfamily: Vespoidea
Family: Formicidae Latreille?, 1809
Subfamily: Dolichoderinae Forel, 1878
Genus: <i>Dolichoderus</i> Lund, 1831
1. <i>Dolichoderus thoracicus</i> (Smith, F., 1860)
Genus <i>Tapinoma</i> Foerster, 1850
2. <i>Tapinoma indicum</i> Forel, 1895
Subfamily: Formicinae Latreille, 1809
Genus: <i>Anoplolepis</i> Santschi, 1914
3. <i>Anoplolepis gracilipes</i> (Smith, F., 1857)
Genus <i>Camponotus</i> Mayr, 1861
4. <i>Camponotus compressus</i> (Fabricius, 1787)
5. <i>Camponotus dolendus</i> Forel, 1892
6. <i>Camponotus parius</i> Emery, 1889
7. <i>Camponotus sericeus</i> (Fabricius, 1798)
Genus: <i>Lepisiota</i>
8. <i>Lepisiota opaca</i> (Forel, 1892)
9. <i>Lepisiota frauenfeldi</i> (Mayr, 1855)
Genus <i>Nylanderia</i>
10. <i>Nylanderia bourbonica</i> (Forel, 1886)
11. <i>Nylanderia indica</i> (Forel, 1894)
Genus <i>Oecophylla</i>
12. <i>Oecophylla smaragdina</i> (Fabricius, 1775)
Genus <i>Paratrechina</i>
13. <i>Paratrechina longicornis</i> (Latreille, 1802)
Genus: <i>Plagiolepis</i>
14. <i>Plagiolepis jerdonii</i> Forel, 1894
Genus: <i>Polyrhachis</i>
15. <i>Polyrhachis dives</i> Smith, F., 1857
16. <i>Polyrhachis lacteipennis</i>
17. <i>Polyrhachis illaudata</i> Walker, 1859
Subfamily: Myrmicinae Lepeletier de Saint- Fargeau, 1835

Genus: <i>Aphaenogaster</i> Mayr, 1853
18. <i>Aphaenogaster feae</i> Emery, 1889
Genus: <i>Carebara</i> Westwood, 1840
19. <i>Carebara affinis</i> (Jerdon, 1851)
Genus: <i>Cardiocondyla</i> Emery, 1869
20. <i>Cardiocondyla wroughtonii</i> (Forel, 1890)
21. <i>Cardiocondyla kagutsuchi</i> Terayama, 1999
Genus <i>Crematogaster</i> Lund, 1831
22. <i>Crematogaster aberrans</i> Forel, 1892
23. <i>Crematogaster buddhae</i> Forel, 1902
24. <i>Crematogaster hodgsoni</i> Forel, 1902
25. <i>Crematogaster rogenhoferi</i> Mayr, 1879
26. <i>Crematogaster subnuda</i> Mayr, 1879
Genus <i>Meranoplus</i> Smith, F., 1853
27. <i>Meranoplus bicolor</i> (Guérin-Méneville, 1844)
28. <i>Meranoplus rothneyi</i> Forel, 1902
Genus: <i>Monomorium</i> Mayr, 1855
29. <i>Monomorium dichroum</i> Forel, 1902
30. <i>Monomorium fioricola</i> (Jerdon, 1851)
31. <i>Monomorium Monomorium</i> Bolton, 1987
32. <i>Monomorium orientale</i> Mayr, 1879
33. <i>Monomorium pharaonic</i> (Linnaeus, 1758)
Genus: <i>Pheidole</i> Westwood, 1839
34. <i>Pheidole indica</i> Mayr, 1879
35. <i>Pheidole megacephala</i> (Fabricius, 1793)
36. <i>Pheidole multidentis</i> Forel, 1902
37. <i>Pheidole pronotalis</i> Forel, 1902
38. <i>Pheidole watsoni</i> Forel, 1902
Genus: <i>Solenopsis</i> Westwood, 1840
39. <i>Solenopsis geminate</i> (Fabricius, 1804)
Genus: <i>Tetramorium</i> Mayr, 1855
40. <i>Tetramorium obesum</i> André, 1887
41. <i>Tetramorium guineense</i> (Bernard, 1953)
Genus: <i>Trichomyrmex</i> Mayr, 1865
42. <i>Trichomyrmex destructor</i> (Jerdon, 1851)
SubFamily: Ponerinae Saint-Fargeau, 1835
Genus: <i>Bothroponera</i> Mayr, 1862
43. <i>Bothroponera sulcate</i> (Mayr, 1867)
Genus: <i>Brachyponera</i> Emery, 1900
44. <i>Brachyponera jerdonii</i> (Forel, 1900)
45. <i>Brachyponera nigrita</i> Emery, 1895)
Genus: <i>Diacamma</i> Mayr, 1862
46. <i>Diacamma rugosum</i> (Le Guillou, 1842)
Genus: <i>Hypoponera</i> Santschi, 1938
47. <i>Hypoponera shattucki</i> Bharti, Akbar, Wachkoo & Singh, 2015
Genus: <i>Pseudoneoponera</i> Donisthorpe, 1943
48. <i>Pseudoneoponera rufipes</i> (Jerdon, 1851)
Subfamily: Pseudomyrmecinae Smith, M.R., 1952
Genus: <i>Tetraponera</i> Smith, F., 1852
49. <i>Tetraponera nigra</i> (Jerdon, 1851)
50. <i>Tetraponera rufonigra</i> (Jerdon, 1851)



## 18. List of Hymenoptera: Vespidae

<b>Taxa</b>
Family VESPIDAE
Subfamily EUMENINAE
Genus <i>Eumenes</i> Latre Me, 1802
1. <i>Eumenesatrophicus</i> (Fabricius, 1798)
2. <i>Eumenesmacrops</i> de Saussure, 1852
Genus <i>Antodynerus</i> de Saussure, 1855
3. <i>Antodynerus flavescens flavescens</i> (Fabricius, 1775)
Genus <i>Antepiponade</i> Saussure, 1855
4. <i>Antepiponabiguttata</i> (Fabricius, 1787)
Subfamily POLISTINAE
Genus <i>Polistes</i> Latreille, 1802
5. <i>Polistes (Gyrostoma) olivaceus</i> (De Geer, 1773)
Genus <i>Ropalidia</i> Guérin-Méneville, 1831
6. <i>Ropalidiabrevita</i> Das & Gupta, 1989
7. <i>Ropalidiavariegata</i> (Smith, 1852)
Subfamily VESPINAE
Genus <i>Vespa</i> Linnaeus, 1758
8. <i>Vespa orientalis</i> Linnaeus, 1771
9. <i>Vespa tropicahaematodes</i> Bequaert, 1936
10. <i>Vespa tropicatropica</i> (Linnaeus, 1758)

## 19. List of Chalcidids

<b>Taxa</b>
1. <i>Antrocephalus fascicornis</i> (Walker, 1871)
2. <i>Antrocephalus phaeospilus</i> Waterston, 1922
3. <i>Brachymeria minuta</i> (Linnaeus, 1767)
4. <i>Brachymeria podagrica</i> (Fabricius, 1787)
Genus <i>Dirhinus</i> Dalman
5. <i>Dirhinus bakeri</i> (Crawford, 1915)
Genus <i>Hockeria</i> Walker
6. <i>Hockeria gibsoni</i> Narendran, 1989
7. <i>Hockeria hayati</i> Narendran, 1989
8. <i>Hockeria manii</i> Narendran, 1989
9. <i>Hockeria opisinae</i> Narendran, 1989
Genus <i>Kriechbaumerella</i> Dalla Torre
10. <i>Kriechbaumerella pulvinata</i> (Masi, 1932)
11. <i>Kriechbaumerella rufimanus</i> (Walker, 1860)

## 20. List of Encyrtidae

1. <i>Acerophagus papaya</i> Noyes and Schauff
2. <i>Anagyrus aquilonaris</i> (Noyes and Hayat)
3. <i>Anagyrus dactylopii</i> (Howard)

4. <i>Anagyrus tricolor</i> (Girault)
5. <i>Anomalicornia tenuicornis</i> Mercet
6. <i>Cheiloneurus bengalensis</i> (Subba Rao)
7. <i>Copidosoma floridanum</i> (Ashmead)
8. <i>Indaphycus planus</i> Hayat
9. <i>Neodusmetia sangwani</i> (Subba Rao)
10. <i>Metaphycus</i> Mercet
11. <i>Ooencyrtus utetheisae</i> (Risbec, 1951)
12. <i>Prochiloneurus albifuniculus</i> (Hayat, Alam and Agarwal)
13. <i>Prochiloneurus pulchellus</i> Silvestri
14. <i>Pseudleptomastix mexicana</i> Noyes and Schauff, 2003 (introduced species)
15. <i>Rhopus</i> sp. Foerster
16. <i>Syrphophagus hofferi</i> (Hayat)
17. <i>Tassonia gloriae</i> (Girault)

## 21. List of Coleopterans: Beetles

S. No	Taxa	Habitat
	Suborder ADEPHAGA Family Gyrinidae	
1.	<i>Dineutus (Protodineutus) indicus</i> Aubé, 1838	Aquatic
2.	<i>Orectochilus productus</i> Régimbart, 1884	Aquatic
3.	Family Carabidae <i>Amblystomus</i> sp.	Predator
4.	<i>Callytron limosum</i> (Saunders, 1834)	Predator
5.	<i>Calochroa sexpunctata</i> Fabricius, 1775	Predator
6.	<i>Cicindela aurovittata</i> Brulle, 1838	Predator
7.	<i>Cylindera (Eugrapha) minuta</i> (Olivier, 1790)	Predator
8.	<i>Neocollyris bonelli</i> (Guerin, 1834)	Predator
9.	<i>Neocollyris crassicornis</i> (Dejean, 1825)	Predator
10.	<i>Pogonus (Pogonus) biroi</i> Csiki, 1907	Predator
11.	Family Haliplidae <i>Halipilus angustifrons</i> Régimbart, 1892	Aquatic
12.	<i>Canthydrus laetabilis</i> (Walker, 1858)	Aquatic
13.	Family Noteridae Family Dytiscidae <i>Cybister javanus</i> Aube, 1838	Aquatic
14.	<i>Cybister limbatus</i> (Fabricius, 1775)	Aquatic
15.	<i>Cybister tripunctatus lateralis</i> (Fabricius, 1798)	Aquatic
16.	<i>Eretes sticticus</i> (Linnaeus, 1767)	Aquatic
17.	<i>Hydaticus</i> sp.	Aquatic
18.	<i>Hydroglyphus fiammulatus</i> (Sharp, 1882)	Aquatic
19.	<i>Hydroglyphus inconstans</i> (Régimbart, 1892)	Aquatic
20.	<i>Hydrovatus acuminatus</i> Motschulsky, 1859	Aquatic
21.	<i>Hydrovatus bonvouloiri</i> Sharp, 1882	Aquatic
22.	<i>Hydrovatus confertus</i> Sharp, 1882	Aquatic
23.	<i>Laccophilus anticatus</i> Sharp, 1890	Aquatic
24.	<i>Laccophilus flexuosus</i> Aube, 1838	Aquatic
25.	<i>Laccophilus inefficiens basalis</i> Motschulsky, 1859)	Aquatic
26.	<i>Laccophilus parvulus</i> Aubé, 1838	Aquatic
27.	<i>Laccophilus sharpi</i> Suborder POLYPHAGA	Aquatic
28.	Family Hydrophilidae <i>Amphiops materpedestris</i> Sharp, 1890	Aquatic
29.	<i>Amphiops mirabilis</i> Sharp, 1890	Aquatic
30.	<i>Berosus (Enoplurus) fairmairei</i> Zaitzev, 1908	Aquatic
31.	<i>Berosus (Enoplurus) indicus</i> Motschulsky, 1861	Aquatic
32.	<i>Enochrus (Methydrus) esuriens</i> (Walker, 1858)	Aquatic
33.	<i>Helochares anchoralis</i> Sharp, 1890	Aquatic

34.	<i>Regimbertia attenuate</i> (Fabricius, 1801)	Aquatic
35.	<i>Stemolophus rufipes</i> (Fabricius, 1792)	Aquatic
36.	Family Spercheidae <i>Spercheus belli belli</i> Champion, 1919	Aquatic
37.	Family Scarabaeidae <i>Adoretus flavus</i> Arrow 1917	Chafer
38.	<i>Adoretus lacustris</i> Arrow 1917	Chafer
39.	<i>Adoretus lasiopygus</i> Burmeister, 1855	Chafer
40.	<i>Adoretus versutus</i> Harold, 1869	Chafer
40.	<i>Adoretus versutus</i> Harold, 1869	Chafer
41.	<i>Anomala bengalensis</i> (Blanchard, 1851)	Chafer
42.	<i>Anomala biharensis</i> Arrow, 1917	Chafer
43.	<i>Anomala bilobate</i> Arrow, 1912	Chafer
44.	<i>Anomala polita</i> (Blanchard, 1851)	Chafer
45.	<i>Heteronychus lioderes</i> Redtenbacher, 1867	Chafer
46.	<i>Oryctes rhinoceros</i> (Linnaeus, 1758)	Chafer
47.	<i>Xylotrupes Gideon</i> (Linnaeus, 1767)	Chafer
48.	<i>Onitis philemon</i> Fabricius, 1801	Dung beetle
49.	<i>Onthophagus (Colobonthophagus) dama</i> (Fabricius, 1798)	Dungbeetle
50.	Family Elateridae <i>Agrypnus</i> sp.	
51.	Family Dermestidae <i>Anthrenus</i> sp.	
52.	<i>Attagenus alfieri</i> Fabricius, 1787	
53.	Family Coccinellidae <i>Coccinella septempunctata</i> (Linnaeus, 1758)	Predator
54.	<i>Coccinella transversalis</i> Fabricius, 1781	Predator
55.	<i>Coelophora unicolor</i> Fabricius, 1792	Predator
56.	<i>Halyzia sanscrita</i> Mulsant, 1853	Predator
57.	<i>Menochilus sexmaculatus</i> (Fabricius, 1781)	Predator
58.	<i>Micraspis discolor</i> (Fabricius, 1798)	Predator
59.	<i>Henosepilachna</i> (Fabricius, 1775)	Chafer
60.	<i>Henosepilachna indica</i> (Mulsant 1850)	Chafer
61.	Family Tenebrionidae <i>Gonocephalum depressum</i> (Fabricius, 1801)	Chafer
62.	<i>Gonocephalum hoffmannseggii</i> (Steven, 1829)	Chafer
63.	<i>Gonocephalum stoeckleini</i> Kaszab, 1952	Chafer
64.	<i>Gonocephalum vagum planatum</i> (Walker, 1858)	Chafer
65.	<i>Scleron reitteri</i> Gebien 1906	Chafer
66.	<i>Scleropatrum strigatum</i> Fabricius, 1798	Chafer
67.	<i>Tribolium castaneum</i> (Herbst 1797)	Chafer
68.	Family Meloidae <i>Hycleus phaleratus</i> (Pallas, 1781)	Chafer
69.	Family Chrysomelidae <i>Aulacophora almora</i> Maulik, 1936	Chafer
70.	<i>Aulacophora lewisii</i> Baly, 1886	Chafer
71.	<i>Aulacophora foveicollis</i> Lucas, 1849	Chafer
72.	<i>Cassida circumdata</i> Herbst, 1799	Chafer
73.	<i>Cassida enervis</i> Boheman, 1862	Chafer
74.	<i>Lema (Lema)</i> (Fabricius, 1798)	Chafer
75.	<i>Lema (Lema) lacertosa</i> Lacordaire, 1845	Chafer
76.	<i>Monolepta signata</i> (Olivier, 1808)	Chafer
77.	<i>Trichobalya bowringi bowringi</i> Baly, 1890	Chafer

## 22. List of Butterflies

<b>Family HesperIIDae</b>
1. <i>Ampittia dioscorides</i> (Fabricius)
2. <i>Barbo cinnara</i> (Wallace, 1866)
3. <i>Lambrix salsala</i> (Moore, 1866)
4. <i>Matapa aria</i> (Moore, 1866)

5. <i>Oriens gola</i> (Moore, 1877)
6. <i>Oriens goloides</i> (Moore, 1881)
7. <i>Pelopidas agna</i> (Moore, 1866)
8. <i>Pelopidas mathias</i> (Fabricius, 1798)
9. <i>Suastus gremius</i> (Fabricius, 1798)
10. <i>Tagiades japedus</i> (Stoll, 1781)
11. <i>Telicota ancilla</i> (Herrich-Schaffer, 1869)
12. <i>Udaspes folus</i> (Cramer, 1775)
<b>Family Lycaenidae</b>
13. <i>Anthene emolus</i> (Godart, 1824)
14. <i>Castalius rosimon</i> (Fabricius, 1775)
15. <i>Catochrysops strabo</i> (Fabricius, 1793)
16. <i>Chilades lajus</i> (Stoll, 1780)
17. <i>Curetis thetis</i> (Drury, 1773)
18. <i>Euchrysops cnejus</i> (Fabricius, 1798)
19. <i>Euchrysops pandava</i> (Horsfield, 1829)
20. <i>Jamides celeno</i> (Cramer, 1775)
21. <i>Loxura atymnus</i> (Stoll, 1780)
22. <i>Neopithecops zalmora</i> (Butler, 1870)
23. <i>Prosotas nora</i> (C. & R. Felder, 1860)
24. <i>Pseudozizeeria maha</i> (Kollar, 1848)
25. <i>Rapala manea</i> (Hewitson, 1863)
26. <i>Rathinda amor</i> (Fabricius, 1775)
27. <i>Spindasis vulcanus</i> (Fabricius, 1775)
28. <i>Tarucus nara</i> (Kollar, 1848)
29. <i>Zizeeria karsandra</i> (Moore, 1865)
30. <i>Zizula hylax</i> (Fabricius, 1775)
<b>Family Nymphalidae</b>
31. <i>Acraea violae</i> (Fabricius, 1775)
32. <i>Ariadne ariadne</i> (Linnaeus, 1758)
33. <i>Ariadne merione</i> (Cramer, 1779)
34. <i>Charaxes solon</i> (Fabricius, 1793)
35. <i>Danaus chrysippus</i> (Linnaeus, 1758)
36. <i>Danaus genutia</i> (Cramer, 1779)
37. <i>Elymnias hypermnestra</i> (Linnaeus, 1758)
38. <i>Euploea core</i> (Cramer, 1780)
39. <i>Euploea klugii</i> (Moore, 1858)
40. <i>Euthalea aconthea</i> (Cramer, 1777)
41. <i>Hypolimnas bolina</i> (Linnaeus, 1758)
42. <i>Hypolimnas misippus</i> (Linnaeus, 1758)
43. <i>Junonia almana</i> (Linnaeus, 1758)
44. <i>Junonia atlites</i> (Linnaeus, 1763)
45. <i>Junonia lemonias</i> (Linnaeus, 1758)
46. <i>Melanitis leda</i> (Linnaeus, 1758)
47. <i>Moduza procris</i> (Cramer, 1777)
48. <i>Mycalesis mineus</i> (Linnaeus, 1758)
49. <i>Mycalesis perseus</i> (Fabricius, 1775)
50. <i>Neptis hylas</i> (Moore, 1758)
51. <i>Neptis jumbah</i> (Moore, 1857)
52. <i>Phalanta phalanta</i> (Drury, 1773)
53. <i>Tirumala limniace</i> (Cramer, 1775)
54. <i>Ypthima baldus</i> (Fabricius, 1775)
55. <i>Ypthima huebneri</i> (Kirby, 1871)
<b>Family Papilionidae</b>
56. <i>Graphium agamemmon</i> (Linnaeus, 1758)

57. <i>Graphium doson</i> (C. & R. Felder, 1864)
58. <i>Pachliopta aristolochiae</i> (Fabricius, 1775)
59. <i>Pachliopta hector</i> (Linnaeus, 1758)
60. <i>Papilio clytia</i> (Linnaeus, 1758)
61. <i>Papilio demoleus</i> (Linnaeus, 1758)
62. <i>Papilio polymnestor</i> (Cramer, 1775)
63. <i>Papilio polytes</i> (Linnaeus, 1758)
<b>Family Pieridae</b>
64. <i>Anaphaeis aurota</i> (Fabricius, 1775)
65. <i>Appias albina</i> (Boisduval, 1836)
66. <i>Appias libythea</i> (Fabricius, 1775)
67. <i>Catopsilia pomona</i> (Fabricius, 1775)
68. <i>Catopsilia pyranthe</i> (Linnaeus, 1758)
69. <i>Cepora nerissa</i> (Fabricius, 1775)
70. <i>Delias eucharis</i> (Drury, 1773)
71. <i>Eurema blanda</i> (Boisduval, 1836)
72. <i>Eurema hecabe</i> (Linnaeus, 1758)
73. <i>Leptosia nina</i> (Fabricius, 1793)
74. <i>Pareronia valeria</i> (Cramer, 1776)
<b>Family Riodinidae</b>
75. <i>Abisara echerius</i> (Stoll, 1790)

## 23. List of Moths

### Family Zygaenidae

1. *Balataea postvitta* Moore, 1879
2. *Gynautocera papilionaria* Guérin- Méneville, 1831
3. *Inope fuliginosa* Moore, 1879
4. *Thyrassia subcordata* Walker, 1854
5. *Trypanophora semihyalina* Kollar, 1844

### Family Saturniidae

6. *Actias selene* Hübner, 1806
7. *Antheraea paphia* (Linnaeus)
8. *Cricula trifenestrata* (Helfer)

### Family Geometridae

9. *Agathia lycaenaria* Kollar, 1844
10. *Ascotis* sp.
11. *Epipristis minimaria* Guenee, 1857
12. *Gelasma goniaria* Felder, 1875
13. *Metoeca foedalis*

14. *Pelagodes* sp.
15. *Problepsis vulgaris* Butler, 1889
16. *Scopula actuaria* Walker, 1861
17. *Scopula aspilataria* Walker, 1861
18. *Scopula pulchellata* Fabricius, 1794
19. *Scopula remotata* Guenee, 1858
20. *Scopula* sp.
21. *Scopula emissaria*
22. *Semiothisa pluviata* Fabricius, 1795
23. *Thalassodes quadraria* (Guenee, 1857)
24. *Timandra convectaria* Walker, 1861
25. *Traminda* sp.

#### **Family Sphingidae**

26. *Acherontia lachesis* Fabricius 1798
27. *Acherontia styx* Westwood 1848
28. *Acosmeryx naga* Moore 1857
29. *Agrius convolvuli* Linnaeus 1758
30. *Ambulyx substrigilis* Westwood 1848
31. *Callambulyx rubricosa* Walker 1856
32. *Cephonodes shylas* Linnaeus 1771
33. *Daphnis hypothous* Cramer 1780
34. *Daphnis nerii* Linnaeus 1758
35. *Eupana cramydon*
36. *Gurelca hyas*
37. *Leucophlebia lineata* Westwood 1847
38. *Macroglossum assimilis* Swainson 1821
39. *Macroglossum glaucoptera*
40. *Meganoton analis* C. Felder & R. Felder 1874
41. *Nephele didyma*
42. *Psilogramma menephron* Cramer 1780
43. *Sataspes infernalis* Westwood 1848

44. *Theretra alecto alecto* Linnaeus 1758  
 45. *Theretra clotho clotho* Drury 1773  
 46. *Theretra nesus* Drury 1773  
 47. *Theretra pinastrina* Martyn, 1797  
 48. *Pergesa acteus*

**Family Lasiocampidae**

49. *Dendrolimus lattipenis* Walker, 1855  
 50. *Metanastria hyrtaca* Cramer, 1782  
 51. *Streblote dorsalis* Walker, 1866  
 52. *Streblote siva* Lefebvre, 1827  
 53. *Trabala visnou* Lefebvre, 1827

**Family Erebidae**

54. *Vamuna ramelana* (Moore, 1900)  
 55. *Mocis frugalis* (Fabricius, 1775)  
 56. *Amata bicincta* (Kollar, 1844)  
 57. *Amata cyssea* (Cramer, 1782)  
 58. *Amata hydatina* (Butler, 1876)  
 59. *Amata insueta* (Swinhoe, 1892)  
 60. *Amata lucina* (Butler, 1876)  
 61. *Amata passalis* (Fabricius, 1781)  
 62. *Amata fortunei* d'Orza, 1869  
 63. *Arctornis submarginata* Walker, 1855  
 64. *Aroa plana* Moore, 1879  
 65. *Aroa socrus* Geyer, 1837  
 66. *Miltochrista obsoleta* (Moore, 1878)  
 67. *Miltochrista congerens* (Felder, 1874)  
 68. *Caeneressa diaphana* (Kollar, 1844)  
 69. *Syntomoides imaon* (Butler, 1879)  
 70. *Chrysorabdia viridata* (Walker, 1865)  
 71. *Creatonotos gangis* Linnaeus, 1763  
 72. *Eilema vicaria* (Walker, 1854)

73. <i>Eressa con nis</i> (Walker, 1854)
74. <i>Euproctis ampala</i> Walker, 1855
75. <i>Euproctis asoetria</i> Huebner, 1818
76. <i>Euproctis digramma</i> Guerin, 1829
77. <i>Euproctis rhoda</i> (Moore, 1879)
78. <i>Euproctis lunata</i> Walker, 1855
79. <i>Euproctis plana</i> Fawcett, 1915
80. <i>Euproctis similis</i> Moore, 1879
81. <i>Euproctis subfasciata</i> Walker, 1865
82. <i>Euproctis annulata</i> Hampson, 1900
83. <i>Lymantriades varians</i> Walker, 1855
84. <i>Barsine radians</i> Moore, 1878
85. <i>Narosodes punctata</i> Walker, 1863
86. <i>Nishada abifera</i> Moore, 1878
87. <i>Orgyia australis</i> Walker, 1855
88. <i>Pantana visum</i> Huebner, 1825
89. <i>Olepa ricini</i> (Fabricius, 1794)
90. <i>Perina nuda</i> Fabricius, 1787
91. <i>Psichotoe duvauceli</i> Boisduval, 1829
92. <i>Somena scintillans</i> Walker, 1856
93. <i>Tigrioides fulveola</i> Hampson, 1900
94. <i>Ranghana punctata</i> Moore, 1878
95. <i>Thyas honesta</i> Huebner, 1806
96. <i>Eudocima</i> sp.
97. <i>Ophiusa coronata</i> Fabricius, 1775
98. <i>Othreis</i> sp.
99. <i>Artena dotata</i> (Fabricius, 1794)
100. <i>Asota ficus</i> Cramer, 1775
101. <i>Asota caricae</i> Fabricius, 1775
<b>Family Noctuidae</b>
102. <i>Chrysodeixis eriosoma</i> Doubleday, 1843



103. *Pseudaletia unipuncta* Haworth, 1809

104. *Spodoptera litura* Fabricius, 1775

#### **Family Crambidae**

105. *Aethaloessa floridalis* Zeller, 1852

106. *Agathodes ostentalis* Geyer, 1837

107. *Agrotera basinotata* Hampson, 1891

108. *Analyta melanopalis* Guenee, 1854

109. *Analyta sigulalis* Guenee, 1854

110. *Ancylolomia chrysographella* (Kollar, 1844)

111. *Antigastra catalaunalis* Duponchel, 1833

112. *Archernis tropicalis* Walker, 1859

113. *Arthroschista hilaralis* Walker, 1859

114. *Autocharis fessalis* Swinhoe, 1887

115. *Bocchoris acamasalis* Walker, 1859

116. *Botyodes asialis* Guenee, 1854

117. *Botyodes flavibasalis* Moore, 1867

118. *Bradina admixtalis* Walker, 1859

119. *Ceratarcha umbrosa* Swinhoe, 1894

120. *Chilo* sp.

121. *Cirrhochrista brizotalis* Walker, 1859

122. *Cnaphalocrocis medinalis* Guenee, 1854

123. *Cnaphalocrocis pauperalis* Strand, 1918

124. *Cnaphalocrocis trapezalis* Guenee, 1854

125. *Cnaphalocrocis trebiusalis* Walker, 1859

126. *Cnaphalocrocis venialialis* Walker, 1859

127. *Crocidolomia binotalis* Zeller, 1852

128. *Cryptographis indica* Saunders, 1891

129. *Cydalima laticostalis* Guenee, 1854

130. *Dichocrocis evaxalis* Walker, 1859

131. *Dysallacta negatalis* Walker, 1859

132. *Ercta elutalis* Walker, 1859

133. *Euclasta defamatalis* Walker, 1859
134. *Eurrhyarodes tricoloralis* Zeller, 1852
135. *Eurrhyarodes bracteolalis* Zeller, 1852
136. *Glyphodes bicolor* Swainson, 1821
137. *Glyphodes bivitalis* Guenee, 1854
138. *Glyphodes caesalis* Walker, 1859
139. *Glyphodes canthusalis* Walker, 1859
140. *Glyphodes pyloalis* Walker, 1859
141. *Glyphodes* sp.
142. *Glyphodes stolalis* Guenee, 1854
143. *Herpetogramma licarsialis* Walker, 1859
144. *Herpetogramma* sp.
145. *Hyalobathra filalis* Guenee, 1854
146. *Hydriris ornatalis* Duponchel, 1832
147. *Hydrorybina bicolor* Moore, 1888
148. *Hymenoptychis sordida* Zeller, 1852
149. *Lepyrodes geometralis* (Guenee, 1854)
150. *Leucinodes apicalis* Hampson, 1896
151. *Leucinodes orbonalis* Guenee, 1854
152. *Loxoneptera albicostalis* Swinhoe, 1906
153. *Maruca testulalis* Geyer, 1832
154. *Maruca vitrata* Fabricius, 1787
155. *Mimudea leucanalisis* Swinhoe, 1890
156. *Nausinoe perspectata* (Fabricius, 1775)
157. *Noorda blitealis* Walker, 1859
158. *Notarcha quaternalis* Zeller, 1852
159. *Palpita vitrealis* (Rossi, 1794)
160. *Paraponyx crisonalis* Walker, 1859
161. *Paraponyx depunctalis* Guenee, 1854
162. *Paraponyx fluctuosalis* Walker, 1852
163. *Parotis marginata* Hampson, 1893

164. *Parotis marinata* (Fabricius 1794)
165. *Parotis vertumnalis* Guenee, 1854
166. *Pleuroptya balteata* (Fabricius 1798)
167. *Polygrammodes sabelialis* Guenee, 1854
168. *Pycnarmon caberalis* Guenee, 1854
169. *Pycnarmon meritalis* Walker, 1859
170. *Pycnarmon virgatalis* Moore 1867
171. *Pygospila tyres* Cramer, 1780
172. *Pyrousta incoloralis* Guenee, 1854
173. *Ramila marginella* Moore 1868
174. *Sameodes cancellalis* Zeller, 1852
175. *Schoenobius adjurellus* Walker, 1863
176. *Schoenobius bipunctifer* Walker, 1863
177. *Schoenobius immeritalis* Walker, 1859
178. *Schoenobius incertellus* Walker, 1863
179. *Scirpophaga auriflua* Zeller, 1863
180. *Scirpophaga incertulus* Walker, 1863
181. *Syllepte lunalis* Guenee, 1854
182. *Syllepte derogata* Fabricius, 1775
183. *Synclera trauducalis* Zeller, 1852
184. *Spoladea recurvalis* (Fabricius, 1775)
185. *Talanga sexpunctalis* Moore, 1877
186. *Tatobotys varanesalis* Walker, 1858
187. *Terastia meticulosalis* Guenee, 1854
188. *Tetridia caletoralis* Walker, 1859

**Family Elaschitidae**

189. *Thudaca obliquella* Walker, 1856

**Family Pyralidae**

190. *Euzophera particella* Ragonot, 1888
191. *Galleria mellonella* Linnaeus, 1758
192. *Herculia nigrivitta* Walker, 1863

193. <i>Herculia suffusalis</i> [Walker, 1866]
194. <i>Hypsopygia mauritalis</i> Boisduval, 1833
195. <i>Oligochroa leucophaeella</i> Zeller, 1863
196. <i>Orthopygia igniflualis</i> Walker, 1859
197. <i>Phycita hemixanthella</i> Hampson, 1856
198. <i>Pyralis manihotalis</i> Guenee, 1854
199. <i>Pyralis pictalis</i> Curtis, 1834
200. <i>Tamaraca torridalis</i> Lederer, 1834
201. <i>Thylacoptila paurosema</i> Meyrick, 1885
<b>Family Pterophoridae</b>
202. <i>Hellinsia homodactyla</i>
<b>Family Uraniidae</b>
203. <i>Phazaca</i> sp.
<b>Family Scythrididae</b>
204. <i>Eretmocera</i> sp.
<b>Family Stathmopodidae</b>
205. <i>Atkinsonia</i> sp.

## 24. List of Dipterans: True Flies

<b>Taxa</b>
Order DIPTERA
Suborder NEMATOCERA
Family LIMONIIDAE Speiser, 1909
Genus <i>Limonia</i> Meigen, 1803
Subgenus <i>Limnobia</i> Meigen, 1818
1. <i>Limonia (Limnobia) irrorata</i> (Wiedemann, 1828)
Subgenus <i>Geranomyia</i> Haliday, 1833
2. <i>Limonia (Geranomyia) flavicosta</i> (Brunetti, 1912)

Family PSYCHODIDAE Bigot, 1854

Genus *Phlebotomus* Loew, 1845

3. *Phlebotomus montana* Sinton, 1924

Subgenus *Euphlebotomus* Theodor, 1948

4. *Phlebotomus (Euphlebotomus) argentipes* (Annandale, 1908)

Genus *Psychodata* Steyskal, 1974

5. *Psychodatavittata* (Brunetti, 1908)

Family CERATOPOGONIDAE Grassi, 1900

Genus *Culicoides* Latreille, 1809

Subgenus *Avaritia* Fox, 1955

6. *Culicoides (Avaritia) actoni* Smith, 1929

7. *Culicoides (Avaritia) orientalis* Macfie, 1932

Subgenus *Hoffmania* Fox, 1948

8. *Culicoides (Hoffmania) innoxius* Sen and Das Gupta, 1959

9. *Culicoides (Hoffmania) peregrinus* Kieffer, 1910

Subgenus *Oecacta* Poey, 1853

10. *Culicoides (Oecacta) schultzei* Enderlein, 1908

Subgenus *Trithecoides* Wirth & Hubert, 1959

11. *Culicoides (Trithecoides) anophelis* Edwards, 1922

Family CHIRONOMIDAE Erichson, 1841

Genus *Clinotanypus* Kieffer, 1913

12. *Clinotanypusvomerus* (Chaudhuri & Debnath), 1984  
Family CULICIDAE Stephens, 1829  
Genus *Anopheles* Meigen, 1818  
Subgenus *Cellia* Theobald, 1903
13. *Anopheles (Cellia) stephensi* Liston, 1901
14. *Anopheles (Cellia) culicifacies* Giles, 1901  
Genus *Culex* Linnaeus, 1758  
Subgenus *Culex* Linnaeus, 1758
15. *Culex (Culex) quinquefasciatus* Say, 1823  
Genus *Aedes* Meigen, 1818  
Subgenus *Stegomyia* Theobald, 1901
16. *Aedes (Stegomyia) aegypti* (Linnaeus, 1762)
17. *Aedes (Stegomyia) albopictus* (Skuse, 1894)  
Genus *Toxorhynchites* Theobald, 1901  
Subgenus *Toxorhynchites* Theobald, 1901
18. *Toxorhynchites (Toxorhynchites) splendens* (Wiedemann, 1819)  
Suborder BRACHYCERA  
Family *Tabanidae* Samouelle, 1819  
Genus *Chrysops* Meigen, 1803
19. *Chrysopsdispar* (Fabricius, 1798)  
Genus *Tabanus* Linnaeus, 1758
20. *Tabanusdorsiger* Wiedemann, 1821  
Subgenus *Tabanus* Linnaeus, 1758

21. *Tabanus (Tabanus) striatus* Fabricius, 1787

22. *Tabanus (Tabanus) rubidus* Wiedemann, 1821

23. *Tabanus (Tabanus) brunnipennis* Ricardo, 1911

Family STRATIOMYIDAE Giebel, 1856

Genus *Hermetia* Latreille, 1804

24. *Hermetia illucens* (Linnaeus, 1758)

Genus *Microchrysa* Loew, 1855

25. *Microchrysa flaviventris* (Wiedemann, 1824)

Family ASILIDAE Kirby & Spence, 1817

Genus *Philodicus* Loew, 1848

26. *Philodicus femoralis* Ricardo, 1921

27. *Philodicus ceylanicus* Schiner, 1868

Genus *Astochia* Becker, 1913

28. *Astochia guptai* Joseph & Parui, 1981

Family DOLICHOPODIDAE Agassiz, 1846

Genus *Chrysosoma* Guerin, 1831

29. *Chrysosoma leucopogon* (Wiedemann, 1824)

Family PHORIDAE Curtis, 1833

Genus *Megaselia* Rondani, 1856

Subgenus *Megaselia* Rondani, 1856

30. *Megaselia (Megaselia) scalaris* (Loew, 1866)

Family SYRPHIDAE Rondani, 1856

Genus *Sphaerophoria* Le Peletier & Audinet-Serville, 1828

31. *Sphaerophoriaindiana* Bigot, 1884

Genus *Mesembrius* Rondani, 1857

32. *Mesembriusbengalensis* (Wiedemann, 1819)

33. *Mesembriusquadrivittatus* (Wiedemann, 1819)

Genus *Mik*, 1897

Subgenus *Mik*, 1897

34. *Eristalinus (Eristalinus) arvorum* (Fabricius, 1787)

35. *Eristalinus (Eristalinus) polychromata* (Brunetti, 1923)

36. *Eristalinus (Eristalinus) quinquestriatus* (Fabricius, 1794)

Genus *Episyrphus* Matsumura & Adachi, 1917

Subgenus *Episyrphus* Matsumura & Adachi, 1917

37. *Episyrphus (Episyrphus) balteatus* (De Geer, 1776)

Genus *Eristalis* Latreille, 1804

Subgenus *Eristalis* Latreille, 1804

38. *Eristalis (Eristalis) tenax* (Linnaeus, 1758)

Genus *Paragus* Latreille, 1804

Subgenus *Paragus* Latreille, 1804

39. *Paragus (Paragus) serratus* (Fabricius, 1805)

Genus *Syritta* Le Peletier & Audinet-Serville, 1828



40. *Syrittaindica* (Wiedemann, 1824)

Family SEPSIDAE Walker, 1833

Genus *Sepsis* Fallén, 1810

Subgenus *Allosepsis* Ozerov, 1992

41. *Sepsis (Allosepsis) indica* Wiedemann, 1824

Family TEPHRITIDAE Newman, 1834

Genus *Bactrocera* Macquart, 1835

Subgenus *Bactrocera* Macquart, 1835

42. *Bactrocera (Bactrocera) dorsalis* (Hendel, 1912)

Subgenus *Zeugodacus* Hendel, 1927

43. *Bactrocera (Zeugodacus) cucurbitae* (Coquillett, 1899)

Genus *Campiglossa* Rondani, 1870

44. *Campiglossacribellata* Bezzi, 1913

Family SCIOMYZIDAE Fallén 1820

Genus *Sepedon* Latreille, 1804

45. *Sepedonplumbella* Wiedemann, 1830

Family DROSOPHILIDAE Loew, 1862

Genus *Drosophila* Fallén, 1823

46. *Drosophilaananassae* Doleschall, 1858

Subgenus *Sophophora* Sturtevant, 1939

47. *Drosophila (Sophophora) melanogaster* (Meigen 1830)

Family HIPPOBOSCIDAE Samouelle, 1819

Genus *Hippobosca* Linnaeus, 1758

48. *Hippoboscavariegata* Megerle, 1803

Family MUSCIDAE Latreille, 1802

Genus *Atherigona* Rondani, 1856

Subgenus *Atherigona* Rondani, 1856

49. *Atherigona (Atherigona) simplex* (Thomas, 1869)

Genus *Musca* Linnaeus, 1758

Subgenus *Musca* Linnaeus, 1758

50. *Musca (Musca) domestica* Linnaeus, 1758

Subgenus *Byomya* Robineau-Desvoidy, 1830

51. *Musca (Byomya) ventrosa* Wiedemann, 1830

Genus *Neomyia* Walker, 1859

52. *Neomyialauta* (Wiedemann, 1830)

53. *timorensis* (Robineau-Desvoidy, 1830)

54. *Neomyiaindica* (Robineau-Desvoidy, 1830)

Family SARCOPHAGIDAE Macquart, 1834

Genus *Sarcophaga* Meigen 1826

55. *Sarcophagaruficornis* Macquart, 1851

Subgenus *Parasarcophaga* Johnston & Tiegs, 1921

56. *Sarcophaga (Parasarcophaga) albiceps* Meigen, 1826

Subgenus *Liosarcophaga* Enderlein, 1928

57. *Sarcophaga (Liosarcophaga) dux* Thomson, 1869

Subgenus *Iranihindia* Rohdendorf, 1961

58. *Sarcophaga (Iranihindia) futilis* Senior-White, 1924

Family RHINIIDAE Brauer and Bergenstamm, 1889

Genus *Stomorphina* Rondani, 1861

59. *Stomorphinadiscolor* (Fabricius, 1794)

Genus *Isomyia* Walker, 1859

60. *Isomyiaviridaurea* (Wiedemann, 1819)

Family CALLIPHORIDAE Townsend, 1915

Genus *Chrysomya* Robineau-Desvoidy, 1830

61. *Chrysomyamegacephala* (Fabricius, 1794)

Genus *Hemipyrellia* Townsend, 1918

62. *Hemipyrellialigurriens* (Wiedemann, 1830)

Genus *Lucilia* Robineau-Desvoidy, 1830

63. *Luciliaporphyrina* (Walker, 1856)

Genus *Bengalia* Robineau-Desvoidy, 1830

64. *Bengaliatorosa* (Wiedemann, 1819)

## 25. List of Gastropods and bivalves

<b>Taxa</b>
Class: GASTROPODA Cuvier, 1795
Subclass: CAENOGASTROPODA Cox, 1960
Order: ARCHITAENIOGLOSSA Haller, 1890
Superfamily: VIVIPAROIDEA Gray, 1847
Family: VIVIPARIDAE Gray, 1847
Subfamily: BELLAMYINAE Rohrbach, 1937
Genus: <i>Filopaludina</i> Habe, 1964
1. <i>Filopaludina bengalensis</i> (Lamarck, 1822)
2. <i>Filopaludina bengalensis</i> (f. <i>doliaris</i> ) (Annandale & Sewell, 1921)
3. <i>Filopaludina bengalensis</i> (f. <i>eburnea</i> ) (Annandale & Sewell, 1921)
Genus: <i>Idiopoma</i> Pilsbry, 1901
4. <i>Idiopoma dissimilis</i> (O. F. Müller, 1774)

<p>Family: AMPULLARIIDAE Gray, 1824  Genus: <i>Pila</i> Röding, 1798  5. <i>Pila globosa</i> (Swainson, 1822)  Superfamily: CERITHIOIDEA J. Fleming, 1822  Family: THIARIDAE Gill, 1871 (1823)  Subfamily: THIARINAE Gill, 1871 (1823)  Genus: <i>Melanoides</i> Olivier, 1804</p>
<p>6. <i>Melanoides tuberculata</i> (O. F. Müller, 1774)  Genus <i>Mieniplotia</i> Low &amp; Tan, 2014</p>
<p>7. <i>Mieniplotia scabra</i> (O. F. Müller, 1774)  Genus: <i>Tarebia</i> H. Adams &amp; A. Adams, 1854</p>
<p>8. <i>Tarebia granifera</i> (Lamarck, 1816)</p>
<p>9. <i>Tarebia lineata</i> (Gray, 1828)  Family: PACHYCHILIDAE P.  Genus: <i>Brotia</i> H. Adams, 1866</p>
<p>10. <i>Brotia costula</i> (Rafinesque, 1833)  Order: LITTORINIMORPHA Golikov &amp; Starobogatov, 1975  Superfamily: TRUNCATELLOIDEA G  Family: BITHYNIIDAE Gray, 1857  Genus: <i>Bithynia</i> Leach, 1818</p>
<p>11. <i>Bithynia pulchella</i> (Benson, 1836)  Genus: <i>Gabbia</i> Tryon, 1865</p>
<p>12. <i>Gabbia orcula</i> (Frauenfeld, 1862)</p>
<p>13. <i>Gabbia orcula</i> (var. <i>producta</i>) (Nevill, 1884)  Subclass: HETEROBRANCHIA  Superorder: HYGROPHILA  Superfamily: LYMNAEOIDEA  Family: LYMNAEIDAE  Subfamily: AMPHIPEPLEINAE  Genus: <i>Radix</i> Montfort, 1810</p>
<p>14. <i>Radix rufescens</i> (Gray, 1822)  Genus: <i>Racesina</i> Vinarski &amp; Bolotov, 2018</p>
<p>15. <i>Racesina luteola</i> (Lamarck, 1822)  Superfamily: PLANORBOIDEA  Family: PHYSIDAE Fitzinger, 1833  Subfamily: PHYSINAE  Tribe: PHYSELLINI Taylor, 2003  Genus: <i>Physella</i> Haldeman, 1842</p>
<p>16. <i>Physella acuta</i> (Draparnaud, 1805)  Family: PLANORBIDAE Rafinesque, 1815  Subfamily: PLANORBINAE  Rafinesque, 1815  Genus: <i>Gyraulus</i> Charpentier, 1837</p>
<p>17. <i>Gyraulus barrackporensis</i> (Clessin, 1886)</p>
<p>18. <i>Gyraulus convexiusculus</i> (Hutton, 1849)</p>
<p>19. <i>Gyraulus euphraticus</i> (Mousson, 1874)  Tribe: SEGMENTININI F.C. Baker, 1945  Genus: <i>Polypylis</i> Pilsbry, 1906</p>
<p>20. <i>Polypylis calathus</i> (Benson, 1850)  Genus: <i>Trochorbis</i> Benson, 1855</p>
<p>21. <i>Trochorbis trochoideus</i> (Benson, 1836)  Family: BULINIDAE P. Fischer &amp; Crosse, 1880  Subfamily: BULININAE P. Fischer &amp; Crosse, 1880  Genus: <i>Indoplanorbis</i> Annandale and Prashad, 1921</p>
<p>22. <i>Indoplanorbis exustus</i> (Deshayes, 1833)</p>

Class: BIVALVIA Linnaeus, 1758
Subclass: PALAEOHETERODONTA Newell, 1965
Order: UNIONIDA Gray, 1854
Superfamily: UNIONOIDEA Rafinesque, 1820
Family: UNIONIDAE Rafinesque, 1820
Subfamily: PARREYSIINAE Henderson, 1935
Genus <i>Lamellidens</i> Simpson, 1900
23. <i>Lamellidens corrianus</i> (Lea, 1834)
24. <i>Lamellidens marginalis</i> (Lamarck, 1819)

## 26. List of Fish

S. No	Taxa	Common Name	IUCN	Remarks
Order ANGUILLIFORMES				
Family OPHICHTHIDAE				
Subfamily: Ophichthinae				
1.	<i>Pisodonophis boro</i> (Hamilton, 1822)	Rice-paddy eel	LC	Food fish
Order OSTEOGLOSSIFORMES				
Family NOTOPTERIDAE				
Subfamily: Notopterinae				
2.	<i>Chitala chitala</i> Hamilton, 1822	Humped, featherback	NT	Food fish
3.	<i>Notopterus notopterus</i> (Pallas, 1769)	Grey, featherback	LC	Food fish
Order CYPRINIFORMES				
Family CYPRINIDAE				
Subfamily: Labeoninae				
4.	<i>Cirrhinus mrigala</i> (Hamilton 1822)	Mrigal	LC	Food fish
5.	<i>Cirrhinus reba</i> (Hamilton 1822)	Reba carp	LC	Food fish
6.	<i>Labeo bata</i> (Hamilton 1822)	Bata	LC	Food fish
7.	<i>Labeo boga</i> (Hamilton 1822)	Boga labeo	LC	Food fish

8.	<i>Labeo calbasu</i> (Hamilton, 1822)	Kalbasu	LC	Food fish
9.	<i>Labeo catla</i> (Hamilton, 1822)	Catla	LC	Food fish
10.	<i>Labeo catla</i> (Hamilton, 1822)	Rohu	LC	Food fish
11.	Subfamily Smiliogastrinae <i>Pethia conchonius</i> (Hamilton, 1822)	Rosy barb	LC	Ornamental & food fish
12.	<i>Pethia phutonio</i> (Hamilton, 1822)	Spotted sail barb	LC	Ornamental & food fish
13.	<i>Pethia terio</i> (Hamilton, 1822)	One spot barb	LC	Ornamental & food fish
14.	<i>Pethia ticto</i> (Hamilton, 1822)	Two spot barb	LC	Ornamental & food fish
15.	<i>Puntius chola</i> (Hamilton, 1822)	Swamp barb	LC	Ornamental & food fish
16.	<i>Puntius sophore</i> (Hamilton, 1822)	Spotfin swamp barb	LC	Ornamental & food fish
17.	<i>Systemus sarana</i> (Hamilton, 1822)	Olive barb	LC	Food fish
18.	Subfamily: Cyprininae <i>Barbonymus gonionotus</i> (Bleeker 1849)	Silver barb	LC	Food fish, Exotic
19.	<i>Carassius auratus</i> (Linnaeus, 1758)	Gold fish	LC	Ornamental fish. Exotic
20.	<i>Cyprinus carpio</i> Linnaeus, 1758	Common carp	VU	Food fish. Exotic
21.	Family DANIONIDAE <i>Amblypharyngodon mola</i> (Hamilton, 1822)	Mola carplet	LC	Ornamental and food fish
22.	<i>Danio rerio</i> (Hamilton, 1822)	Zebra danio	LC	Ornamental fish
23.	<i>Laubuka laubuca</i> (Hamilton, 1822)	Indian Glass barb	LC	Ornamental & food fish
24.	<i>Raiamas bola</i> (Hamilton, 1822)	Trout barb	LC	Ornamental & food fish
25.	<i>Rasbora daniconius</i> (Hamilton, 1822)	Blackline rasbora	LC	Ornamental & food fish

26.	<i>Salmostoma bacaila</i> (Hamilton, 1822)	Large razorbelly	LC	Food fish
27.	<i>Salmostoma phulo</i> (Hamilton, 1822)	Finescale razorbelly minnow	LC	Food fish
28.	<i>Esomus danricus</i> (Hamilton, 1822)	Flying barb	LC	Ornamental & food fish
29.	Family XENOCYPRIDIDAE <i>Hypophthalmichthys molitrix</i>	Silver carp	NT	Food fish. Exotic
30.	<i>Hypophthalmichthys nobilis</i> (Richardson, 1845)	Bighead carp	DD	Food fish. Exotic
31.	<i>Ctenopharyngodon Idella</i> (Valenciennes, 1844)	Grass Carp	NA	Food fish. Exotic
32.	Family COBITIDAE <i>Lepidocephalichthys guntea</i> (Hamilton, 1822)	Guntea loach	LC	Widely distributed, food fish
33.	Order CHARACIFORMES Family SERRASALMIDAE <i>Piaractus brachypomus</i> (Cuvier 1818)	Red-bellied pacu	NA	Food fish. Exotic
34.	Order SILURIFORMES Family LORICARIIDAE Subfamily: Hypostominae <i>Pterygoplichthys anisitsi</i> Eigenmann & Kennedy, 1903	Snow pleco	NA	Ornamental fish. Exotic
35.	<i>Pterygoplichthys disjunctivus</i> (Weber, 1991)	Vermiculated sailfin catfish	NA	Ornamental fish. Exotic
36.	<i>Pterygoplichthys multiradiatus</i> (Hancock, 1828)	Orinoco sailfin catfish	NA	Ornamental fish. Exotic
37.	<i>Pterygoplichthys pardalis</i> (Castelnau, 1855)	Amazon sailfin catfish	NA	Ornamental fish. Exotic
38.	Family CHACIDAE <i>Chaca chaca</i> (Hamilton, 1822)	Squarehead chaca	LC	
39.	Family ALIIDAE <i>Ailia coila</i> (Hamilton, 1822)	Gangetis ailia	NT	Ornamental & food fish
40.	<i>Silonia silondia</i> (Hamilton, 1822)	Silondia vacha	LC	Food & game fish
41.	Family HORABAGRIDAE	Indian potasi	LC	Ornamental & food fish

	<i>Pachypterus atherinoides</i> (Bloch 1794)			
42.	Family BAGRIDAE <i>Mystus bleekeri</i> (Day, 1877)	Day,s Mystus	LC	Food fish
43.	<i>Mystus gulio</i> Hamilton, 1822	Long whiskered catfish	LC	Food fish
44.	<i>Mystus tengara</i> (Hamilton, 1822)	Tengara Mystus	LC	Food fish
45.	<i>Mystus vittatus</i> (Bloch 1794)	Striped dwarf catfish	LC	Food fish
46.	<i>Rita rita</i> (Hamilton, 1822)	Rita	LC	Food fish
47.	<i>Sperata aor</i> (Hamilton, 1822)	Long whiskered catfish	LC	Food fish
48.	Family PANGASIDAE <i>Pangasianodon hypophthalmus</i> (Sauvage 1878)	Striped catfish	EN	Food fish. Exotic
49.	<i>Pangasius pangasius</i> (Hamilton 1822)	Pungas	LC	Ornamental & food fish
50.	Family SILURIDAE <i>Ompok pabda</i> (Hamilton, 1822)	Pabdah	NT	Food fish
51.	Family SILURIDAE <i>Wallago attu</i> (Schneider, 1801)	Boal	NT	Food & game fish
52.	Family CLARIIDAE <i>Clarias gariepinus</i> (Burchell, 1822)		LC	Food fish. Exotic
53.	<i>Clarias magur</i> (Hamilton, 1822)	Magur	EN	Food fish
54.	Family HETEROPNEUSTIDAE <i>Heteropneustes fossilis</i> (Bloch, 1794)	Stinging catfish	LC	Food fish
55.	Order GOBIIFORMES Family GOBIIDAE Subfamily Gobiinae <i>Glossogobius giuris</i> (Hamilton, 1822)	Tank Goby	LC	Food fish
56.	Order SYNBRANCHIFORMES Family MASTACEMBELIDAE <i>Macrogathus aral</i> (Schneider, 1801)	One stripe spinyeel	LC	Food fish



57.	<i>Macrogathus pancalus</i> Hamilton, 1822	Barred spiny eel	LC	Food fish
58.	<i>Mastacembelus armatus</i> (La Cepède, 1800)	Zig zag eel	LC	Food fish
59.	Family SYNBRANCHIDAE <i>Monopterus cuchia</i> (Hamilton, 1822)	Cuchia	LC	Food fish
60.	Order ANABANTIFORMES Family ANABANTIDAE <i>Anabas testudineus</i> (Bloch, 1792)	Climbing perch	DD	Food fish
61.	<i>Anabas cobojus</i> (Hamilton, 1822)	Gangetic koi	DD	Food fish
62.	Family OSPHRONEMIDAE Subfamily Trichogastrinae <i>Trichogaster fasciata</i> Bloch & Schneider 1801	Stripled gourami		
63.	<i>Trichogaster lalius</i> (Hamilton, 1822)	Dwarf gourami	LC	Ornamental & Food fish
64.	Family CHANNIDAE <i>Channa gachua</i> (Hamilton, 1822)	Dwarf snakehead	LC	Ornamental & Food fish
65.	<i>Channa marulius</i> (Hamilton, 1822)	Giant snakehead	LC	Food fish
66.	<i>Channa punctate</i> (Bloch, 1793)	Spotted snakehead	LC	Food fish
67.	<i>Channa striata</i> (Bloch, 1793)	Banded snakehead	LC	Food fish
68.	Family Badidae <i>Badis badis</i> (Hamilton, 1822)	Badis	LC	Ornamental fish
69.	Order Pleuronectiformes Family Cynoglossidae <i>Cynoglossus cynoglossus</i> (Hamilton, 1822)	Bengal tongue sole	LC	Food fish
70.	Order Cichliformes Family Cichlidae <i>Oreochromis mossambica</i> (Peters, 1852)	Mozambique tilapia	VU	Food fish. Exotic
71.	<i>Oreochromis niloticus</i> (Linnaeus, 1758)	Nile tilapia		Food fish. Exotic
72.	Order CYPRINODONTIFORMES	Blue panchax		Biological

	Family APLOCHEILIDAE <i>Aplocheilus panchax</i> Hamilton, 1822			
73.	Order Mugiliformes Family Mugilidae <i>Chelon.parsia</i> (Hamilton, 1822)	Goldspot mullet	NA	Food fish
74.	<i>Rhinomugil corsula</i> (Hamilton, 1822)	Corsula mullet	LC	Food fish
75.	Order Tetraodontiformes Family Tetraodontidae <i>Leiodon cutcutia</i> (Hamilton, 1822)	Ocellated pufferfish	LC	Ornamental fish
76.	Order Perciformes Family Latidae <i>Lates Calcarifer</i> (Bloch 1790)	Barramundi	LC	Food fish
77.	<i>Chanda nama</i> Hamilton, 1822	Elongate glass perchlet	LC	Ornamental & Food fish
78.	<i>Parambassis lala</i> (Hamilton, 1822)	High fin glassy perchlet	NT	Ornamental & Food fish
79.	<i>Parambassis ranga</i> (Hamilton, 1822)	Indian glassy fish	LC	Ornamental & Food fish

\*NE: Not Evaluated, DD: Data Deficient, LC: Least Concern, NT: Near Threatened, VU: Vulnerable, EN: Endangered, CR: Critically Endangered, EW: Extinct in the Wild, EX:Extinct

## 27. List of Amphibians and Reptiles

S. No.	Taxa	Common Name	Local Name
<b>Lizards</b>			
1.	FAMILY: AGAMIDAE <i>Calotes versicolor</i> (Daudin, 1802) Common Garden Lizard	Girgiti	
2.	FAMILY: GEKKONIDAE <i>Hemidactylus flaviviridis</i> Ruppell, 1835 Yellow-Green House Gecko	Tiktiki	
3.	<i>Hemidactylus frenatus</i> Dumeril	Asian House Gecko	Chhoto Tiktiki
4.	<i>Hemidactylus parvimaclatus</i>	Ground Gecko	Metho Tiktiki
5.	FAMILY: SCINCIDAE <i>Eutropis carinata</i> (Schneider, 1799)	Common Grass Skink	Baro Ghaser Anjoni
6.	<i>Eutropis macularia</i> (Blyth, 1853) Bronze Grass Skink	Tamate Ghaser Anjoni	
7.	<i>Riopa albopunctata</i> (Gray, 1846) White-Spotted Supple Skink	Futkijukto choto anjoni	

8.	FAMILY: VARANIDAE <i>Varanus salvator</i> (Laurenti, 1768)	Water Monitor Lizard	Jal Godhika, Tarkel, Gorkel
9.	<i>Varanus flavescens</i> (Hardwicke & Gray, 1827)	Yellow Monitor Lizard	Sonali Godhika, Sonali Gui
<b>Turtle</b>			
10.	FAMILY: TRIONYCHIDAE <i>Lissemys punctata</i> (Bonnaterre, 1789) Indian Flap-Shell Turtle	Chiti Kachim	
<b>Snake</b>			
11.	FAMILY: COLUBRIDAE <i>Amphiesma stolatum</i> (Linnaeus, 1758)	Buff-Striped Keelback	Hele
12.	<i>Atretium schistosum</i> (Daudin, 1803) Olive Keelback Water Snake	Kerul	
13.	<i>Cerberus rynchops</i> (Schneider, 1799) Dog-Faced Water Snake	Kukur Mukho	
14.	<i>Enhydris enhydris</i> (Schneider, 1799)	Common Smooth	Metuli
15.	<i>Lycodon aulicus</i> (Linnaeus, 1758) Common Wolf Snake	Gharchiti	
16.	<i>Ptyas mucosus</i> (Linnaeus, 1758)	Indian Rat Snake	Daras, Dhamna
17.	<i>Fowlea piscator</i> (Schneider, 1799)	Checkered Keelback	Jaldhora
18.	<i>Fowlea schnurrenbergeri</i> (Kramer, 1977)	Bar-Necked Keelback	Jaldhora
19.	<i>Xenochrophis cerasogaster</i> (Cantor, 1839)	Painted Keelback	Khorichonch
20.	<i>Chrysopelea ornata</i> (Shaw, 1802)	Ornamental Flying Snake	Kalnagini
21.	<i>Oligodon arnensis</i> (Shaw, 1802)	Banded Kukri Snake	Udaykal
22.	<i>Ahaetulla anomala</i> Annandale, 1906	Common Vine Snake	Laudaga
23.	<i>Dendrelaphis tristis</i> (Daudin, 1803)	Common Bronze-Back	Betanchra
24.	<i>Dendrelaphis pictus</i> (Gmelin, 1789)	Painted Bronze-back	Kharichur
<b>FAMILY: ELAPIDAE</b>			
25.	<i>Bungarus caeruleus</i> (Schneider, 1801)	Common Krait	Kalaj
26.	<i>Bungarus fasciatus</i> (Schneider, 1801)	Banded Krait	Sankhamuthi
27.	<i>Naja kaouthia</i> Lesson, 1831 Monocellate Cobra	Keute	
<b>FAMILY: VIPERIDAE</b>			
28.	<i>Daboia russellii</i> (Shaw & Nodder, 1797)	Russell's Viper	Chandrabora
<b>FAMILY: TYPHLOPIDAE</b>			
29.	<i>Indotyphlops braminus</i> (Daudin, 1803)	Common Blind	Puye saap
<b>Amphibians</b>			
1.	<i>Duttaphrynus melanostictus</i> (Schneider, 1799)	Common Indian Toad	
2.	<i>Duttaphrynus stomaticus</i> (Lutken, 1864)	Marbled Toad	
3.	<i>Microhyla ornata</i> (Dumeril & Bibron, 1841)	Ornate Narrow-Mouthed Frog	
4.	<i>Kaloula taprobanica</i> (Parker, 1934)	Indian Painted Frog	
5.	<i>Euphlyctis cyanophlyctis</i> (Schneider, 1799)	Skittering Frog	

6.	<i>Euphlyctis hexadactylus</i> (Lesson, 1834)	Green Pond Frog	
7.	<i>Hoplobatrachus tigerinus</i> (Daudin, 1802)	Indian Bull Frog	
8.	<i>Fejervarya syhadrensis</i> (Annandale, 1919)	Syhadra Cricket Frog	
9.	<i>Hylarana tytleri</i> (Theobald, 1868)	Reed Frog	
10.	<i>Polypedates maculatus</i> (Gray, 1834)	Common Indian Tree Frog	

## 28. List of Birds

S. No.	Common Name	Species	Status
1.	Little Grebe	<i>Podiceps ruficollis</i>	R/f
2.	Grey Pelican	<i>Pelecanus philippensis</i>	lx
3.	Cormorant	<i>Phalacrocorax Carbo</i>	M/u
4.	Indian Shag	<i>Phalacrocorax</i>	R/f
5.	Little Cormorant	<i>Phalacrocorax niger</i>	R/c
6.	Darter	<i>Anhinga Rufa</i>	lx
7.	Grey Heron	<i>Ardea cinerea</i>	R/u
8.	Purple Heron	<i>Ardea purpurea</i>	R/u
9.	Pond Heron	<i>Ardeola grayii</i>	R/u
10.	Cattle Egret	<i>Bubulcus ibis</i>	
11.	Large Egret	<i>Ardea alba</i>	R/u
12.	Smaller Egret	<i>Egretta intermedia</i>	R/u
13.	Little Egret	<i>Egretta garzetta</i>	R/u
14.	Night Heron	<i>Nycticorax nycticorax</i>	
15.	Chestnut Bittern	<i>Lxobrychus</i>	
16.	Yellow Bittern	<i>Lxobrychus sinensis</i>	
17.	Black Bittern	<i>Lxobrychus</i>	
18.	Openbill Stork	<i>Anastomus oscitans</i>	
19.	Spoonbill	<i>Platalea leucorodia</i>	
20.	Greylag Goose	<i>Anser anser</i>	
21.	Bar-headed Goose	<i>Anser indicus</i>	
22.	Lesser Whistling Teal	<i>Dendrocygna javanica</i>	

23.	Ruddy Shelduck	<i>Tadorna ferruginea</i>	
24.	Common Shelduck	<i>Tadorna fadoma</i>	
25.	Pintail	<i>Anas acuta</i>	
26.	Common Teal	<i>Anas Crecca</i>	M/u
27.	Spot-billed Duck	<i>Anas Poecilorhyncha</i>	
28.	Mallard	<i>Anas platyrhynchos</i>	
29.	Gadwall	<i>Anas strepera</i>	M/f
30.	Wigeon	<i>Anas Penelope</i>	
31.	Garganey	<i>Anas querquedula</i>	
32.	Shoveler	<i>Anas clypeata</i>	
33.	Red-crested Pochard	<i>Netta rufina</i>	
34.	Common Pochard	<i>Aythya ferina</i>	
35.	White Eyed Pochard	<i>Aythya nyroca</i>	
36.	Baer's Pochard	<i>Aythya baeri</i>	
37.	Tufted Pochard	<i>Aythya fuligula</i>	M/f
38.	Cotton Teal	<i>Nettapus coromandelianus</i>	R/r
39.	Comb Duck	<i>Sakidiomis melanotos</i>	
40.	Red-breasted merganser	<i>Mergus serrator</i>	
41.	Marsh Harrier	<i>Circus aeruginosus</i>	M/u
42.	Crested Serpent Eagle S	<i>Spilornis cheela</i>	
43.	Osprey	<i>Pandion heliaetus</i>	M/r
44.	Common Bastard Quail	<i>Tumix suscicator</i>	
45.	Water Rail	<i>Rallus Aquaticus</i>	
46.	Blue-breasted Banded Rail	<i>Rallus striatus</i>	
47.	Banded Crake	<i>Rallina eurizonoides</i>	
48.	Baillon's Crake	<i>Porzana pusilla</i>	
49.	Spotted Crake	<i>Porzana porzana</i>	
50.	Ruddy Crake	<i>Porzana fusca</i>	
51.	Brown Crake	<i>Amauromis akool</i>	
52.	White-breasted waterhen	<i>Amauromis hoenicurus</i>	R/f

53.	Watercock	<i>Gallixrex cineres</i>	
54.	Moorhen	<i>Gallixrex cineres</i>	R/f
55.	Purple Moorthen	<i>Porphyrio porphyrio</i>	R/u
56.	Coot	<i>Fulica atra</i>	M/u
57.	Pheasant-tailed Jacana	<i>Hydrophasianus chirurgus</i>	
58.	Bronze-winged Jacana	<i>Metropidius indicus</i>	R/u
59.	Grey-headed Lapwing	<i>Vanellus cinereus</i>	
60.	Red-wattled Lapwing	<i>Vanellus indicus</i>	R/u
61.	Spur-winged Lapwing	<i>Vanellus spinosus</i>	
62.	Yellow-wattled Lapwing	<i>Vanellus malabaricus</i>	
63.	Grey Plover	<i>Pluvialis Squatarola</i>	
64.	Eastern Golden Plover	<i>Pluvialis dominica</i>	
65.	Ringed Plover	<i>Charadrius hiaticufa</i>	
66.	Little Ringed Plover	<i>Charadrius dubius</i>	
67.	Kentish Plover	<i>Chandrius</i>	
68.	Curlew	<i>Numenius arquata</i>	
69.	Black-tailed Godwit	<i>Limosa limosa</i>	
70.	Spotted Redshank	<i>Tringa erythropus</i>	
71.	Marsh Sandpiper	<i>Tringa stagnatilis</i>	
72.	Greenshank	<i>Tringa nebularia</i>	
73.	Blue-throated	<i>Megalaima asiatica</i>	
74.	Lineated Barbet	<i>Megalaima lineata</i>	M
75.	Coppersmith Barbet	<i>Megalaima haemacephala</i>	
76.	Indian Roller	<i>Coracias benghalensis</i>	M
77.	Common Kingfisher	<i>Alcedo atthi</i>	
78.	White-throated Kingfisher	<i>Halcyon smyrnensis</i>	
79.	Stork-billed Kingfisher	<i>Halcyon capensis</i>	
80.	Pied Kingfisher	<i>Ceryle rudis</i>	M
81.	Common Hawk-Cuckoo	<i>Hierococcyx varius</i>	
82.	Grey-bellied Cuckoo	<i>Cacomantis passerinus</i>	

83.	Asian Koel	<i>Eudynamys scolopacea</i>	
84.	Greater Coucal	<i>Centropus sinensis</i>	
85.	Rose-ringed Parakeet	<i>Psittacula krameri</i>	
86.	House Swift	<i>Apus affinis</i>	
87.	Asian Palm-Swift	<i>Cypsiurus balasiensis</i>	
88.	Rock Pigeon	<i>Columba livia</i>	
89.	Spotted Dove	<i>Streptopelia chinensis</i>	
90.	Common Snipe	<i>Gallinago gallinago</i>	
91.	Wood Sandpiper	<i>Tringa glareola</i>	M
92.	Green Sandpiper	<i>Tringa ochropus</i>	M
93.	Common Sandpiper	<i>Tringa hypoleucos</i>	M
94.	Little Stint	<i>Calidris minuta</i>	M
95.	Small Pratincole	<i>Glareola lactea</i>	M
96.	Pacific Golden Plover	<i>Pluvialis fulva</i>	M
97.	Black Kite	<i>Milvus migrans</i>	
98.	Black-headed Ibis	<i>Threskiornis melanocephalus</i>	
99.	Brown Shrike	<i>Lanius cristatus</i>	M
100.	Long-tailed Shrike	<i>Lanius schach tricolor</i>	
101.	Rufous Treepie	<i>Corvus macrorhynchos</i>	
102.	House Crow	<i>Corvus splendens</i>	
103.	Black Drongo	<i>Dicrurus leucophaeus</i>	
104.	Chestnut-tailed Starling	<i>Sturnus malabaricus</i>	
105.	Asian Pied Starling	<i>Sturnus contra</i>	
106.	Common Myna	<i>Acridotheres tristis</i>	
107.	Barn Swallow	<i>Hirundo rustica</i>	
108.	Oriental Magpie- Robin	<i>Copsychus saularis</i>	
109.	Black Redstart	<i>Phoenicurus ochruros</i>	
110.	Red-vented Bulbul	<i>Pycnonotus cafer</i>	
111.	Plain Prinia	<i>Prinia inornata</i>	
112.	Zitting Cisticola	<i>Cisticola juncidis</i>	

113.	Blyth's Reed- Warbler	<i>Acrocephalus</i>	M
114.	Clamorous Reed-Warbler	<i>Acrocephalus dumetorum</i>	M
115.	Striated Grassbird	<i>Megalurus stentoreus</i>	
116.	Common Tailorbird	<i>Orthotomus palustris</i>	
117.	Purple Sunbird	<i>Nectarinia sutorius</i>	
118.	White-browed Wagtail	<i>Motacilla asiatica</i>	
119.	White Wagtail	<i>Motacilla alba maderaspatensis</i>	M
120.	Yellow Wagtail	<i>Motacilla fiava</i>	M
121.	Grey Wagtail	<i>Motacilla cinerea</i>	M
122.	Paddyfield Pipit	<i>Anthus rufulus</i>	
123.	House Sparrow	<i>Passer domesticus</i>	

[Status: c = common; seen in very high numbers and/or on many occasions in suitable habitat. f = fairly common; seen on more than one occasion or in fairly high numbers; u = uncommon; seen on one /two occasions and / or one or two individuals; v = vagrant; lx = locally extinct; ? = unknown; R = Resident; M = Migrant; P = Partly migrant]

## 29. List of Mammals

Taxa	Occurrence	IUCN Status
Order Carnivora	Rare	LC
Family Viverridae		
1. Asian Palm Civet <i>Paradoxurus hermaphrodites</i> (Pallas, 1777)		
Family Herpestidae	Rare	LC
2. Bengal Marsh Mongoose <i>Herpestes palustris</i> Ghose, 1965		
Family Canidae	Rare	LC
3. Asiatic Jackal <i>Canis aureus</i> Linnaeus, 1758		
Order Rodentia		
Family Sciuridae	Common	LC
4. Northern Palm Squirrel <i>Funambulus pennantii</i> Wroughton, 1905		
Family Muridae	Common	LC
5. Greater Bandicoot Rat <i>Bandicota indica</i> (Bechstein, 1800)		
6. Common House Rat <i>Rattus rattus</i> (Linnaeus, 1758)	Common	LC



7. House mouse <i>Mus musculus</i> Linnaeus, 1758	Common	LC
Order Eulipotyphla	Rare	LC
Family Soricidae		
8. Asian House Shrew <i>Suncus murinus</i> (Linnaeus, 1766)		
Order Chiroptera	Common	LC
Family Pteropodidae		
9. Indian Flying Fox <i>Pteropus giganteus</i> (Brünnich, 1782)		
10. Greater Short nosed Fruit Bat <i>Cynopterus sphinx</i> (Vahl, 1797)	Common	LC
11. Lesser Short nosed Fruit Bat <i>Cynopterus brachyotis</i> (Müller, 1838)	Common	LC
12. Leschenault's Rousette <i>Rousettus leschenaultii</i> (Desmarest, 1820)	Common	LC
Family Vespertilionidae	Common	LC
13. Indian Pipistrelle <i>Pipistrellus coromandra</i> (Gray, 1838)		

\*NE: Not Evaluated, DD: Data Deficient, LC: Least Concern, NT: Near Threatened, VU: Vulnerable, EN: Endangered, CR: Critically Endangered, EW: Extinct in the Wild, EX: Extinct

### 30. List of Fungi

	Family	Species
Bracket Macrofungi	Bondarzewiaceae	<i>Amyloporus campbellii</i> (Berk.) Ryvarden
	Ganodermataceae	<i>Ganoderma australe</i> (Fr.) Pat. <i>Ganoderma curtisii</i> (Berk.) Murrill <i>Ganoderma colossus</i> (Fr.) C.F. Baker <i>Ganoderma mediosinense</i> J.D. Zhao
	Hymenochaetaceae	<i>Fuscoporia senex</i> (Nees & Mont.) Ghob. -Nejh. <i>Inonotus pachyphloeus</i> (Pat.) T. Wagner & M. Fisch.

		<i>Phellinus allardii</i> (Bres.) S. Ahmad <i>Phellinus gilvus</i> (Schwein.) Pat.
	Meripilaceae	<i>Rigidoporus lineatus</i> (Pers.) Ryvarden
	Polyporaceae	<i>Cellulariella acuta</i> (Berk.) Zmitr. & Malysheva <i>Corioloopsis occidentalis</i> (Klotzsch) Murrill <i>Earliella scabrosa</i> (Pers.) Gilb. & Ryvarden <i>Favolus grammocephalus</i> (Berk.) Imazeki <i>Funalia caperata</i> (Berk.) Zmitr. & Malysheva <i>Leiotrametes lactinea</i> (Berk.) Welti & Courtec. <i>Leiotrametes menziesii</i> (Berk.) Welti & Courtec. <i>Pycnoporus sanguineus</i> (L.) Murrill <i>Truncospora tephropora</i> (Mont.) Zmitr.
Pseudogilled Macrofungi	Schizophyllaceae	<i>Schizophyllum commune</i> Fr.
	Serpulaceae	<i>Serpula similis</i> (Berk. & Broome) Ginns
Gilled Macrofungi	Agaricaceae	<i>Agaricus</i> sp.
	Hymenogastraceae	<i>Gymnopilus purpureosquamulosus</i> Høil.
	Lyophyllaceae	<i>Termitomyces striatus</i> (Beeli) R. Heim
	Pluteaceae	<i>Pluteus chrysaegis</i> (Berk. & Broome) Petch
		<i>Volvariella volvacea</i> (Bull.) Singer
	Polyporaceae	<i>Lentinus polychrous</i> Lév.
<i>Lentinus sajor-caju</i> (Fr.) Fr.		

	Tricholomataceae	<i>Macrocybe gigantea</i> (Masseé) Pegler & Lodge
	Psathyrellaceae	<i>Coprinopsis</i> sp. <i>Psathyrella</i> sp.
Crust Macrofungi	Lachnocladiaceae	<i>Scytinostroma duriusculum</i> (Berk. & Broome) Donk
	Meripilaceae	<i>Rigidoporus vinctus</i> (Pers.) Ryvarden
	Peniophoraceae	<i>Duportella tristicula</i> (Berk. & Broome) Reinking
	Phanerochaetaceae	<i>Porostereum spadiceum</i> (Pers.) Hjortstam & Ryvarden
	Polyporaceae	<i>Grammothele fuligo</i> (Berk. & Broome) Ryvarden
Stereoid Macrofungi	Meruliaceae	<i>Stereopsis dimiticum</i> (Rehill & B.K. Bakshi)
	Stereopsidaceae	<i>Gyrodontium sacchari</i> (Spreng.) Hjortstam
Toothed Macrofungi	Coniophoraceae	<i>Gyrodontium sacchari</i> (Spreng.) Hjortstam
	Meruliaceae	<i>Flavodon avus</i> (Klotzsch) Ryvarden
Jelly Macrofungi	Auriculariaceae	<i>Auricularia auricula-judae</i> (Bull.) Quél. <i>Auricularia nigricans</i> (Sw.) Birkebak, Looney & Sánchez-García
	Dacrymycetaceae	
	Tremellaceae	
Stink horn	Phallaceae	
Carbon and Cushion Macrofungi	Hypoxylaceae	<i>Daldinia bambusicola</i> Y.M. Ju, Macrofungi J.D. Rogers & F. San Martín <i>Daldinia concentrica</i> (Bolton) Ces. & De Not. <i>Hypoxylon haematostroma</i> Mont.

	Xylariaceae	<i>Xylaria multiplex</i> (Kunze) Fr. <i>Xylaria polymorpha</i> (Pers.) Grev.
--	-------------	---

### 31. List of Freshwater Algae

S. No.	Taxa	Family	Mode of occurrence
<b>Cyanophyceae</b>			
1	<i>Limnococcus limneticus</i> (Lemmerm.) Komárk. & al.	Merismopediaceae	Planktonic
2	<i>Merismopedia minima</i> Beck	Merismopediaceae	Planktonic
3	<i>Merismopedia tenuissima</i> Lemmerm.	Merismopediaceae	Planktonic
4	<i>Merismopedia tranquilla</i> (Ehrenb.) Trevis.	Merismopediaceae	Planktonic
5	<i>Synechocystis aquatilis</i> Sauv.	Merismopediaceae	Planktonic
6	<i>Microcystis aeruginosa</i> (Kütz.) Kütz.	Microcystaceae	Planktonic
7	<i>Lemmermanniella pallida</i> (Lemmerm.) Geitler	Synechococcaceae	Planktonic
8	<i>Rhabdoderma irregulare</i> (Naumann) Geitler	Synechococcaceae	Planktonic
9	<i>Synechococcus elongatus</i> (Nägeli) Nägeli	Synechococcaceae	Planktonic
10	<i>Chroococcus disperses</i> (Keisll.) Lemmerm.	Chroococcaceae	Planktonic; Attached
11	<i>Chroococcus turgidus</i> (Kütz.) Nägeli	Chroococcaceae	Planktonic
12	<i>Pseudanabaena catenate</i> Lauterb.	Pseudanabaenaceae	Planktonic
13	<i>Leptolyngbya subtilis</i> (West) Anagn.	Leptolyngbyaceae	Attached
14	<i>Planktolyngbya limnetica</i> (Lemmerm.) Komárk. -Legn. & Cornberg	Leptolyngbyaceae	Planktonic
15	<i>Spirulina major</i> Kütz. ex Gomont	Spirulinaceae	Planktonic
16	<i>Spirulina subsalsa</i> Oerst. ex Gomont	Spirulinaceae	Planktonic
17	<i>Spirulina gigantea</i> Schmidle	Spirulinaceae	Planktonic
18	<i>Microcoleus autumnalis</i> (Gomont) Strunecky & al.	Microcoleaceae	Attached
19	<i>Planktothrix isothrix</i> (Skuja) Komárek & Komárk.	Microcoleaceae	Planktonic
20	<i>Planktothrix rubescens</i> (DC. ex Gomont) Anagn. & Komárk	Microcoleaceae	Planktonic
21	<i>Kamptonema chlorinum</i> (Kütz. ex Gomont) Strunecky & al.	Oscillatoriaceae	Attached
22	<i>Kamptonema formosum</i> (Bory ex Gomont) Strunecky & al.	Oscillatoriaceae	Attached
23	<i>Lyngbya hieronymusii</i> Lemmerm.	Oscillatoriaceae	Attached
24	<i>Oscillatoria limosa</i> C.Agardh ex Gomont	Oscillatoriaceae	Planktonic; Attached
25	<i>Oscillatoria ornata</i> Kütz. ex Gomont	Oscillatoriaceae	Attached

26	<i>Oscillatoria crassa</i> (C.B.Rao) Anagn.	Oscillatoriaceae	Attached
27	<i>Oscillatoria subbrevis</i> Schmidle	Oscillatoriaceae	Planktonic; Attached
28	<i>Oscillatoria princeps</i> Vaucher ex Gomont	Oscillatoriaceae	Attached
29	<i>Phormidium tergestinum</i> (Rabenh ex Gomont) Anagn. & Komárek	Oscillatoriaceae	Planktonic
30	<i>Phormidium chalybeum</i> (Martens ex Gomont) Anagn. & Komárek	Oscillatoriaceae	Planktonic
31	<i>Scytonema pseudohofmanni</i> Bharadwaja	Scytonemataceae	Attached
32	<i>Anabaena iyengarii</i> Bharadwaja	Nostocaceae	Planktonic
33	<i>Anabaena catenula</i> Kütz. ex Bornet & Flahault	Nostocaceae	Planktonic
34	<i>Cylindrospermum stagnale</i> Bornet & Flahault	Nostocaceae	Attached
<b>Chlorophyceae</b>			
1	<i>Chlamydomonas mucicola</i> Schmidle	Chlamydomonadaceae	Planktonic
2	<i>Edaphochlamys debaryana</i> (Gorozh.) Pröschold & Darienko	Chlamydomonadaceae	Planktonic
3	<i>Eudorina elegans</i> Ehrenb.	Volvocaceae	Planktonic
4	<i>Pandorina morum</i> (O.F. Müll.) Bory	Volvocaceae	Planktonic
5	<i>Oocystis borgei</i> J. Snow	Oocystaceae	Attached
6	<i>Chlorococcum infusionum</i> (Schrank) Menegh.	Chlorococcaceae	Attached
7	<i>Lacunastrum gracillimum</i> (West & G.S. West) McManus	Hydrodictyceae	Planktonic
8	<i>Goniochloris mutica</i> (A. Braun) Fott	Hydrodictyceae	Attached
9	<i>Pediastrum boryanum</i> var. <i>cornutum</i> (Racib.) Sulek	Hydrodictyceae	Planktonic; Attached
10	<i>Pediastrum duplex</i> Meyen	Hydrodictyceae	Planktonic
11	<i>Pediastrum sarmae</i> Keshri & Mullick	Hydrodictyceae	Planktonic
12	<i>Pediastrum subgranulatum</i> (Racib.) Komárek & Jánková	Hydrodictyceae	Attached
13	<i>Pseudopediastrum boryanum</i> (Turpin) E.Hegew.	Hydrodictyceae	Attached
14	<i>Stauridium privum</i> (Printz) E. Hegew.	Hydrodictyceae	Planktonic; Attached
15	<i>Stauridium tetras</i> (Ehrenb.) E.Hegew.	Hydrodictyceae	Planktonic; Attached
16	<i>Tetraedron caudatum</i> (Corda) Hansg.	Hydrodictyceae	Attached
17	<i>Tetraedron gracile</i> (Reinsch) Hansg.	Hydrodictyceae	Planktonic
18	<i>Tetraedron minimum</i> (A. Braun) Hansg.	Hydrodictyceae	Planktonic
19	<i>Tetraedron trigonum</i> (Nägeli) Hansg.	Hydrodictyceae	Planktonic
20	<i>Coelastropsis costata</i> (Korshikov) Fott & Kalina	Scenedesmaceae	Attached
21	<i>Coelastrum microporum</i> Nägeli	Scenedesmaceae	Attached
22	<i>Desmodesmus brasiliensis</i> (Bohlin) E. Hegew.	Scenedesmaceae	Planktonic

23	<i>Desmodesmus pannonicus</i> (Hortob.) E. Hegew.	Scenedesmaceae	Planktonic
24	<i>Desmodesmus hystrix</i> (Lagerh.) E. Hegew.	Scenedesmaceae	Planktonic
25	<i>Scenedesmus armatus</i> (Chodat) Chodat	Scenedesmaceae	Planktonic
26	<i>Scenedesmus ecornis</i> (Ehrenb.) Chodat	Scenedesmaceae	Planktonic
27	<i>Scenedesmus lafevrii</i> var. <i>manguinii</i> M.Lafèvre & Bourr.	Scenedesmaceae	Planktonic
28	<i>Tetradesmus dimorphus</i> (Turpin) M.J. Wynne	Scenedesmaceae	Planktonic
29	<i>Tetradesmus incrassatulus</i> (Bohlin) M.J. Wynne	Scenedesmaceae	Planktonic
30	<i>Tetradesmus lagerheimii</i> M.J. Wynne & Guiry	Scenedesmaceae	Planktonic
31	<i>Tetradesmus obliquus</i> (Turpin) M.J. Wynne	Scenedesmaceae	Planktonic; Attached
32	<i>Kirchneriella lunaris</i> (Kirchn.) K. Möbius	Selenastraceae	Planktonic; Attached
33	<i>Kirchneriella obesa</i> (West) West & G.S. West	Selenastraceae	Planktonic; Attached
34	<i>Monoraphidium convolutum</i> (Corda) Komárk.-Legn.	Selenastraceae	Planktonic
35	<i>Monoraphidium contortum</i> (Thur.) Komárk.-Legn.	Selenastraceae	Planktonic
36	<i>Monoraphidium griffithii</i> (Berk.) Komárk.-Legn.	Selenastraceae	Planktonic
37	<i>Monoraphidium pseudobraunii</i> J.H. Belcher & Swale) Heynig"	Selenastraceae	Planktonic
38	<i>Monoraphidium pusillum</i> (Printz) Komárk.-Legn.	Selenastraceae	Planktonic
39	<i>Ulothrix</i> sp.	Ulotrichaceae	Attached
40	<i>Trentepohlia aurea</i> (L.) C.Mart.	Trentepohliaceae	Attached
41	<i>Trentepohlia willei</i> (Tiffany) Printz	Trentepohliaceae	Attached
42	<i>Cladophora glomerata</i> (L.) Kütz.	Cladophoraceae	Attached
43	<i>Oedogonium varians</i> Wittr. & P.L undell ex Hirn	Oedogoniaceae	Attached
44	<i>Closterium costatum</i> Corda ex Ralfs	Closteriaceae	Planktonic
45	<i>Closterium littorale</i> F. Gay	Closteriaceae	Planktonic; Attached
46	<i>Spirogyra</i> sp.	Zygnemataceae	Attached
<b>Xanthophyceae</b>			
1	<i>Characium angustum</i> A. Braun	Characiaceae	Attached
<b>Euglenophyceae</b>			
1	<i>Cryptoglena skujae</i> Marin & Melkonian	Euglenaceae	Attached
2	<i>Euglena deses</i> Ehrenb.	Euglenaceae	Planktonic
3	<i>Euglena gracilis</i> G.A. Klebs	Euglenaceae	Planktonic
4	<i>Euglena granulata</i> (G.A. Klebs) F. Schmitz	Euglenaceae	Planktonic

5	<i>Euglena sanguinea</i> Ehrenb.	Euglenaceae	Planktonic
6	<i>Euglena tuberculata</i> Swirenko	Euglenaceae	Planktonic
7	<i>Euglena viridis</i> (O.F. Müll.) Ehrenb.	Euglenaceae	Planktonic
8	<i>Euglenaria anabaena</i> (Mainx) Karnkowska & E.W. Linton	Euglenaceae	Planktonic
9	<i>Euglenaria clavata</i> (Skuja) Karnkowska & E.W.	Euglenaceae	Planktonic
10	<i>Eugleniformis proxima</i> (P.A. Dang.) M.S. Benn. & Triemer	Euglenaceae	Planktonic
11	<i>Monomorphina pyrum</i> (Ehrenb.) Mereschk.	Euglenaceae	Attached
12	<i>Strombomonas triquetra</i> (Playfair) Deflandre	Euglenaceae	Planktonic; Attached
13	<i>Trachelomonas abrupt</i> var. minor Deflandre	Euglenaceae	Planktonic
14	<i>Trachelomonas intermedia</i> P.A. Dang.	Euglenaceae	Planktonic
15	<i>Trachelomonas volzii</i> var. intermedia Playfair	Euglenaceae	Planktonic
16	<i>Peranemopsis trichophora</i> (Ehrenb.) L.S. Péter	Peranemidae	Planktonic
17	<i>Lepocinclis acus</i> (O.F. Müll.) Marin & Melkonian	Phacaceae	Planktonic
18	<i>Lepocinclis fusca</i> (Klebs) Kosmala & Zakry	Phacaceae	Planktonic
19	<i>Lepocinclis fusiformis</i> (H.J. Carter) Lemmerm.	Phacaceae	Planktonic
20	<i>Lepocinclis ovum</i> (Ehrenb.) Lemmerm.	Phacaceae	Planktonic
21	<i>Lepocinclis salina</i> F.E. Fritsch	Phacaceae	Planktonic
22	<i>Phacus acuminatus</i> A. Stokes	Phacaceae	Planktonic
23	<i>Phacus caudatus</i> K. Hübner	Phacaceae K. Hübner	Planktonic
24	<i>Phacus curvicauda</i> Svirenko	Phacaceae	Planktonic
25	<i>Phacus glaber</i> Pochm.	Phacaceae	Planktonic
26	<i>Phacus hamatus</i> Pochm.	Phacaceae	Planktonic
27	<i>Phacus helikoides</i> (Ehrenb.) Dujard.	Phacaceae	Planktonic
29	<i>Phacus tortus</i> (Lemmerm.) Skvortzov	Phacaceae	Planktonic
30	<i>Phacus triqueter</i> Perty (Ehrenb.)	Phacaceae	Planktonic
<b>Bacillariophyceae</b>			
1	<i>Aulacoseira</i> sp.	Aulacosieraceae	Planktonic
2	<i>Cyclotella meneghiniana</i> Kütz.	Stephanodiscaceae	Planktonic
3	<i>Diademsis confervacea</i> Kütz.	Diadesmidiaceae	Planktonic; Attached
4	<i>Achnantheidium minutissimum</i> (Kütz.) Czarnecki	Achnanthidiaceae	Attached
5	<i>Anomoeoneis sphaerophora</i> Pfitzer	Anomoeoneidaceae	Attached
6	<i>Cymbella lanceolata</i> (C. Agardh) C. Agardh	Cymbellaceae	Planktonic
8	<i>Gyrosigma acuminatum</i> (Kütz.) Rabenh.	Naviculaceae	Attached

9	<i>Navicula cryptocephala</i> Kütz.	Naviculaceae	Attached
10	<i>Navicula peregrina</i> (Ehrenb.) Kütz.	Naviculaceae	Attached
11	<i>Navicula phyllepta</i> Kütz.	Naviculaceae	Attached
12	<i>Navicula tripunctata</i> (O.F. Müll.) Bory	Naviculaceae	Planktonic; Attached
13	<i>Craticula cuspidate</i> (Kütz.) D.G. Mann	Stauroneidaceae	Attached
14	<i>Craticula halophila</i> (Grunow) D.G. Mann	Stauroneidaceae	Attached
15	<i>Halamphora coffaeiformis</i> (C. Agardh) Mereschk.	Amphipleuraceae	Planktonic
16	<i>Nitzschia acicularis</i> (Kütz.) W.Sm.	Bacillariaceae	Planktonic
17	<i>Nitzschia frustulum</i> (Kütz.) Grunow	Bacillariaceae	Attached
18	<i>Nitzschia fruticose</i> Hust.	Bacillariaceae	Attached
19	<i>Nitzschia palea</i> (Kütz.) W.Sm.	Bacillariaceae	Attached

### 32. List of Bryophytes

Species	Family
<b>Marchantiophyta (Liverworts)</b>	
1. <i>Cyathodium cavernarum</i> Kunze ex Lehm.	Cyathodiaceae
2. <i>Lejeunea alata</i> Gottsche	Lejeuneaceae
3. <i>Lejeunea devendrae</i> (Sushil K. Singh) P.K. Verma & K.K. Rawat	Lejeuneaceae
4. <i>Lopholejeunea sikkimensis</i> Steph.	Lejeuneaceae
5. <i>Riccia billardierei</i> Mont. & Nees	Ricciaceae
6. <i>Riccia discolor</i> Lehm. & Lindenb.	Ricciaceae
7. <i>Spruceanthus minutilobulus</i> (Udar & U.S. Awasthi) Sushil K. Singh	Lejeuneaceae
<b>Bryophyta (Mosses)</b>	
8. <i>Bryum plumosum</i> Dozy & Molk.	Bryaceae
9. <i>Entodontopsis tavoyensis</i> (Hook.) W.R. Buck & Ireland	Stereophyllaceae
10. <i>Erpodium mangiferae</i> Müll. Hal.	Erpodiaceae
11. <i>Fissidens crenulatus</i> Mitt.	Fissidentaceae
12. <i>Fissidens sylvaticus</i> Griff.	Fissidentaceae
13. <i>Hydrogonium arcuatum</i> (Griff.) Wijk & Margad.	Pottiaceae
14. <i>Hydrogonium consanguineum</i> (Thwaites & Mitt.) Hilp.	Pottiaceae
15. <i>Hyophila involuta</i> (Hook.) A. Jaeger	Pottiaceae
16. <i>Hyophila nymaniana</i> (M. Fleisch.) M. Menzel	Pottiaceae

### 33. List of Agro-Flora (Cultivation crops)

S. No.	Local Name	Scientific Name	Family
1.	Bhindi	<i>Abelmoschus esculentus</i> (L.) Moench	Malvaceae
2.	Bel	<i>Aegle marmelos</i> (L.) Corrêa	Rutaceae
3.	Lalsag	<i>Amaranthus cruentus</i> L.	Amaranthaceae
4.	Kanta Note	<i>Amaranthus spinosus</i> L.	Amaranthaceae
5.	Dengua	<i>Amaranthus tricolor</i> L.	Amaranthaceae
6.	Note	<i>Amaranthus viridis</i> L.	Amaranthaceae
7.	OI	<i>Amorphophallus paeoniifolius</i> (Dennst.) Nicolson	Araceae
8.	Supari	<i>Areca catechu</i> L.	Arecaceae



9.	Kathal	<i>Artocarpus heterophyllus</i> Lam.	Moraceae
10.	Bramhi	<i>Bacopa monnieri</i> (L.) Wettst.	Scrophulariaceae
11.	Pui	<i>Basella alba</i> L.	Basellaceae
12.	Taal	<i>Borassus fiabellifer</i> L.	Arecaceae
13.	Phul Kophi	<i>Brassica oleracea</i> var. botrytis L.	Brassicaceae
14.	Lanka	<i>Capsicum annuum</i> L.	Solanaceae
15.	Pepe	<i>Carica papaya</i> L.	Caricaceae
16.	Karamcha	<i>Carissa carandas</i> L.	Apocynaceae
17.	Bethu	<i>Chenopodium album</i> L.	Chenopodiaceae
18.	Lebu	<i>Citrus aurantiifolia</i> (Christm.) Swingle	Rutaceae
19.	Pati Lebu	<i>Citrus limon</i> (L.) Osbeck	Rutaceae
20.	Batabi Lebu	<i>Citrus maxima</i> (Burm.) Merr.	Rutaceae
21.	Kundri	<i>Coccinia grandis</i> (L.) Voigt	Cucurbitaceae
22.	Narkel	<i>Cocos nucifera</i> L.	Arecaceae
23.	Kochu	<i>Colocasia esculenta</i> (L.) Schott	Araceae
24.	Pat	<i>Corchorus olitorius</i> L.	Tiliaceae
25.	Kumro	<i>Cucurbita maxima</i> Duchesne	Cucurbitaceae
26.	Holud	<i>Curcuma longa</i> L.	Zingiberaceae
27.	Aans	<i>Dimocarpus longan</i> Lour.	Sapindaceae
28.	Lombu	<i>Dysoxylum alliaceum</i> (Blume) Blume	Meliaceae
29.	Surjamukhi	<i>Helianthus annuus</i> L.	Asteraceae
30.	Kulekhara	<i>Hygrophila schulli</i> M.R. Almeida & S.M. Almeida	Acanthaceae
31.	Kalmi	<i>Ipomoea aquatica</i> Forssk.	Convolvulaceae
32.	Sim	<i>Lablab purpureus</i> (L.) Sweet	Fabaceae
33.	Lau	<i>Lagenaria siceraria</i> (Molina) Standl.	Cucurbitaceae
34.	Katbel	<i>Limonia acidissima</i> L.	Rutaceae
35.	Tometo	<i>Lycopersicon esculentum</i> Mill.	Solanaceae
36.	Aam	<i>Mangifera indica</i> L.	Anacardiaceae
37.	Safeda	<i>Manilkara zapota</i> (L.) P. Royen	Sapotaceae
38.	Susni	<i>Marsilea minuta</i> L.	Marsileaceae
39.	Uchhe	<i>Momordica charantia</i> L.	Cucurbitaceae
40.	Sojne	<i>Moringa oleifera</i> Lam.	Moringaceae
41.	Kari Pata	<i>Murraya koenigii</i> (L.) Spreng.	Rutaceae
42.	Kola	<i>Musa paradisiaca</i> L.	Musaceae
43.	Shaluk	<i>Nymphaea pubescens</i> Willd.	Nymphaeaceae
44.	Tulsi	<i>Ocimum tenuiflorum</i> L.	Lamiaceae
45.	Dhan	<i>Oryza sativa</i> L.	Poaceae
46.	Khejur	<i>Phoenix sylvestris</i> (L.) Roxb.	Arecaceae
47.	Hari fal	<i>Phyllanthus acidus</i> (L.) Skeels	Euphorbiaceae
48.	Pyara	<i>Psidium guajava</i> L.	Myrtaceae
49.	Bedana	<i>Punica granatum</i> L.	Punicaceae
50.	Mulo	<i>Raphanus sativus</i> L.	Brassicaceae
51.	Til	<i>Sesamum indicum</i> L.	Pedaliaceae
52.	Begun	<i>Solanum melongena</i> L.	Solanaceae
53.	Palak	<i>Spinacia oleracea</i> L.	Amaranthaceae
54.	Aamra	<i>Spondias pinnata</i> (L. f.) Kurz	Anacardiaceae
55.	Jam	<i>Syzygium cumini</i> (L.) Skeels	Myrtaceae
56.	Golap jam	<i>Syzygium samarangense</i> (Blume) Merr. & L.M. Perry	Myrtaceae
57.	Tetul	<i>Tamarindus indica</i> L.	Caesalpiniaceae
58.	Pani fal	<i>Trapa natans</i> L.	Trapaceae
59.	Ghatkanchu	<i>Typhonium flagelliforme</i> (Lodd.) Blume	Araceae
60.	Bhutta	<i>Zea mays</i> L.	Poaceae

### 34. List of Floristic Diversity

S. No.	Name	Family	Habit
1	<i>Ranunculus sceleratus</i> L.	Ranunculaceae	Herb
2	<i>Magnolia champaca</i> (L.) Baill. ex Pierre	Magnoliaceae	Tree
3	<i>Annona reticulata</i> L.	Annonaceae	Tree
4	<i>Polyalthia longifolia</i> (Sonn.) Thwaites	Annonaceae	Tree
5	<i>Cissampelos pareira</i> L.	Menispermaceae	Climber
6	<i>Tinospora cordifolia</i> (Willd.) Miers	Menispermaceae	Climber
7	<i>Stephania japonica</i> (Thunb.)	Menispermaceae	Climber
8	<i>Nymphaea nouchali</i> Burm.f.	Nymphaeaceae	Herb
9	<i>Nymphaea pubescens</i> Willd.	Nymphaeaceae	Herb
10	<i>Nelumbo nucifera</i> Gaertn.	Nelumbonaceae	Herb
11	<i>Argemone mexicana</i> L.	Papaveraceae	Herb
12	<i>Brassica juncea</i> (L.) Czern.	Brassicaceae	Herb
13	<i>Rorippa indica</i> (L.) Hiern	Brassicaceae	Herb
14	<i>Capparis zeylanica</i> L.	Capparaceae	Scandent shrub
15	<i>Cleome gynandra</i> L.	Capparaceae	Herb
16	<i>Cleome rutidosperma</i> DC.	Capparaceae	Herb
17	<i>Cleome viscosa</i> L.	Capparaceae	Herb
18	<i>Portulaca oleracea</i> L.	Portulacaceae	Herb
19	<i>Abelmoschus esculentus</i> (L.) Moench	Malvaceae	Undershrub
20	<i>Abutilon indicum</i> (L.) Sweet	Malvaceae	Undershrub
21	<i>Hibiscus mutabilis</i> L.	Malvaceae	Shrub
22	<i>Hibiscus rosa-sinensis</i> L.	Malvaceae	Shrub
23	<i>Hibiscus vitifolius</i> L.	Malvaceae	Undershrub
24	<i>Malachra capitata</i> (L.) L.	Malvaceae	Undershrub
25	<i>Malvastrum coromandelianum</i> (L.) Garcke	Malvaceae	Undershrub
26	<i>Sida acuta</i> Burm.f.	Malvaceae	Undershrub
27	<i>Sida cordata</i> (Burm.f.) Borss. Waalk.	Malvaceae	Undershrub
28	<i>Sida rhombifolia</i> L.	Malvaceae	Undershrub
29	<i>Urena lobata</i> L.	Malvaceae	Herb
30	<i>Urena sinuata</i> L.	Malvaceae	Herb
31	<i>Ceiba pentandra</i> (L.) Gaertn.	Bombacaceae	Tree
32	<i>Abroma augusta</i> (L.) L.f.	Sterculiaceae	Shrub
33	<i>Melochia corchorifolia</i> L.	Sterculiaceae	Undershrub
34	<i>Corchorus aestuans</i> L.	Tiliaceae	Undershrub
35	<i>Oxalis corniculata</i> L.	Oxalidaceae	Herb
36	<i>Averrhoa bilimbi</i> L.	Averrhoaceae	Tree
37	<i>Aegle marmelos</i> (L.) Corrêa	Rutaceae	Tree
38	<i>Citrus aurantiifolia</i> (Christm.) Swingle	Rutaceae	Tree
39	<i>Citrus limon</i> (L.) Osbeck	Rutaceae	Tree
40	<i>Citrus maxima</i> (Burm.) Merr.	Rutaceae	Tree
41	<i>Glycosmis pentaphylla</i> (Retz.) DC.	Rutaceae	Undershrub
42	<i>Limonia acidissima</i> L.	Rutaceae	Tree
43	<i>Murraya koenigii</i> (L.) Spreng.	Rutaceae	Shrub
44	<i>Murraya paniculata</i> (L.) Jack	Rutaceae	Tree
45	<i>Azadirachta indica</i> A. Juss.	Meliaceae	Tree
46	<i>Melia azedarach</i> L.	Meliaceae	Tree
47	<i>Swietenia mahogani</i> L.	Meliaceae	Tree
48	<i>Ziziphus oenopolia</i> (L.) Mill.	Rhamnaceae	Shrub
49	<i>Cayratia trifolia</i> (L.) Domin	Vitaceae	Climber
50	<i>Cardiospermum halicacabum</i> L.	Sapindaceae	Climber
51	<i>Dimocarpus longan</i> Lour.	Sapindaceae	Tree

52	<i>Anacardium occidentale</i> L.	Anacardiaceae	Tree
53	<i>Mangifera indica</i> L.	Anacardiaceae	Tree
54	<i>Spondias pinnata</i> (L. f.) Kurz	Anacardiaceae	Tree
55	<i>Moringa oleifera</i> Lam.	Moringaceae	Tree
56	<i>Bauhinia acuminata</i> L.	Caesalpiniaceae	Tree
57	<i>Caesalpinia bonduc</i> (L.) Roxb. (Roxb.) K. Larsen & S.S. Larsen	Caesalpiniaceae	Shrub
58	<i>Cassia fistula</i> L.	Caesalpiniaceae	Tree
59	<i>Cassia javanica</i> subsp. <i>nodosa</i>	Caesalpiniaceae	Tree
60	<i>Delonix regia</i> (Bojer ex Hook.) Raf.	Caesalpiniaceae	Tree
61	<i>Peltophorum pterocarpum</i> (DC.) Baker ex K. Heyne	Caesalpiniaceae	Tree
62	<i>Senna alata</i> (L.) Roxb.	Caesalpiniaceae	Shrub
63	<i>Senna obtusifolia</i> (L.) Irwin & Barneby	Caesalpiniaceae	Undershrub
64	<i>Senna occidentalis</i> (L.) Link	Caesalpiniaceae	Undershrub
65	<i>Senna siamea</i> (Lam.) H.S. Irwin & Barneby	Caesalpiniaceae	Tree
66	<i>Senna tora</i> (L.) Roxb.	Caesalpiniaceae	Undershrub
67	<i>Tamarindus indica</i> L.	Caesalpiniaceae	Tree
68	<i>Acacia auriculiformis</i> A. Cunn. ex Benth.	Mimosaceae	Tree
69	<i>Acacia nilotica</i> (L.) Willd. ex Delile ssp. <i>indica</i> (Benth.) Brenan	Mimosaceae	Tree
70	<i>Albizia lebbek</i> (L.) Benth.	Mimosaceae	Tree
71	<i>Albizia saman</i> (Jacq.) Merr.	Mimosaceae	Tree
72	<i>Lysiloma latisiliquum</i> (L.) Benth.	Mimosaceae	Tree
73	<i>Neptunia oleracea</i> Lour.	Mimosaceae	Herb
74	<i>Pithecellobium dulce</i> (Roxb.) Benth.	Mimosaceae	Tree
75	<i>Aeschynomene aspera</i> L.	Fabaceae	Herb
76	<i>Aeschynomene indica</i> L.	Fabaceae	Herb
77	<i>Alysicarpus vaginalis</i> (L.) DC.	Fabaceae	Herb
78	<i>Butea monosperma</i> (Lam.) Taub.	Fabaceae	Tree
79	<i>Dalbergia latifolia</i> Roxb.	Fabaceae	Tree
80	<i>Desmodium gangeticum</i> (L.) DC.	Fabaceae	Herb
81	<i>Desmodium triflorum</i> (L.) DC.	Fabaceae	Herb
82	<i>Erythrina variegata</i> L.	Fabaceae	Tree
83	<i>Sesbania sesban</i> (L.) Merr.	Fabaceae	Shrub
84	<i>Smithia sensitiva</i> Aiton	Fabaceae	Herb
85	<i>Teramnus labialis</i> (L.f.) Spreng.	Fabaceae	Climber
86	<i>Myriophyllum tetrandrum</i> Roxb.	Haloragaceae	Herb
87	<i>Combretum indicum</i> (L.) DeFilippis	Combretaceae	Climber
88	<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight & Arn.	Combretaceae	Tree
89	<i>Terminalia catappa</i> L.	Combretaceae	Tree
90	<i>Corymbia citriodora</i> (Hook.) K.D. Hill & L.A.S. Johnson	Myrtaceae	Tree
91	<i>Callistemon lanceolatus</i> (Sm.) Sweet	Myrtaceae	Tree
92	<i>Psidium guajava</i> L.	Myrtaceae	Tree
93	<i>Syzygium cumini</i> (L.) Skeels	Myrtaceae	Tree
94	<i>Syzygium samarangense</i> (Blume) Merr. & L.M. Perry	Myrtaceae	Tree
95	<i>Ammannia baccifera</i> L.	Lythraceae	Herb
96	<i>Lagerstroemia speciosa</i> (L.) Pers.	Lythraceae	Tree
97	<i>Lawsonia inermis</i> L.	Lythraceae	Shrub
98	<i>Punica granatum</i> L.	Punicaceae	Shrub
99	<i>Ludwigia adscendens</i> (L.) H. Hara	Onagraceae	Herb
100	<i>Ludwigia hyssopifolia</i> (G. Don) Exell	Onagraceae	Herb

101	<i>Ludwigia perennis</i> L.	Onagraceae	Herb
102	<i>Trapa natans</i> L.	Trapaceae	Herb
103	<i>Passiflora foetida</i> L.	PassiLoraceae	Climber
104	<i>Passiflora suberosa</i> L.	PassiLoraceae	Climber
105	<i>Carica papaya</i> L.	Caricaceae	Tree
106	<i>Coccinia grandis</i> (L.) Voigt	Cucurbitaceae	Climber
107	<i>Cucurbita coccinea</i> G. Don	Cucurbitaceae	Climber
108	<i>Momordica charantia</i> L.	Cucurbitaceae	Climber
109	<i>Glinus lotoides</i> L.	Aizoaceae	Herb
110	<i>Glinus oppositifolius</i> (L.) Aug. DC.	Aizoaceae	Herb
111	<i>Trianthema portulacastrum</i> L.	Aizoaceae	Herb
112	<i>Centella asiatica</i> (L.) Urb.	Apiaceae	Herb
113	<i>Foeniculum vulgare</i> Mill.	Apiaceae	Herb
114	<i>Oenanthe javanica</i> (Blume) DC.	Apiaceae	Herb
115	<i>Alangium salvifolium</i> (L.f.) Wangerin	Alangiaceae	Tree
116	<i>Dentella repens</i> (L.) J.R. Forst. & G. Forst.	Rubiaceae	Herb
117	<i>Ixora chinensis</i> Lam.	Rubiaceae	Shrub
118	<i>Morinda angustifolia</i> Roxb.	Rubiaceae	Shrub
119	<i>Mussaenda philippica</i> A. Rich.	Rubiaceae	Shrub
120	<i>Neolamarckia cadamba</i> (Roxb.) Bosser	Rubiaceae	Tree
121	<i>Oldenlandia biflora</i> L.	Rubiaceae	Herb
122	<i>Oldenlandia corymbosa</i> L.	Rubiaceae	Herb
123	<i>Paederia foetida</i> L.	Rubiaceae	Climber
124	<i>Ageratum conyzoides</i> (L.) L.	Asteraceae	Herb
125	<i>Blumea lacera</i> (Burm.f.) DC.	Asteraceae	Herb
126	<i>Caesulia axillaris</i> Roxb.	Asteraceae	Herb
127	<i>Cyanthillium cinereum</i> (L.) H. Rob.	Asteraceae	Herb
128	<i>Eclipta prostrata</i> (L.) L.	Asteraceae	Herb
129	<i>Eleutheranthera ruderalis</i> (Sw.) Sch. Bip.	Asteraceae	Herb
130	<i>Enydra fluctuans</i> DC.	Asteraceae	Herb
131	<i>Grangea maderaspatana</i> (L.) Poir.	Asteraceae	Herb
132	<i>Helianthus annuus</i> L.	Asteraceae	Undershrub
133	<i>Laphangium luteoalbum</i> (L.) Tzvelev	Asteraceae	Herb
134	<i>Mikania micrantha</i> Kunth	Asteraceae	Climber
135	<i>Parthenium hysterophorus</i> L.	Asteraceae	Herb
136	<i>Pluchea indica</i> (L.) Less.	Asteraceae	Herb
137	<i>Pseudelephantopus spicatus</i> (B. Juss. ex Aubl.) Rohr ex C.F. Baker	Asteraceae	Herb
138	<i>Sonchus arvensis</i> L.	Asteraceae	Herb
139	<i>Tridax procumbens</i> (L.) L.	Asteraceae	Herb
140	<i>Xanthium strumarium</i> L.	Asteraceae	Undershrub
141	<i>Synedrella nodiflora</i> (L.) Gaertn.	Asteraceae	Herb
142	<i>Sphagneticola calendulacea</i> (L.) Pruski	Asteraceae	Herb
143	<i>Sphenoclea zeylanica</i> Gaertn.	Sphenocleaceae	Herb
144	<i>Manilkara zapota</i> (L.) P. Royen	Sapotaceae	Tree
145	<i>Nyctanthes arbor-tristis</i> L.	Oleaceae	Tree
146	<i>Allamanda cathartica</i> L.	Apocynaceae	Climber
147	<i>Alstonia scholaris</i> (L.) R. Br.	Apocynaceae	Tree
148	<i>Cascabela thevetia</i> (L.) Lippold	Apocynaceae	Tree
149	<i>Carissa carandas</i> L.	Apocynaceae	Tree
150	<i>Catharanthus roseus</i> (L.) G. Don	Apocynaceae	Herb
151	<i>Kopsia fruticosa</i> (Roxb.) A. DC.	Apocynaceae	Tree
152	<i>Nerium oleander</i> L.	Apocynaceae	Shrub
153	<i>Rauvolfia tetraphylla</i> L.	Apocynaceae	Undershrub

154	<i>Tabernaemontana divaricata</i> (L.) R. Br. ex Roem. & Schult.	Apocynaceae	Shrub
155	<i>Calotropis gigantea</i> (L.) R. Br.	Asclepiadaceae	Shrub
156	<i>Pergularia daemia</i> (Forssk.) Chiov.	Asclepiadaceae	Climber
157	<i>Dregea volubilis</i> (L.f.) Benth. ex Hook.f.	Asclepiadaceae	Climber
158	<i>Nymphoides cristata</i> (Roxb.) Kuntze	Menyanthaceae	Herb
159	<i>Nymphoides indica</i> (L.) Kuntze	Menyanthaceae	Herb
160	<i>Heliotropium curassavicum</i> L.	Boraginaceae	Herb
161	<i>Heliotropium indicum</i> L.	Boraginaceae	Herb
162	<i>Heliotropium ovalifolium</i> Forssk.	Boraginaceae	Herb
163	<i>Cuscuta campestris</i> Yunck.	Convolvulaceae	Climber
164	<i>Evolvulus alsinoides</i> (L.) L.	Convolvulaceae	Herb
165	<i>Evolvulus nummularius</i> (L.) L.	Convolvulaceae	Herb
166	<i>Hewittia malabarica</i> (L.) Suresh	Convolvulaceae	Climber
167	<i>Ipomoea aquatica</i> Forssk.	Convolvulaceae	Herb
168	<i>Ipomoea carnea</i> Jacq. subsp. <i>fistulosa</i> (Mart. ex Choisy) D.F. Austin	Convolvulaceae	Undershrub
169	<i>Ipomoea obscura</i> (L.) Ker Gawl.	Convolvulaceae	Climber
170	<i>Ipomoea marginata</i> (Desr.) Verdc.	Convolvulaceae	Climber
171	<i>Ipomoea triloba</i> L.	Convolvulaceae	Climber
172	<i>Operculina turpethum</i> (L.) Silva Manso	Convolvulaceae	Climber
173	<i>Capsicum annum</i> L.	Solanaceae	Herb
174	<i>Cestrum diurnum</i> L.	Solanaceae	Shrub
175	<i>Datura metel</i> L.	Solanaceae	Herb
176	<i>Lycopersicon esculentum</i> Mill.	Solanaceae	Herb
177	<i>Nicotiana plumbaginifolia</i> Viv.	Solanaceae	Herb
178	<i>Physalis minima</i> L.	Solanaceae	Herb
179	<i>Solanum americanum</i> Mill.	Solanaceae	Herb
180	<i>Solanum indicum</i> L.	Solanaceae	Herb
181	<i>Solanum melongena</i> L.	Solanaceae	Herb
182	<i>Solanum sisymbriifolium</i> Lam.	Solanaceae	Herb
183	<i>Solanum torvum</i> Sw.	Solanaceae	Herb
184	<i>Bacopa monnieri</i> (L.) Wettst.	Scrophulariaceae	Herb
185	<i>Limnophila heterophylla</i> (Roxb.) Benth.	Scrophulariaceae	Herb
186	<i>Lindernia oppositifolia</i> (Retz.) Mukherjee	Scrophulariaceae	Herb
187	<i>Mazus pumilus</i> (Burm.f.) Steenis	Scrophulariaceae	Herb
188	<i>Mecardonia procumbens</i> (Mill.) Small	Scrophulariaceae	Herb
189	<i>Scoparia dulcis</i> L.	Scrophulariaceae	Herb
190	<i>Utricularia aurea</i> Lour.	Lentibulariaceae	Herb
191	<i>Utricularia stellaris</i> L.f.	Lentibulariaceae	Herb
192	<i>Tabebuia aurea</i> (Silva Manso) Benth. & Hook.f. ex S. Moore	Bignoniaceae	Tree
193	<i>Spathodea campanulata</i> P. Beauv.	Bignoniaceae	Tree
194	<i>Sesamum indicum</i> L.	Pedaliaceae	Herb
195	<i>Acanthus ilicifolius</i> L.	Acanthaceae	Undershrub
196	<i>Andrographis paniculate</i> (Burm. f.) Wall. ex Nees	Acanthaceae	Herb
197	<i>Asystasia gangetica</i> (L.) T. Anderson	Acanthaceae	Herb
198	<i>Barleria prionitis</i> L.	Acanthaceae	Undershrub
199	<i>Hemigraphis hirta</i> (Vahl) T. Anderson	Acanthaceae	Herb
200	<i>Hygrophila schulli</i> M.R. Almeida & S.M. Almeida	Acanthaceae	Herb
201	<i>Hygrophila triflora</i> (Roxb. ex Nees) Fosberg & Sachet	Acanthaceae	Herb
202	<i>Justicia adhatoda</i> L.	Acanthaceae	Undershrub

203	<i>Justicia gendarussa</i> Burm.f.	Acanthaceae	Undershrub
204	<i>Justicia simplex</i> D. Don	Acanthaceae	Herb
205	<i>Ruellia prostrata</i> Poir.	Acanthaceae	Herb
206	<i>Ruellia tuberosa</i> L.	Acanthaceae	Herb
207	<i>Clerodendrum indicum</i> (L.) Kuntze	Verbenaceae	Undershrub
208	<i>Clerodendrum infortunatum</i> L.	Verbenaceae	Undershrub
209	<i>Duranta erecta</i> L.	Verbenaceae	Shrub
210	<i>Gmelina arborea</i> Roxb.	Verbenaceae	Tree
211	<i>Lantana camara</i> L.	Verbenaceae	Shrub
212	<i>Lippia alba</i> (Mill.) N.E. Br. ex Britton & P. Wilson	Verbenaceae	Undershrub
213	<i>Phyla nodiflora</i> (L.) Greene	Verbenaceae	Herb
214	<i>Tectona grandis</i> L.f.	Verbenaceae	Tree
215	<i>Vitex negundo</i> L.	Verbenaceae	Shrub
216	<i>Volkameria inermis</i> L.	Verbenaceae	Shrub
217	<i>Anisomeles indica</i> (L.) Kuntze	Lamiaceae	Undershrub
218	<i>Leonurus sibiricus</i> L.	Lamiaceae	Undershrub
219	<i>Leucas aspera</i> (Willd.) Link	Lamiaceae	Herb
220	<i>Hyptis suaveolens</i> (L.) Poit	Lamiaceae	Herb
221	<i>Ocimum tenuiflorum</i> L.	Lamiaceae	Herb
222	<i>Boerhavia diffusa</i> L.	Lamiaceae	Herb
223	<i>Boerhavia repens</i> L.	Lamiaceae	Herb
224	<i>Mirabilis jalapa</i> L.	Lamiaceae	Herb
225	<i>Achyranthes aspera</i> L.	Amaranthaceae	Herb
226	<i>Aerva lanata</i> (L.) Juss.	Amaranthaceae	Herb
227	<i>Alternanthera ficoidea</i> (L.) Sm.	Amaranthaceae	Herb
228	<i>Alternanthera paronychioides</i> A.St.-Hil.	Amaranthaceae	Herb
229	<i>Alternanthera philoxeroides</i> (Mart.) Griseb.	Amaranthaceae	Herb
230	<i>Alternanthera sessilis</i> (L.) R.Br. ex DC.	Amaranthaceae	Herb
231	<i>Amaranthus cruentus</i> L.	Amaranthaceae	Herb
232	<i>Amaranthus hybridus</i> L.	Amaranthaceae	Herb
233	<i>Amaranthus spinosus</i> L.	Amaranthaceae	Herb
234	<i>Amaranthus tricolor</i> L.	Amaranthaceae	Herb
235	<i>Amaranthus viridis</i> L.	Amaranthaceae	Herb
236	<i>Celosia argentea</i> L.	Amaranthaceae	Herb
237	<i>Gomphrena globosa</i> L.	Amaranthaceae	Herb
238	<i>Chenopodium album</i> L.	Chenopodiaceae	Herb
239	<i>Suaeda maritima</i> (L.) Dumort.	Chenopodiaceae	Herb
240	<i>Basella alba</i> L.	Basellaceae	Climber
241	<i>Persicaria barbata</i> (L.) H. Hara	Polygonaceae	Herb
242	<i>Persicaria glabra</i> (Willd.) M. Gómez	Polygonaceae	Herb
243	<i>Persicaria hydropiper</i> (L.) Delarbre	Polygonaceae	Herb
244	<i>Persicaria orientalis</i> (L.) Spach	Polygonaceae	Herb
245	<i>Polygonum plebeium</i> R. Br.	Polygonaceae	Herb
246	<i>Rumex dentatus</i> L.	Polygonaceae	Herb
247	<i>Peperomia pellucida</i> (L.) Kunth	Piperaceae	Herb
248	<i>Litsea glutinosa</i> (Lour.) C.B. Rob.	Lauraceae	Tree
249	<i>Acalypha indica</i> L.	Euphorbiaceae	Herb
250	<i>Chrozophora rotleri</i> (Geiseler)	Euphorbiaceae	Undershrub
251	<i>Codiaeum variegatum</i> (L.) Rumph. ex A. Juss.	Euphorbiaceae	Shrub
252	<i>Croton bonplandianus</i> Baill.	Euphorbiaceae	Undershrub
253	<i>Euphorbia antiquorum</i> L.	Euphorbiaceae	Shrub
254	<i>Euphorbia heterophylla</i> L.	Euphorbiaceae	Undershrub
255	<i>Euphorbia hirta</i> L.	Euphorbiaceae	Herb
256	<i>Euphorbia indica</i> Lam.	Euphorbiaceae	Herb

257	<i>Euphorbia nivulia</i> Buch. -Ham.	Euphorbiaceae	Shrub
258	<i>Euphorbia prostrata</i> Aiton	Euphorbiaceae	Herb
259	<i>Euphorbia thymifolia</i> L.	Euphorbiaceae	Herb
260	<i>Euphorbia tithymaloides</i> L.	Euphorbiaceae	Undershrub
261	<i>Excoecaria agallocha</i> L.	Euphorbiaceae	Tree
262	<i>Flueggea virosa</i> (Roxb. ex Willd.) Voigt	Euphorbiaceae	Shrub
263	<i>Jatropha curcas</i> L.	Euphorbiaceae	Shrub
264	<i>Jatropha gossypifolia</i> L.	Euphorbiaceae	Shrub
265	<i>Trewia nudiflora</i> L.	Euphorbiaceae	Tree
266	<i>Putranjiva roxburghii</i> Wall.	Euphorbiaceae	Tree
267	<i>Phyllanthus acidus</i> (L.) Skeels	Euphorbiaceae	Tree
268	<i>Phyllanthus amarus</i> Schumach. & Thonn.	Euphorbiaceae	Herb
269	<i>Phyllanthus emblica</i> L.	Euphorbiaceae	Tree
270	<i>Phyllanthus reticulatus</i> Poir.	Euphorbiaceae	Shrub
271	<i>Phyllanthus urinaria</i> L.	Euphorbiaceae	Herb
272	<i>Ricinus communis</i> L.	Euphorbiaceae	Shrub
273	<i>Pilea microphylla</i> (L.) Liebm.	Urticaceae	Herb
274	<i>Pouzolzia zeylanica</i> (L.) Benn.	Urticaceae	Herb
275	<i>Cannabis sativa</i> L.	Cannabaceae	Shrub
276	<i>Trema orientalis</i> (L.) Blume	Cannabaceae	Tree
277	<i>Artocarpus heterophyllus</i> Lam.	Moraceae	Tree
278	<i>Ficus benghalensis</i> L.	Moraceae	Tree
279	<i>Ficus hispida</i> L.f.	Moraceae	Tree
280	<i>Ficus racemosa</i> L.	Moraceae	Tree
281	<i>Ficus religiosa</i> L.	Moraceae	Tree
282	<i>Streblus asper</i> Lour.	Moraceae	Shrub
283	<i>Casuarina equisetifolia</i> L.	Casuarinaceae	Tree
284	<i>Ceratophyllum demersum</i> L.	Ceratophyllaceae	Herb
285	<i>Hydrilla verticillata</i> (L.f.) Royle	Hydrocharitaceae	Herb
286	<i>Nechamandra alternifolia</i> (Roxb. ex Wight) Thwaites	Hydrocharitaceae	Herb
287	<i>Ottelia alismoides</i> (L.) Pers.	Hydrocharitaceae	Herb
288	<i>Vallisneria spiralis</i> L.	Hydrocharitaceae	Herb
289	<i>Vanda tessellata</i> (Roxb.) Hook. ex G. Don	Orchidaceae	Herb
290	<i>Curcuma longa</i> L.	Zingiberaceae	Herb
291	<i>Musa paradisiaca</i> L.	Musaceae	Herb
292	<i>Canna indica</i> L.	Cannaceae	Herb
293	<i>Dioscorea bulbifera</i> L.	Dioscoreaceae	Climber
294	<i>Eichhornia crassipes</i> (Mart.) Solms	Pontederiaceae	Herb
295	<i>Monochoria hastata</i> (L.) Solms	Pontederiaceae	Herb
296	<i>Monochoria vaginalis</i> (Burm.f.) C. Presl	Pontederiaceae	Herb
297	<i>Commelina benghalensis</i> L.	Commelinaceae	Herb
298	<i>Commelina diffusa</i> Burm.f.	Commelinaceae	Herb
299	<i>Commelina longifolia</i> Lam.	Commelinaceae	Herb
300	<i>Commelina paludosa</i> Blume	Commelinaceae	Herb
301	<i>Murdannia vaginata</i> (L.) G. Brückn.	Commelinaceae	Herb
302	<i>Cyanotis axillaris</i> (L.) D. Don ex Sweet	Commelinaceae	Herb
303	<i>Areca catechu</i> L.	Arecaceae	Tree
304	<i>Borassus flabellifer</i> L.	Arecaceae	Tree
305	<i>Cocos nucifera</i> L.	Arecaceae	Tree
306	<i>Dypsis lutescens</i> (H. Wendl.) Beentje & J. Dransf.	Arecaceae	Shrub
307	<i>Phoenix sylvestris</i> (L.) Roxb.	Arecaceae	Tree
308	<i>Typha angustifolia</i> L.	Typhaceae	Herb
309	<i>Typha domingensis</i> Pers.	Typhaceae	Herb

310	<i>Alocasia macrorrhizos</i> (L.) G. Don	Araceae	Herb
311	<i>Colocasia esculenta</i> (L.) Schott	Araceae	Herb
312	<i>Cryptocoryne spiralis</i> (Retz.) Fisch. ex Wydler	Araceae	Herb
313	<i>Pistia stratiotes</i> L.	Araceae	Herb
314	<i>Typhonium flagelliforme</i> (Lodd.) Blume	Araceae	Herb
315	<i>Lemna minor</i> L.	Lemnaceae	Herb
316	<i>Lemna perpusilla</i> Torrey	Lemnaceae	Herb
317	<i>Spirodela polyrrhiza</i> (L.) Schleid.	Lemnaceae	Herb
318	<i>Wolffia globosa</i> (Roxb.) Hartog & Plas	Lemnaceae	Herb
319	<i>Sagittaria guayanensis</i> Kunth	Alismataceae	Herb
320	<i>Sagittaria montevidensis</i> Cham. & Schtdl.	Alismataceae	Herb
321	<i>Sagittaria sagittifolia</i> L.	Alismataceae	Herb
322	<i>Najas graminea</i> Delile	Najadaceae	Herb
323	<i>Aponogeton natans</i> (L.) Engl. & K. Krause	Aponogetonaceae	Herb
324	<i>Potamogeton crispus</i> L.	Potamogetonaceae	Herb
325	<i>Bolboschoenus maritimus</i> subsp. <i>afinis</i> (Roth.) T. Koyama	Cyperceae	Herb
326	<i>Bulbostylis densa</i> (Wall.) Hand. -Mazz.	Cyperceae	Herb
327	<i>Cyperus alopecuroides</i> Rottb.	Cyperceae	Herb
328	<i>Cyperus articulatus</i> L.	Cyperceae	Herb
329	<i>Cyperus compressus</i> L.	Cyperceae	Herb
330	<i>Cyperus corymbosus</i> Rottb.	Cyperceae	Herb
331	<i>Cyperus difformis</i> L.	Cyperceae	Herb
332	<i>Cyperus distans</i> L.f.	Cyperceae	Herb
333	<i>Cyperus kyllingia</i> Endl.	Cyperceae	Herb
334	<i>Cyperus nutans</i> Vahl	Cyperceae	Herb
335	<i>Cyperus pangorei</i> Rottb.	Cyperceae	Herb
336	<i>Cyperus pilosus</i> Vahl	Cyperceae	Herb
337	<i>Cyperus polystachyos</i> Rottb.	Cyperceae	Herb
338	<i>Cyperus pumilus</i> L.	Cyperceae	Herb
339	<i>Cyperus rotundus</i> L.	Cyperceae	Herb
340	<i>Eleocharis spiralis</i> (Rottb.) Roem. & Schult.	Cyperceae	Herb
341	<i>Fimbristylis eragrostis</i> (Nees) Hance	Cyperceae	Herb
342	<i>Fimbristylis polytrichoides</i> (Retz.) Vahl	Cyperceae	Herb
343	<i>Fimbristylis schoenoides</i> (Retz.) Vahl	Cyperceae	Herb
344	<i>Kyllinga brevifolia</i> Rottb.	Cyperceae	Herb
345	<i>Mariscus aristatus</i> (Rottb.) T. Tang & F.T. Wang	Cyperceae	Herb
346	<i>Pycnus polystachyos</i> (Rottb.) P. Beauv.	Cyperceae	Herb
347	<i>Schoenoplectiella articulata</i> (L.) Lye	Cyperceae	Herb
348	<i>Brachiaria reptans</i> (L.) C.A. Gardner & C.E. Hubb.	Poaceae	Herb
349	<i>Chloris barbata</i> Sw.	Poaceae	Herb
350	<i>Chrysopogon aciculatus</i> (Retz.) Trin.	Poaceae	Herb
351	<i>Coix aquatica</i> Roxb.	Poaceae	Herb
352	<i>Cynodon dactylon</i> (L.) Pers.	Poaceae	Herb
353	<i>Dactyloctenium aegyptium</i> (L.) Willd.	Poaceae	Herb
354	<i>Echinochloa colona</i> (L.) Link	Poaceae	Herb
355	<i>Echinochloa crus-galli</i> (L.) P.Beauv.	Poaceae	Herb
356	<i>Eleusine indica</i> (L.) Gaertn.	Poaceae	Herb
357	<i>Eragrostis amabilis</i> (L.) Wight & Arn.	Poaceae	Herb
358	<i>Hygroryza aristata</i> (Retz.) Nees ex Wight & Arn.	Poaceae	Herb
359	<i>Imperata cylindrica</i> (L.) Rausch.	Poaceae	Herb
360	<i>Isachne globosa</i> (Thunb.) Kuntze	Poaceae	Herb



361	<i>Leersia hexandra</i> Sw.	Poaceae	Herb
362	<i>Leptochloa chinensis</i> (L.) Nees	Poaceae	Herb
363	<i>Panicum paludosum</i> Roxb.	Poaceae	Herb
364	<i>Paspalidium flavidum</i> (Retz.) A. Camus	Poaceae	Herb
365	<i>Paspalidium punctatum</i> (Burm.) A. Camus	Poaceae	Herb
366	<i>Paspalum conjugatum</i> P.J. Bergius	Poaceae	Herb
367	<i>Paspalum scrobiculatum</i> L.	Poaceae	Herb
368	<i>Phragmites karka</i> (Retz.) Trin. ex Steud.	Poaceae	Herb
369	<i>Phyllostachys aurea</i> Rivière & C. Rivière	Poaceae	Herb
370	<i>Vetiveria zizanioides</i> (L.) Nash	Poaceae	Herb
371	<i>Zoysia matrella</i> (L.) Merr.	Poaceae	Herb
372	<i>Azolla imbricate</i> (Roxb. ex Griff.) Nakai	Salviniaceae	Herb
373	<i>Salvinia molesta</i> D.S. Mitch.	Salviniaceae	Herb
374	<i>Salvinia natans</i> (L.) All.	Salviniaceae	Herb
375	<i>Marsilea minuta</i> L.	Marsileaceae	Herb
376	<i>Ceratopteris thalictroides</i> (L.) Brongn.	Pteridaceae	Herb
377	<i>Adiantum</i> sp.	Pteridaceae	Herb
378	<i>Pteris vittata</i> L.	Pteridaceae	Herb
379	<i>Asplenium nidus</i> L.	Aspleniaceae	Herb
380	<i>Drynaria quercifolia</i> (L.) J. Sm.	Polypodiaceae	Herb
381	<i>Platycladus orientalis</i> (L.) Franco	Cupressaceae	Herb

### 35. List of Invasive Alien Species (Vascular Plants)

S. No.	Plant Name	Family	Common Name	Habitat	Nativity
1	<i>Ageratum conyzoides</i> (L.) L.	Asteraceae	Goat weed	Herb	Tropical America
2	<i>Alternanthera paronychioides</i> A.St.-Hil.	Amaranthaceae	Smooth joy weed	Herb	Tropical America
3	<i>Alternanthera philoxeroides</i> (Mart.) Griseb.	Amaranthaceae	Alligator weed	Herb	Tropical America
4	<i>Argemone mexicana</i> L.	Papaveraceae	Prickly poppy	Herb	Tropical Central & South America
5	<i>Blumea lacera</i> (Burm.f.) DC.	Asteraceae	Lettuce-leaf Blumea	Herb	Tropical America
6	<i>Calotropis gigantea</i> (L.) R. Br.	Asclepiadaceae	Crown flower	Shrub	Tropical Africa
7	<i>Celosia argentea</i> L.	Amaranthaceae	Wool flower	Herb	Tropical Africa
8	<i>Chloris barbata</i> Sw.	Poaceae	Peacock plume grass	Herb	Tropical America
9	<i>Chrozophora rottleri</i> (Geiseler) A. Juss. ex Spreng.	Euphorbiaceae	Suryavarti	Under shrub	Tropical Africa
10	<i>Cleome gynandra</i> L.	Capparaceae	Wild Spider flower	Herb	Tropical America

11	<i>Cleome rutidosperma</i> DC.	Capparaceae	Fringed spider flower	Herb	Tropical America
12	<i>Cleome viscosa</i> L.	Capparaceae	Dog mustard	Herb	Tropical America
13	<i>Corchorus aestuans</i> L.	Tiliaceae	East Indian mallow	Under shrub	Tropical America
14	<i>Croton bonplandianus</i> Baill.	Euphorbiaceae	Bonpland's Croton	Under shrub	Temperate South America
15	<i>Cyperus difformis</i> L.	Cyperaceae	Rice sedge	Herb	Tropical America
16	<i>Datura metel</i> L.	Solanaceae	Thorn apple	Herb	Tropical America
17	<i>Echinochloa colona</i> (L.) Link	Poaceae	Shama millet	Herb	Tropical South America
18	<i>Echinochloa crus-galli</i> (L.) P. Beauv.	Poaceae	Barnyard millet	Herb	Tropical South America
19	<i>Eclipta prostrata</i> (L.) L.	Asteraceae	False daisy	Herb	Tropical America
20	<i>Eichhornia crassipes</i> (Mart.) Solms*	Pontederiaceae	Water hyacinth	Herb	Tropical America
21	<i>Euphorbia heterophylla</i> L.	Euphorbiaceae	Wild Poinsettia	Under shrub	Tropical America
22	<i>Euphorbia hirta</i> L.	Euphorbiaceae	Snake weed	Herb	Tropical America
23	<i>Euphorbia indica</i> Lam.	Euphorbiaceae	Snake weed	Herb	Tropical South America
24	<i>Evolvulus nummularius</i> (L.) L.	Convolvulaceae	Round leaf bind weed	Herb	Tropical America
25	<i>Grangea maderaspatana</i> (L.) Poir.	Asteraceae	Madras carpet	Herb	Tropical South America
26	<i>Hyptis suaveolens</i> (L.) Poit	Lamiaceae	American mint	Herb	Tropical America
27	<i>Imperata cylindrica</i> (L.) Raeusch.*	Poaceae	Cogon grass	Herb	Tropical America
28	<i>Ipomoea carnea</i> Jacq. subsp. <i>fistulosa</i> (Mart. ex Choisy)	Convolvulaceae	Bush morning glory	Shrub	Tropical America
29	<i>Ipomoea obscura</i> (L.) Ker Gawl.	Convolvulaceae	Obscure morning glory	Climber	Tropical Africa
30	<i>Lantana camara</i> L.*	Verbenaceae	Wild sage	Shrub	Tropical America
31	<i>Ludwigia adscendens</i> (L.) H. Hara	Onagraceae	Water primrose	Herb	Tropical America
32	<i>Ludwigia perennis</i> L.	Onagraceae	Paddy clove	Herb	Tropical Africa
33	<i>Lysiloma latisiliquum</i> (L.) Benth. *	Mimosaceae	Horse tamarind	Tree	Tropical America
34	<i>Malachra capitata</i> (L.) L.	Malvaceae	Brazil jute	Under shrub	Tropical America
35	<i>Malvastrum coromandelianum</i> (L.) Garcke"	Malvaceae	False mallow	Under shrub	Tropical America
36	<i>Mecardonia procumbens</i> (Mill.) Small	Scrophulariaceae	Baby jump-up	Herb	Tropical North America
37	<i>Melochia corchorifolia</i> L.	Sterculiaceae	Chocolate weed	Under shrub	Tropical America
38	<i>Mikania micrantha</i> Kunth*	Asteraceae	Mile-a-minute	Climber	Tropical America

39	<i>Monochoria vaginalis</i> (Burm.f.) C. Presl	Pontederiaceae	Oval leaf pond weed	Herb	Tropical America
40	<i>Nicotiana plumbaginifolia</i> Viv.	Solanaceae	Tex-mex tobacco	Herb	Tropical America
41	<i>Oxalis corniculata</i> L.	Oxalidaceae	Indian sorrel	Herb	Europe
42	<i>Parthenium hysterophorus</i> L.	Asteraceae	Congress weed	Herb	Tropical North America
43	<i>Passiflora foetida</i> L.	Passifloraceae	Stinking passion flower	Climber	Tropical South America
44	<i>Peperomia pellucida</i> (L.) Kunth	Piperaceae	Rat's ear	Herb	Tropical South America
45	<i>Pilea microphylla</i> (L.) Liebm.	Urticaceae	Gun powder plant	Herb	Tropical South America
46	<i>Pistia stratiotes</i> L.	Araceae	Water lettuce	Herb	Tropical America
47	<i>Portulaca oleracea</i> L.	Portulacaceae	Purslane	Herb	Tropical South America
48	<i>Ruellia tuberosa</i> L.	Acanthaceae	Menow weed	Herb	Tropical America
49	<i>Salvinia molesta</i> D.S. Mitch. *	Salviniaceae	Kariba weed	Herb	Brazil
50	<i>Scoparia dulcis</i> L.	Scrophulariaceae	Sweet broom weed	Herb	Tropical America
51	<i>Senna alata</i> (L.) Roxb.	Caesalpiniaceae	Roman candle	Shrub	West Indies
52	<i>Senna obtusifolia</i> (L.) Irwin & Barneby	Caesalpiniaceae	Sickle senna	Under shrub	Tropical America
53	<i>Senna occidentalis</i> (L.) Link	Caesalpiniaceae	Coffee weed	Under shrub	Tropical South America
54	<i>Sida acuta</i> Burm.f.	Malvaceae	Morning mallow	Under shrub	Tropical America
55	<i>Solanum americanum</i> Mill.	Solanaceae	Nightshade	Herb	Tropical America
56	<i>Solanum torvum</i> Sw.	Solanaceae	Turkey berry	Under shrub	West Indies
57	<i>Synedrella nodiflora</i> (L.) Gaertn.	Asteraceae	Node weed	Herb	West Indies
58	<i>Tridax procumbens</i> (L.) L.	Asteraceae	Mexican daisy	Herb	Tropical Central America
59	<i>Typha domingensis</i> Pers. ( <i>T. angustata</i> Bory.)	Typhaceae	Cat tail	Herb	Tropical America
60	<i>Urena lobata</i> L.	Malvaceae	Bur mallow	Herb	Tropical Africa
61	<i>Xanthium strumarium</i> L.	Asteraceae	Cocklebur	Under shrub	Tropical America

\* - Species included in the list of 100 of the world's worst invasive alien species (Lowe et. al., 2000).

## AI Wetland Mitra

The concept of 'Wetland Mitra' is an informal, voluntary and non-statutory network of concerned citizens to foster and promote community engagement in wetlands conservation and management efforts. An effective Wetland Mitra' network is poised to enable wetland management to gain access to local views, rights and capacities for supporting wetlands management. The network is also aimed as a communication and outreach vehicle for promoting awareness on the value of wetlands, and management and conservation efforts. By involving themselves within the Wetland Mitra network, citizens gain an opportunity of shaping wetlands management by bringing onboard indigenous land local knowledge, and views of diverse stakeholder groups.

Key role and responsibilities of Wetland Mitra are as follows:

- a. Promote awareness on the relevance, values and functions of the wetlands with local communities, students, resident welfare groups and other stakeholders.
- b. Build a team of Wetland Mitras and engage in its restoration, rejuvenation & management
- c. Participate in wetlands management planning and implementation processes
- d. Alert authorities on any activity detrimental to wetlands

The following four members were enrolled in our Wetland Mitra Network of the East Kolkata Wetlands so far:

1. Mr. Altup Sekh, Headmaster, Kheadaha High School (HS).
2. Mr. Manas Haldar, Headmaster, Bamanghata High School.
3. Mr. Sukumar Mandal, Assistant Headmaster, Kheadaha High School (HS).
4. Dr. Subir K. Ghosh, Expert/Researcher.

The following duly filled Form is to be submitted to the East Kolkata Wetlands Management Authority to become a Wetland Mitra of the East Kolkata Wetlands:

**BE a Wetland Mitra**

**Together we can rejuvenate, conserve our wetlands**

**Name:** \_\_\_\_\_ **Gender : Male/ Female**

**Occupation:** \_\_\_\_\_

**Contact Details:**

**Phone:** \_\_\_\_\_

**Email:** \_\_\_\_\_

**Address:** \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

I am willing to commit \_\_\_\_\_ hours per month to work towards restoring/  
rejuvenating the health of our wetland East Kolkata Wetlands

**My areas of interest:**

**Wetland Assessment/ surveys**

**Awareness/ Mobilisation**

**Bird surveys**

**Monitoring**

**Clean up drives**


**Others (specify)**

\_\_\_\_\_

\_\_\_\_\_  
**Signature**

## A6 Schedule of Rates

### I. Boundary Pillar

<b>WEST BENGAL ZOO AUTHORITY</b> <i>(A Registered Society under the Department of Forests, Government of West Bengal)</i>	
Ref: 476/WBZA/ D-18 /15-16	Date : 25 / 01 /2016
Chief Technical Officer, EKWMA, Poura Bhavan, 4th Floor, Block - FD-415/A, Sector III Bidhannagar, Kolkata - 700106.	
Sub. : Demarcation of EKWMA Boundary - vetting of estimates.	
Ref. : a) Your letter vide no. CTO/EN//030/348/2012-13/16 dated 25.01.2016 b) Interaction held between concerned Executives of West Bengal Zoo Authority and EKWMA.	
Please be informed that the details behind derivation of rates for each pillar i.e. (i) Rs.1800/- for construction and erection (ii) Rs.800/- for surveying land periphery have been observed by this department and further interacted with concerned persons at your office. The ground reality and working practice in respect of respective construction and execution plan of above mentioned works were discussed at length and considered for derivation.	
The above rates based on prevalent PWD, schedule of works have been derived as hereunder and has been duly vetted by Chief Engineer this department.	
a) Rs.1553/- each pillar for constructions & erection as enclosed vide Annexure 1/ (i) & (ii).	
b) Rs.1225/- each pillar for surveying the land periphery as enclosed vide Annexure 2.	
All above are for your information and necessary course of action.	
 Member Secretary West Bengal Zoo Authority	
2 <sup>nd</sup> Floor, KB – 19, Sector – III, Salt Lake City, Kolkata – 700 098, West Bengal Phone: +91 33 23355010 (Direct), 23355011 (EPABX), FAX: +91 33 23350012 E-mail: <a href="mailto:mswbza@gmail.com">mswbza@gmail.com</a> , <a href="mailto:wbzooauthority@yahoo.in">wbzooauthority@yahoo.in</a>	
sd	

Estimate for Construction of Boundary Pillar						
Sl. No.	Description	Unit	Quantity	Rate Rs.	Amount Rs.	Total Amount(Rs.)
1.00 P-1 2.	Earth work in excavation of foundation trenches or drains in all sorts of soil (including mixed soil but excluding laterite or sandstone) including removing, spreading or stacking the spils within a lead of 75m. As directed. The item includes necessary trimming the sides of trenches, leveling dressing and ramming the bottom, bailing out water as required complete. a) Depth of excavation not exceeding 1500 mm. Column 1 Nos. x 0.600 x 0.600 x 0.750 = 0.270 Cum 0.270 Cum	Cum	0.270	120.47	32.53	
2.00 P-1 3.	Earth work in filling in foundation trenches or plinth with good earth in layers not exceeding 150 mm. including water and raming etc. layer by layer complete. (Payment to be made on basis of measurement of finished quantity of work). a) With earth obtained from excavation of foundation. 1/3 x 0.270 = 0.090 Cum	Cum	0.090	78.31	7.05	
3.00 P-11 1.	Single brick flat soling of picked jhama bricks including ramming and dressing bed to proper level and filling joints with powdered earth or local sand. 1 Nos. x 0.375 x 0.500 = 0.188 sqm 0.188 sqm	Sqm	0.188	377.00	70.69	
4.00 P-11 5	Ordinary Cement Concrete (mix 1:2:4) with graded stone chips (20 mm down) excluding shuttering and reinforcement if any, in ground floor. As per IS456-2000 a) Pakur Variety Base 1 Nos. x 0.450 x 0.250 x 0.150 = 0.017 Cum Pedestal 1 Nos. x 0.250 x 0.250 x 1.050 = 0.066 Cum 0.083 Cum	Cum	0.083	6064.02	500.28	
5.00 P-26 12.	Hire and labour charges for shuttering with centreing and necessary staging upto 4 m. using approved stout props and thick hard wood planks of approved thickness with required bracing for concrete slabs, beams, eplumms, lintels curved or straight including fitting, fixing and striking out after completion of works. (upto roof of ground floor) a) 25 mm. to 30 mm. thick wooden shuttering as per decision & direction of Engineer-In-Chargr Base 1 Nos. x 1.400 x 0.150 = 0.210 sqm Pedestal 1 Nos. x 1.000 x 1.050 = 1.050 sqm 1.260 sqm	Sqm	1.260	360.00	453.60	
6.00 P-27 15.	Reinforcement for reinforced concrete work in all sorts of structures i/c. distribution bars, stirrups, binders etc. including supply of rods, straightening and removal of loose rust (if necessary) cutting to requisite length, hooking and bending to correct shape, placing in proper position and binding with 16 gauge black annealed wire at every inter-section, complete as per drawing and direction (For all floors) 1.0% of 0.083-Cum = 0.006 MT	MT	0.006	68508.00	443.67	
					Total -	1507.82
7.00	Add for provisional contingency @3%					45.23
					Grand Total -	1,553.05
					Or say	1,553.00


(Rupees One Thousand Five Hundred Fifty Three Only)

Note : The rate based on P.W.D. Schedule of Rate 2014

The above estimate of Rs. 1553.05 is correct.

Chief Engineer  
West Bengal Zoo Authority

Member Secretary  
West Bengal Zoo Authority



*Hyamal R.*  
Chief Engineer  
West Bengal Zoo Authority

3  
Member Secretary  
West Bengal Zoo Authority



Estimate for Survey Of Land Perifery						
Sl. No.	Description	Unit	Quantity	Rate Rs.	Amount Rs.	Total Amount(Rs.)
1.00	Cost of doing works, in relation to survey works, by the way of using prismatic compass, measuring distance between station to station with chain, cutting bushes, making visibility between the stations etc.					
	a) Labour for showing ranging rod and helping placing prismatic compass properly.	each	1 Nos.	245.00	245.00	
	b) Labour for measuring distances between stations	each	2 Nos.	245.00	490.00	
	c) Labour for cutting bushes, clearing obstacles etc.	each	2 Nos.	245.00	490.00	
					<b>Total -</b>	<b>1225.00</b>

( Rupees One Thosuang Two Hundred Twenty five Only )

Note : The rate based on **F.W.D. Schedule of Rate 2014**

*Hymal R*  
 Chief Engineer  
 West Bengal Zoo Authority

*B*  
 Member Secretary  
 West Bengal Zoo Authority

## 2. Dredging and desilting

### **ABSTRACT OF COST FOR THE WORK**

Abstract of cost for the work Re- excavation & De-silting of Lalkuthi-2 from 0.00 m to 3000.00 m in connection with improvement of sewage Distribution system at East Kolkata Wetlands area.

Item No.	Description of Items	Quantity	Unit	Rate (Rs.)	Amount (Rs.)
1.	Bailing out water by pump including all leads and lifts and including making all arrangements for disposal, where continuous flow or seepage of water is encountered and rating of pumps and hours of operation is specifically ascertained as per direction of the Engineer-in-charge.  (ref-vide item no.4.01, page no. C-35 of Unified Schedule of Rates )	700.00	HP-Hr	23.00	16100.00
2.	Earthwork in excavation in the canals / drainage channels / borrow pits / River bed / slope of bank as noted below as per designed section & throwing the spoils in layers of 25 cm. for making banks as per profile or to deposit the same in any other place within an initial lead of 30 metre and initial lift of 1.5 metre including breaking clods, rough dressing etc. complete as per direction of the Engineer-in-charge.  (a) Ordinary soil / local soil / soil mixed with moorum or kankar / silt, dry or moist / slushy silt / sticky soil.  (ref-vide item no.1.02, page no. C-1 of Unified Schedule of Rates )	12074.00	Cum	109.00	1316066.00
3.	Earthwork from borrow pits over existing embankment or for repairs to embankment, closing breaches etc. within an initial lead of 30 metre and lift of 1.50 metre including breaking clods and depositing the same in layers not exceeding 25 cm. and rough dressing etc. complete as per profile.  (ref-vide item no.1.01, page no. C-1 of Unified Schedule of Rates )	30.12	Cum	81.00	2439.72
4.	Extra rate over Item Nos.2 & 3 for all kinds of soil for every additional lead of 30 metres or part thereof beyond the initial lead of 30 metres.  (ref-vide item no.1.04, page no. C-1 of Unified Schedule of Rates ) (a) One additional lead	5500.00	Cum	1 x 12.00	66000.00
5.	Extra rate over Item Nos.2 & 3 for all kinds of soil for every additional lift of 1.50 metre or part thereof beyond the initial lift of 1.50 metre.  (ref-vide item no.1.05, page no. C-1 of Unified Schedule of Rates ) (a) one additional lift (b) Two additional lift	6500.00 5200.00	Cum Cum	1 x 12.00 2 x 12.00	78000.00 124800.00

Item No.	Description of Items	Quantity	Unit	Rate (Rs.)	Amount (Rs.)
6.	Supplying to site empty cement polythene bags (capacity 50kg) in good condition including stacking, incidental charges etc. all complete (capacity 50 kg).  (a) Polythene bags  (ref-vide item no.2.24, page no. C-11 of Unified Schedule of Rates )	1100	Each	6.00	6600.00
7.	Labour charges for filling empty cement gunny bag / polythene bag (capacity 50 kg) with all kinds of earth (wet or dry), sand or aggregates(stone / brick) and sewing as done in case of cement bags ,( tying with a knot after filling will not be accepted ) , and stacking within a lead of 30 metre as per direction of Engr-in-charge (cost is excluding the cost of any types of filling materials). NB:- Volume of earth/ sand filled bags should not be less than 0.028 cum/ bag.  (ref-vide item no.2.27, page no. C-12 of Unified Schedule of Rates )	1100	Each	7.00	7700.00
8.	Dumping in position earth / sand filled gunny / poly bags (capacity 50 kg) within a lead of 30 metre inclusive of all lifts.  (ref-vide item no.2.29, page no. C-12 of Unified Schedule of Rates )	1100	Each	3.00	3300.00
9.	Removing earth/sand filled poly bags/gunny bags from the channel bed/slope and initially depositing the same near channel bank at a place as directed by the Engineer-in Charge for measurement of Bag. & then disposing off the same at a distance not less than 150 metre from the channel site.  (ref-vide item no.2.32, page no. C-12 of Unified Schedule of Rates )	810	Each	6.00	4860.00
10.	Supplying, dressing and stacking at site bamboo pins above 5 cm. diameter and upto 7.5 cm. diameter. (Dia. to be measured at mid length). (ref-vide item no.2.09, page no. C-09 of Unified Schedule of Rates )	1200.00	Meter	29.00	34800.00
11.	Driving bamboo pins in standing or flowing water including scaffolding and driving appliances. (a) Where boats are not necessary.  (ref-vide item no.2.09, page no. C-09 of Unified Schedule of Rates )	900.00	Meter	11.00	9900.00
12.	Supplying, fitting and fixing with iron nails half split bamboo wailing pieces (av. dia. of bamboo not less than 5.0 cm.)  (ref-vide item no.2.11, page no. C-09 of Unified Schedule of Rates )	400.00	Meter	22.00	8800.00
13.	Taking out old bamboo pins from piling and stacking the same within 150 metre from the site of work.  (a) Where boats are not necessary.  (ref-vide item no.2.12, page no. C-10 of Unified Schedule of Rates )	900.00	Meter	8.00	7200.00

<b>Item No.</b>	<b>Description of Items</b>	<b>Quantity</b>	<b>Unit</b>	<b>Rate (Rs.)</b>	<b>Amount (Rs.)</b>
14.	Survey Cost (Cost for survey works towards pre-work and post-work section and tabulation of data in the section paper including charges for surveyors, skilled and skilled labours, bamboo pegs rope etc. and all other incidental charges completed as per direction. Survey instruments will have to supply by the agency. Cross section is to be taken at an interval not more than 60 meter apart).	3.0	Km	9000.00	27000.00
	<b>Total</b>				<b>17,13,565.72</b>
	<b>Contingency @5%</b>				<b>85,678.29</b>
	<b>Grand Total</b>				<b>17,99,244.01</b>
	<b>Rounded Off</b>				<b>17,99,244.00</b>
	(Rupees: Seventeen lakhs ninety nine thousand two hundred forty four only)				



Save Wetlands to Save Life



Department of Environment  
Government of West Bengal